# Generalized quantum asymptotic equipartition

Kun Fang  $*1$ , Hamza Fawzi  $*2$ , and Omar Fawzi  $*3$ 

<sup>1</sup> School of Data Science, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, China <sup>2</sup>Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, United Kingdom 3 Inria, ENS de Lyon, UCBL, LIP, 69342, Lyon Cedex 07, France

#### Abstract

The asymptotic equipartition property (AEP) states that in the limit of a large number of independent and identically distributed (i.i.d.) random experiments, the output sequence is virtually certain to come from the typical set, each member of which is almost equally likely. This property is a form of the law of large numbers and lies at the heart of information theory.

In this work, we prove a generalized quantum AEP beyond the i.i.d. framework where the random samples are drawn from two sets of quantum states. In particular, under suitable assumptions on the sets, we prove that all operationally relevant divergences converge to the quantum relative entropy between the sets. More specifically, both the quantum hypothesis testing relative entropy (a smoothed form of the min-relative entropy) and the smoothed max-relative entropy approach the regularized relative entropy between the sets. Notably, the asymptotic limit has explicit convergence guarantees and can be efficiently estimated through convex optimization programs, despite the regularization, provided that the sets have efficient descriptions.

As the first application, the generalized AEP directly implies a new generalized quantum Stein's lemma for conducting quantum hypothesis testing between two sets of quantum states. This addresses open questions raised by Brandão et al. [IEEE TIT 66(8):5037–5054, 2020] and Mosonyi et al. [IEEE TIT 68(2):1032-1067, (2022)], which seek a Stein's lemma with computational efficiency. Moreover, we propose a new framework for quantum resource theory in which state transformations are performed without requiring precise characterization of the states being manipulated, making it more robust to imperfections. We demonstrate the reversibility (also referred to as the second law) of such a theory and identify the regularized relative entropy as the "unique" measure of the resource in this new framework.

As the second application, we introduce a quantum version of adversarial hypothesis testing, which was first studied in the classical case by Brandão et al. [ITCS 2014], where the tester plays against an adversary who possesses internal quantum memory and controls the quantum device. We show that the optimal error exponent is precisely characterized by a new notion of quantum channel divergence, named the minimum output channel divergence. Interestingly, by proving the chain rule property of the minimum output channel divergence, we show that the adaptive strategies employed by the adversary provide no advantage over non-adaptive ones.

As the third application, we derive a relative entropy accumulation theorem stating that the smoothed min-relative entropy between two sequential processes of quantum channels can be lower bounded by the sum of the regularized minimum output channel divergences. A special case of this theorem gives a version of the  $H_{\text{max}}$  entropy accumulation theorem established in Dupuis et al. [CMP 379(3):867-913, 2020] and Metger et al. [FOCS 2022].

As the fourth application, we apply our generalized AEP to quantum resource theories and provide improved and efficient bounds for entanglement distillation, magic state distillation, and the entanglement cost of quantum states and channels. These serve as illustrative examples to demonstrate the general idea to leverage the computational efficiency of our results when the problem of interest is hard to solve directly.

<sup>\*</sup>kunfang@cuhk.edu.cn

<sup>†</sup> h.fawzi@damtp.cam.ac.uk

<sup>‡</sup> omar.fawzi@ens-lyon.fr

Finally, we believe the technical tools established in this work — including the variational formula, superadditivity for the measured relative entropy between two sets of states, the chain rule property for the quantum relative entropy, and the minimum output channel divergence — will find more applications in quantum information. In particular, we expect that the measured relative entropy, which is at the heart of all the proofs in this work, will play a more prominent role in quantum information.

# **Contents**



## <span id="page-2-0"></span>1 Introduction

The asymptotic equipartition property (AEP) is a form of the law of large numbers in information theory, stating that  $-\frac{1}{n}$  $\frac{1}{n} \log p(X_1, X_2, \dots, X_n)$  is close to the entropy  $H(X)$ , where  $X_1, X_2, \dots, X_n$  are independent, identically distributed (i.i.d.) random variables and  $p(X_1, X_2, \dots, X_n)$  is the probability of observing the sequence  $X_1, X_2, \cdots, X_n$  [\[Cov99\]](#page-55-0). The AEP was first stated by Shannon in his seminal 1948 paper [\[Sha48\]](#page-58-0), where he proved the result for i.i.d. processes and stated the result for staionary ergodic processes. McMillan [\[McM53\]](#page-57-0) and Breiman [\[Bre57\]](#page-55-1) then proved the AEP for ergodic finite alphabet sources. The result is now referred to as the AEP or the Shannon–McMillan–Breiman theorem. This property lies in the heart of information theory, forming the mathematical foundation for applications such as data compression  $<sup>1</sup>$  $<sup>1</sup>$  $<sup>1</sup>$ , channel coding [\[Sha48\]](#page-58-0) and cryptography [\[HV93,](#page-56-0) [VV95,](#page-58-1) [Cac97\]](#page-55-2).</sup>

The AEP can be represented in a more generic form in terms of entropic quantities:

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} \frac{1}{n} \mathbb{D}_{\varepsilon}(P^{\otimes n} \| Q^{\otimes n}) = D(P \| Q),\tag{1}
$$

where P is a probability distribution, Q is a nonnegative function on a finite set and  $\mathbb D$  is a divergence of interest,  $\varepsilon$  is a smoothing parameter and D is the Kullback–Leibler divergence also called the relative entropy. For instance, considering  $Q$  to be a constant function equal to 1 and  $D$  be the min-relative entropy and max-relative entropy, then the above equation reduce to the Shannon–McMillan–Breiman theorem.

The AEP has been generalized to quantum information theory, by replacing the probability distribution with the density matrix of a quantum state and using quantum divergences,

$$
\lim_{\varepsilon \to 0} \lim_{n \to \infty} \frac{1}{n} \mathbb{D}_{\varepsilon}(\rho^{\otimes n} \| \sigma^{\otimes n}) = D(\rho \| \sigma), \tag{2}
$$

where  $\rho$  is the density matrix,  $\sigma$  is a positive semidefinite operator. In particular, Hiai and Petz [\[HP91\]](#page-56-1) proved the case when  $\mathbb{D}_{\varepsilon}$  is the quantum hypothesis testing relative entropy and D is the Umegaki relative entropy. Ogawa and Nagaoka [\[NO00\]](#page-57-1) then strenghtened the result by removing the dependence of  $\varepsilon$  in the outer limit. These together formed the well-known quantum Stein's lemma in quan-tum information theory. Tomamichel et al. [\[TCR09\]](#page-58-2) proved the case when  $\rho_{AB}$  is a bipartite quantum state and  $\sigma_{AB} = I_A \otimes \rho_B$  and  $\mathbb{D}_{\varepsilon}$  is the quantum min-relative entropy and quantum max-relative entropy [\[Ren05\]](#page-58-3). As the classical AEP plays an essential role in classical information theory, the quantum AEP has also found plenty of applications, including quantum data compression [\[Sch95\]](#page-58-4), quantum state merging [\[Ber09\]](#page-54-0), quantum channel coding [\[DMHB13\]](#page-55-3), quantum cryptography [\[Ren05\]](#page-58-3) and quantum resource theory [\[FWTD19\]](#page-56-2).

For many problems, the situation is not so simple and  $\rho^{\otimes n}$  and  $\sigma^{\otimes n}$  are not fully known and do not necessarily possess the i.i.d. structure. All one knows is that they belong to two sets,  $\mathscr{A}_n$  and  $\mathscr{B}_n$ , respectively, making the existing quantum AEPs not applicable. Such as scenario has been extensively studied in the classical information theory and applied to practical problems (refer to [\[LM02\]](#page-57-2) and references therein) such as classification with training sequences (e.g. speech recognition, signal detection and digital communication) and detection of messages via unknown channels (e.g. radar target detection, identification problem and watermark detection). This motivates us to explore a more general notion of the quantum AEP by extending beyond the i.i.d. structure of the source and beyond the singleton case. Specifically, we consider the following limit:

<span id="page-2-2"></span>
$$
\lim_{n \to \infty} \frac{1}{n} \mathbb{D}_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = ? \quad \text{with} \quad \mathbb{D}_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) := \inf_{\substack{\rho_n \in \mathscr{A}_n \\ \sigma_n \in \mathscr{B}_n}} \mathbb{D}_{\varepsilon}(\rho_n \| \sigma_n)
$$
\n(3)

where  $\mathscr{A}_n$  is a set of quantum states and  $\mathscr{B}_n$  is a set of positive semidefinite operators acting on  $\mathcal{H}^{\otimes n}$ , where  $H$  is a finite-dimensional Hilbert space. This is a very general framework that encompasses almost

<span id="page-2-1"></span><sup>1</sup> The AEP has been proved to be almost equivalent to the source coding theorem by [\[VH97\]](#page-58-5), which reinforces the prominent role played by the AEP in information theory.

all existing studies of quantum AEP in the literature. For instance, if  $\mathbb{D}_{\varepsilon}$  is the hypothesis testing relative entropy, then  $\mathcal{A}_n$  is seen as a composite null hypothesis and  $\mathcal{B}_n$  as a composite alternate hypothesis. Specific examples that were recently studied include composite hypothesis testing where  $\mathscr{A}_n$  and  $\mathscr{B}_n$  are convex mixtures of i.i.d. states [\[BBH21\]](#page-54-1), as well as the generalized quantum Stein's lemma, where  $\mathcal{A}_n$ is an i.i.d. state and  $\mathcal{B}_n$  is a set of quantum states [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3).

#### <span id="page-3-0"></span>1.1 Main contributions

Under some assumptions on the sets  $\mathscr{A}_n$  and  $\mathscr{B}_n$ , in particular the stability of the sets and their polars under tensor products, we prove that the limit in Eq.  $(3)$  exists for both the hypothesis testing relative entropy  $D_{H,\varepsilon}$  (a smoothed form of the min-relative entropy) and the max-relative entropy  $D_{\max,\varepsilon}$ , which represent two extreme cases in the family of quantum divergences. Both limits converge to the quantum relative entropy between the sets. More specifically, we establish the following:

<span id="page-3-1"></span>**Theorem 1.** *(Generalized AEP, informal) Let*  $\{\mathcal{A}_n\}_{n\in\mathbb{N}}$  *and*  $\{\mathcal{B}_n\}_{n\in\mathbb{N}}$  *be two sequences of sets of states such that each set is convex, compact, permutation invariant, and the sequences as well as their polar sets are closed under tensor product. Then for any*  $\varepsilon \in (0,1)$ *, it holds* 

$$
nD^{\infty}(\mathscr{A}||\mathscr{B}) - O(n^{2/3}\log n) \le D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \le nD^{\infty}(\mathscr{A}||\mathscr{B}) + O(n^{2/3}\log n),\tag{4}
$$

$$
nD^{\infty}(\mathscr{A}||\mathscr{B}) - O(n^{2/3}\log n) \le D_{\max,\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \le nD^{\infty}(\mathscr{A}||\mathscr{B}) + O(n^{2/3}\log n),\tag{5}
$$

where  $D^{\infty}(\mathscr{A}|\mathscr{B}) := \lim_{n\to\infty} \frac{1}{n}D(\mathscr{A}_n|\mathscr{B}_n)$ , which can be estimated with explicit convergence guar*antees and can be efficiently computed given that*  $\mathscr{A}_m$  *and*  $\mathscr{B}_m$  *have efficient descriptions.* 

A central ingredient to the proof of this result is the superadditivity of the measured relative entropy between two sets of quantum states under the stability assumption of the polar set,

$$
D_{\mathrm{M}}(\mathscr{A}_{m+k}||\mathscr{B}_{m+k}) \ge D_{\mathrm{M}}(\mathscr{A}_{m}||\mathscr{B}_{m}) + D_{\mathrm{M}}(\mathscr{A}_{k}||\mathscr{B}_{k}). \tag{6}
$$

Some remarks concerning the statement are in order:

- (*Generality.*) All assumptions, except for the one regarding polar sets, are standard in most existing literature (e.g., [\[BP10,](#page-55-4) [BHLP20,](#page-54-2) [HY24,](#page-56-3) [Lam24\]](#page-57-3)). The polar assumption, while technical, is satisfied in many cases of interest. For instance, it holds when the set is a singleton of i.i.d. state, as in the existing quantum AEP; when the set consists of the identity operator tensored with all density operators, which is relevant for conditional quantum entropies; or when it is the image of a quantum channel or a set of quantum channels, a scenario that naturally arises in quantum channel discrimination.
- (*Relaxation.*) Even in cases where the polar assumption is not directly satisfied, we can relax the original problem to meet the required conditions. This approach provides an efficient approximation to the original problem, which may be difficult to solve directly. We illustrate this approach in the main text through applications in entanglement manipulation and magic state distillation and anticipate that this approach has the potential for far-reaching applications beyond the specific cases discussed here.
- (*Efficiency.*) The leading order in the generalized AEP is characterized by a regularized quantity, which will be shown to be necessary in general through specific examples. However, the asymptotic limit comes with explicit convergence guarantees and can still be efficiently estimated via convex optimization programs, despite the regularization, provided that  $\mathscr{A}_n$  and  $\mathscr{B}_n$  have efficient descriptions.
- (*Explicit finite n estimate.*) Our generalized AEP has a leading term that is independent of  $\varepsilon$ , corresponding to the strong converse property in information theory. Moreover, while the leading order is characterized by a regularized quantity, we can still provide an explicit estimate for finite

 $n$ , making its convergence controllable in the finite block length regime. Even though the bound we obtain on the second order term is  $O(n^{2/3} \log n)$  instead of the usual  $O(\sqrt{n})$ , this is a rare case where we can explicitly bound the convergence rate for a regularized quantity.

• (*Divergences.*) The results apply to the two extreme cases of quantum divergence; therefore, any divergence that lies between the hypothesis testing relative entropy and the max-relative entropy, or is equivalent to these, will yield a similar result.

On top of the generalized AEP, we provide several applications, including fundamental tasks such as quantum state discrimination and quantum channel discrimination with the corresponding Stein's lemma, a relative entropy accumulation theorem with potential applications in quantum cryptography, and the quantum resource theory of entanglement and magic, which are important for quantum networking and fault-tolerant quantum computing.

Application 1: quantum hypothesis testing between two sets of states. Quantum hypothesis testing is a pivotal concept in quantum information theory, forming the foundation for numerous applications, including quantum channel coding, quantum illumination, and the operational interpretation of quantum divergences. A central task in hypothesis testing is to discriminate between quantum states corresponding to the different hypotheses, with the goal of determining which state is actually provided. In this work, we investigate quantum hypothesis testing between two sets of quantum states,  $\mathscr{A}_n$  and  $\mathscr{B}_n$ . The fundamental difficulty in analyzing such a hypothesis testing is that both hypotheses are composite and extend beyond the i.i.d. structure. We need to control the type-I error  $\alpha(\mathscr{A}_n, M_n)$  within a constant threshold whenever the sample is drawn from  $\mathcal{A}_n$ , in order to prevent the extremely bad case from passing the test. The asymmetric setting then seeks to determine the optimal exponent at which the type-II error probability decays  $\beta(\mathcal{B}_n, M_n)$ , known as the *Stein's exponent*, while keeping the type-I error within a fixed threshold  $\varepsilon$ . Specifically, the goal is to evaluate

$$
\beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) := \inf_{0 \le M_n \le I} \left\{ \beta(\mathscr{B}_n, M_n) : \alpha(\mathscr{A}_n, M_n) \le \varepsilon \right\}.
$$
 (7)

<span id="page-4-0"></span>Theorem 2. *(Another generalized quantum Stein's lemma, informal.) Given the same assumptions on*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  as stated in Theorem [1.](#page-3-1) Then for any  $\varepsilon \in (0,1)$ , it holds

$$
\lim_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = D^{\infty}(\mathscr{A} \| \mathscr{B}).
$$
\n(8)

This result extends the renowned quantum Stein's lemma to a more general setting. Notably, by utilizing the generalized AEP in Theorem [26,](#page-16-0) the rate of convergence in our version of Stein's lemma can be explicitly determined, and the Stein's exponent can be efficiently estimated via convex optimization programs, despite the regularization involved, provided that  $\mathcal{A}_n$  and  $\mathcal{B}_n$  have efficient descriptions. This addresses open questions raised of significant interest in the community, particularly those seeking a Stein's lemma with efficient estimation, as discussed in [\[BHLP20,](#page-54-2) Section 3.5] and [\[MSW22,](#page-57-4) Section VI].

The discussion on the reduction to the classical case and the comparison with different variants of the quantum Stein's lemma is presented as follows:

- (*Reduction to the classical case.*) If  $\{\mathcal{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathcal{B}_n\}_{n\in\mathbb{N}}$  are classical probability distributions, then the measured relative entropy coincides with the quantum relative entropy. In this case, we can show that  $D(\mathscr{A}_n|\mathscr{B}_n)$  is additive. Therefore, the Stein's exponent  $D^{\infty}(\mathscr{A}|\mathscr{B})$  can be replaced by the one-shot quantity  $D(\mathscr{A}_1|\mathscr{B}_1)$  without regularization.
- (*Quantum Stein's lemma*.) Let  $\mathscr{A}_n = \{\rho^{\otimes n}\}\$  and  $\mathscr{B}_n = \{\sigma^{\otimes n}\}\$  be two singletons. Then it is easy to check that these sets satisfy all the required assumptions. Therefore, our generalized quantum Stein's lemma covers the quantum Stein's lemma [\[HP91,](#page-56-1) [NO00\]](#page-57-1) as a special case.
- (*Generalized quantum Stein's lemma.*) Theorem [2](#page-4-0) is incomparable to the generalized quantum Stein lemma of [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3). It is weaker in the sense that we have the additional assumption on the stability of the polar sets under tensor product for the alternate hypothesis  $\mathcal{B}_n$ , but it is stronger in two ways: it allows for a composite null hypothesis  $\mathscr{A}_n$  whereas the results of [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3) are restricted to  $\mathscr{A}_n = \{\rho^{\otimes n}\}\$ , and in addition we can obtain efficient and controlled approximations of the Stein exponent as presented in Theorem [1.](#page-3-1)
- (*Composite quantum Stein's lemma.*) The composite quantum hypothesis testing studied in [\[BBH21,](#page-54-1) [MSW22\]](#page-57-4) does not satisfy the stability assumption of the sets. Therefore, it is not comparable to our result or the previous generalized quantum Stein's lemma [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3).
- (*Quantum Stein's lemma for restricted measurements.*) In [\[BHLP20,](#page-54-2) Theorem 16], the authors proved a quantum Stein's lemma for restricted measurements, assuming that the measurements and the set of quantum states are compatible [\[Pia09\]](#page-58-6). In this work, we do not impose constraints on the performed measurements. As a result, the measurements and the set of quantum states in our result may not be compatible in their context, making our result not directly comparable to theirs. It is also worth noting that our Stein's lemma possesses the strong converse property (i.e., its convergence is independent of  $\varepsilon$ ), whereas the Stein's lemma for restricted measurements [\[BHLP20,](#page-54-2) Theorem 16] only exhibits a weak converse property (i.e., it requires taking  $\varepsilon \to 0$ ).

As the existing generalized quantum Stein's lemma is deeply connected to the second law of quantum resource theory, we propose a new framework for quantum resource theory in which state transformations are performed without requiring precise characterization of the states being manipulated, making it more robust to imperfections. We demonstrate that such a theory is reversible under asymptotically resource non-generating operations, identifying the regularized quantum relative entropy between two sets as the "unique" measure of resource in this new framework.

Application 2: adversarial quantum channel discrimination. Consider a scenario in which a tester is working with an untrusted quantum device that generates a quantum state upon request. This device guarantees that the state is produced by either the quantum channel  $N$  (the bad case) or the quantum channel  $M$  (the good case). The tester is permitted to request multiple samples from the device and perform measurements to determine which channel is being used. However, because the device is untrusted, it may have access to the environmental system of the channel and also possess an internal memory that correlates with the generated samples, actively misleading the tester and undermining their ability to correctly identify the channel. The key question here is: *How effectively can the tester distinguish between the good and bad cases while playing against the adversary?* Such a setting has been studied by [\[BHLP20\]](#page-54-2) in the classical case. In this work, we investigate the fully quantum setting.

Let  $\mathcal{A}_n$  and  $\mathcal{B}_n$  be the sets of quantum states generated by any adaptive strategies from the adversary. Similar to quantum state discrimination, we denote  $\beta_{\varepsilon}(\mathscr{A}_n||\mathscr{B}_n)$  as the optimal type-II error probability when the type-I error is within a constant threshold  $\varepsilon$ . We will show that the optimal exponent at which  $\beta_{\varepsilon}(\mathscr{A}_n||\mathscr{B}_n)$  decays is given by a new notion of quantum channel divergence, which we call the *minimum output channel divergence*, defined as

$$
D^{\inf}(\mathcal{N}||\mathcal{M}) := \inf_{\substack{\rho \in \mathcal{D}(A) \\ \sigma \in \mathcal{D}(A)}} D(\mathcal{N}_{A \to B}(\rho_A) || \mathcal{M}_{A \to B}(\sigma_A)), \tag{9}
$$

where  $\mathscr{D}$  is the set of all density matrices.

Theorem 3. *(Adversarial quantum Stein's lemma, informal.) Let* N *and* M *be two quantum channels to distinguish. Then for any*  $\varepsilon \in (0,1)$ *, it holds* 

$$
\lim_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = D^{\inf, \infty}(\mathcal{N} \| \mathcal{M}),\tag{10}
$$

where  $D^{\inf,\infty}(\mathcal{N}|\mathcal{M}) := \lim_{n\to\infty} \frac{1}{n}D^{\inf}(\mathcal{N}^{\otimes n}|\mathcal{M}^{\otimes n})$  and it can be efficiently approximated via a *convex program. Moreover, the optimal exponent can be achieved by non-adaptive strategies, indicating that adaptive strategies by the adversary offer no advantage over non-adaptive ones.*

The key ingredient of its proof is a new chain rule property for the quantum relative entropy, which lower bounds the entropy of a large system into the sum of the entropies of its individual subsystems:

<span id="page-6-1"></span>
$$
D(\mathcal{N}_{A\to B}(\rho_{RA})||\mathcal{M}_{A\to B}(\sigma_{RA})) \ge D(\rho_R || \sigma_R) + D^{\inf,\infty}(\mathcal{N} || \mathcal{M}).
$$
\n(11)

This provides the first quantum analog of the chain rule for Kullback–Leibler divergence in the lower bound direction, complementing the chain rule for quantum relative entropy that establishes an upper bound on the entropy of a global system in the previous work [\[FFRS20\]](#page-56-4).

Application 3: a relative entropy accumulation theorem. The entropy accumulation theorem is a technique to find bounds on the operationally relevant uncertainty present in the outputs of a sequential process as a sum of the worst case uncertainties of each step [\[DFR20\]](#page-55-5). More specifically, a variant for the max-entropy  $H_{\text{max}}^{\varepsilon}$  states that for channels  $\mathcal{N}_i \in \text{CPTP}(R_{i-1}:R_iB_iC_i)$ , we have

$$
H_{\max}^{\varepsilon}(B_1 \dots B_n | C_1 \dots C_n)_{\mathcal{N}_n \circ \dots \circ \mathcal{N}_1(\rho_{R_0})} \leq \sum_{i=1}^n \sup_{\omega_{R_{i-1}}} H(B_i | C_i)_{\mathcal{N}_i(\omega)} + O(\sqrt{n}), \tag{12}
$$

where  $H_{\max}^{\varepsilon}(B|C)_{\rho} \approx -\inf_{\sigma_B \in \mathcal{D}} D_{\text{H},\varepsilon}(\rho_{AB}||\text{id}_{A} \otimes \sigma_B)^2$  $H_{\max}^{\varepsilon}(B|C)_{\rho} \approx -\inf_{\sigma_B \in \mathcal{D}} D_{\text{H},\varepsilon}(\rho_{AB}||\text{id}_{A} \otimes \sigma_B)^2$  and  $H(A|B)_{\rho} = -\inf_{\sigma_B} D(\rho_{AB}||\text{id}_{A} \otimes \sigma_B)$ . This version can be found in [\[MFSR22\]](#page-57-5). It is natural to ask whether such a statement can be generalized to divergences between arbitrary sequential processes of channels, and not only ones corresponding to entropies. This was first asked as an open question in [\[MFSR22\]](#page-57-5) for  $D_{\text{max},\varepsilon}$ . Here we establish such a theorem for  $D_{H,\varepsilon}$ .

**Theorem 4.** *(Relative entropy accumulation, informal.) Consider quantum operations*  $\mathcal{N}_i \in \text{CPTP}(R_{i-1} :$  $R_iA_i$ ),  $M_i \in \mathbb{CP}(R_{i-1}:R_iA_i)$  and states  $\rho_{R_0}, \sigma_{R_0}$ . Let  $\varepsilon \in (0,1)$ , then

$$
D_{\mathrm{H},\varepsilon} \bigg( \operatorname{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{R_0}) \bigg| \operatorname{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{R_0}) \bigg) \n\geq \sum_{i=1}^n D^{\inf,\infty}(\operatorname{Tr}_{R_i} \circ \mathcal{N}_i | \operatorname{Tr}_{R_i} \circ \mathcal{M}_i) - O(n^{2/3} \log n). \tag{13}
$$

Our proof makes important use of the measured relative entropy, in particular to establish a chain rule analogous to  $(11)$ . We leave it as an open question for future work whether this new proof technique can lead to better entropy accumulation theorems. Note that a dual relative entropy accumulation theorem that proves upper bounds on  $D_{\text{max},\varepsilon}$  is unknown.

Application 4: efficient bounds for quantum resource theory. The generalized quantum Stein's lemma [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3) has found broad applications in quantum resource theory. However, the optimal Stein exponents derived in these works are often challenging to compute, limiting their practical utility. In contrast, our new generalized Stein's lemma, presented in Theorem [2,](#page-4-0) offers computational efficiency. To leverage this advantage, even when the task of interest does not directly satisfy all the required assumptions, one can follow a general methodology of relaxing the set of interest to one that does, particularly regarding the polar assumption. For example, in entanglement theory, the set of separable states can be relaxed to the Rains set, which then satisfies all necessary assumptions. Similarly, in fault-tolerant quantum computing, the set of stabilizer states can be relaxed to the set of states with nonpositive mana, which also fulfills all the required conditions. This helps to establish efficient benchmarks to the task of interest and improve the existing bounds. We apply this general methodology to quantum resource theories and provide improved and efficient bounds for entanglement distillation, magic state distillation, and the entanglement cost of quantum states and channels.

<span id="page-6-0"></span><sup>&</sup>lt;sup>2</sup> The exact definition is not in terms of  $D_{\text{H},\varepsilon}$  but a closely related smoothed version of  $D_{\text{S},1/2}$ .

### <span id="page-7-0"></span>1.2 Organization

The organization of the remaining content is as follows. In Section [2,](#page-7-1) we introduce necessary notations and preliminary results on convex optimization and quantum divergences that will be used throughout this work. In Section [3,](#page-11-0) we present the variational formula and superadditivity of the measured relative entropy between two sets of quantum states, which serve as the main technical tools for proving our main results. In Section [4,](#page-15-0) we present our main result on the generalized AEP and its formal proof. In Section [5,](#page-24-0) we derive another generalized quantum Stein's lemma by applying the generalized AEP, and introduce a new framework of quantum resource theory and prove its reversibility. In Section [6,](#page-31-0) we introduce the adversarial quantum channel discrimination and present its corresponding Stein's lemma via the minimun output channel divergence. In Section [7,](#page-41-0) we establish a relative entropy accumulation theorem. In Section [8,](#page-45-0) we present efficient bounds for quantum resource theory by relaxing the originally hard problem to the one that fits our framework. In Section [9,](#page-53-0) we give a conclusion of this work, discussing the significance of this work and some open problem for further study.

## <span id="page-7-1"></span>2 Preliminaries

## <span id="page-7-2"></span>2.1 Notations

In this section we set the notations and define several quantities that will be used throughout this work. Some frequently used notations are summarized in Table [1.](#page-7-4) Note that we label different physical systems by capital Latin letters and use these labels as subscripts to guide the reader by indicating which system a mathematical object belongs to. We drop the subscripts when they are evident in the context of an expression (or if we are not talking about a specific system).

<span id="page-7-4"></span>

<b>Notations</b>	Descriptions
$\mathcal{H}_A$	Hilbert space on system $A$
A	Dimension of $\mathcal{H}_A$
$\mathscr{L}(A)$	Linear operators on $\mathcal{H}_A$
$\mathscr{H}(A)$	Hermitian operators on $\mathcal{H}_A$
$\mathscr{H}_+(A)$	Positive semidefinite operators on $\mathcal{H}_A$
$\mathscr{H}_{++}(A)$	Positive definite operators on $\mathcal{H}_A$
$\mathscr{D}(A)$	Density matrices on $\mathcal{H}_A$
$\mathscr{A}, \mathscr{B}, \mathscr{C}$	Set of linear operators
$\mathscr{C}^\circ$	Polar set $\mathscr{C}^\circ := \{X : \text{Tr}[XY] \leq 1, \forall Y \in \mathscr{C}\}\$ of $\mathscr{C}$
$\mathscr{C}_{\scriptscriptstyle\!+\!\!\!}^\circ$	Polar set restricted to positive semidefinite cone $\mathscr{C}^{\circ} \cap \mathscr{H}_+$
$\mathscr{C}_{\scriptscriptstyle\!+\!\!1}^\circ$	Polar set restricted to positive definite operators $\mathscr{C}^{\circ} \cap \mathscr{H}_{++}$
<b>CPTP</b>	Completely positive and trace preserving maps
$\rm CP$	Completely positive maps
$\log(x)$	Logarithm of $x$ in base two

Table 1: Overview of notational conventions.

#### <span id="page-7-3"></span>2.2 Polar set and support function

In the following, we introduce the definitions of the polar set and support function, along with some fundamental results that will be used in our discussions.

**Definition 5.** Let  $\mathscr C$  be a convex set. Then its polar set is defined by

$$
\mathscr{C}^{\circ} := \{ X : \text{Tr}[XY] \le 1, \forall Y \in \mathscr{C} \}. \tag{14}
$$

Let  $\mathscr{C}_+^{\circ} := \mathscr{C}^{\circ} \cap \mathscr{H}_+$  and  $\mathscr{C}_{++}^{\circ} := \mathscr{C}^{\circ} \cap \mathscr{H}_{++}$  be the intersections with positive semidefinite operators *and positive definite operators, respectively. The support function of*  $\mathscr{C}$  *at*  $\omega$  *is defined by* 

$$
h_{\mathscr{C}}(\omega) := \sup_{\sigma \in \mathscr{C}} \text{Tr}[\omega \sigma]. \tag{15}
$$

It is clear from the definitions that  $\omega \in \mathscr{C}^{\circ}$  if and only if  $h_{\mathscr{C}}(\omega) \leq 1$ . The support function is defined as a convex program. Its dual program can be written as follows.

<span id="page-8-1"></span>**Lemma 6.** Let  $\mathscr{C} \subseteq \mathscr{H}_+$  be a convex set and  $\omega \in \mathscr{H}_+$ . Then  $h_{\mathscr{C}}(\omega) = \inf\{t > 0 : \omega \in t\mathscr{C}_+^{\circ}\}.$ 

*Proof.* We start from ">" direction. Let  $s > h_{\mathscr{C}}(\omega) \geq 0$ . We have  $h_{\mathscr{C}}(\omega/s) = h_{\mathscr{C}}(\omega)/s < 1$ . So  $\omega/s \in \mathscr{C}_+^{\circ}$  and therefore  $\omega \in s\mathscr{C}_+^{\circ}$ . This implies that s is a feasible solution for  $\inf\{t > 0 : \omega \in t\mathscr{C}_+^{\circ}\}\,$ , and we get  $s \ge \inf\{t > 0 : \omega \in t\mathcal{C}_+^{\circ}\}\)$ . As this holds for any  $s > h_{\mathscr{C}}(\omega)$ , we get  $h_{\mathscr{C}}(\omega) \ge \inf\{t > 0 : \omega \in t\mathcal{C}_+^{\circ}\}\$ .  $\omega \in t\mathscr{C}_{+}^{\circ}$ . Now we prove the " $\leq$ " direction. Let z be any feasible solution to  $\inf\{t > 0 : \omega \in t\mathscr{C}_{+}^{\circ}\}.$ We have  $\omega = zX$  with  $X \in \mathcal{C}_+^{\circ}$ . Then  $h_{\mathcal{C}}(\omega) = zh_{\mathcal{C}}(X) \leq z$  because  $X \in \mathcal{C}_+^{\circ}$  implies  $h_{\mathcal{C}}(X) \leq 1$ . As this holds for any feasible solution z, we have  $h_{\mathscr{C}}(\omega) \le \inf\{t > 0 : \omega \in t\mathscr{C}_+^{\circ}$ }. ⊓⊔

**Definition 7.** Let  $\mathcal{H}_1$  and  $\mathcal{H}_2$  be finite-dimensional Hilbert spaces. Consider three sets  $\mathcal{A}_1 \subseteq \mathcal{H}_+(\mathcal{H}_1)$ ,  $\mathscr{A}_2 \subseteq \mathscr{H}_+(\mathcal{H}_2)$ , and  $\mathscr{A}_{12} \subseteq \mathscr{H}_+(\mathcal{H}_1 \otimes \mathcal{H}_2)$ . We call  $\{\mathscr{A}_1, \mathscr{A}_2, \mathscr{A}_{12}\}\$ is closed under tensor product if for *any*  $X_1$  ∈  $\mathscr{A}_1$ ,  $X_2$  ∈  $\mathscr{A}_2$ , we have  $X_1$  ⊗  $X_2$  ∈  $\mathscr{A}_{12}$ . In short, we write  $\mathscr{A}_1$  ⊗  $\mathscr{A}_2$  ⊆  $\mathscr{A}_{12}$ .

The following lemma provides an equivalent condition for determining if the polar sets of interest are closed under tensor product, which can be easier to validate for specific examples.

<span id="page-8-2"></span>**Lemma 8.** Let  $\mathcal{H}_1$  and  $\mathcal{H}_2$  be finite-dimensional Hilbert spaces. Consider three sets  $\mathcal{A}_1 \subseteq \mathcal{H}_+(\mathcal{H}_1)$ ,  $\mathscr{A}_2 \subseteq \mathscr{H}_+(\mathcal{H}_2)$ , and  $\mathscr{A}_{12} \subseteq \mathscr{H}_+(\mathcal{H}_1 \otimes \mathcal{H}_2)$ . Their polar sets are closed under tensor product if and only *if their support functions are sub-multiplicative. That is,*

$$
(\mathscr{A}_1)^{\circ}_{+}\otimes (\mathscr{A}_2)^{\circ}_{+} \subseteq (\mathscr{A}_{12})^{\circ}_{+} \iff h_{\mathscr{A}_{12}}(X_1 \otimes X_2) \leq h_{\mathscr{A}_{1}}(X_1)h_{\mathscr{A}_{2}}(X_2), \ \forall X_i \in \mathscr{H}_{+}(\mathcal{H}_i). \tag{16}
$$

*Proof.* We start from " $\Rightarrow$ " direction. Consider any  $X_1 \in \mathcal{H}_+(\mathcal{H}_1), X_2 \in \mathcal{H}_+(\mathcal{H}_2)$ . Let  $t_1$  and  $t_2$  be any feasible solutions of the dual programs in Lemma [6](#page-8-1) for  $h_{\mathscr{A}_1}(X_1)$  and  $h_{\mathscr{A}_2}(X_2)$ , respectively. Then we have  $X_1 \in t_1(\mathcal{A}_1)^\circ_+$  $\frac{1}{4}$  and  $X_2 \in t_2(\mathcal{A}_2)^\circ_+$ <sup>o</sup>. This implies  $X_1 \otimes X_2 \in t_1t_2(\mathscr{A}_1)_+^{\circ} \otimes (\mathscr{A}_2)_+^{\circ} \subseteq t_1t_2(\mathscr{A}_{12})_+^{\circ}$  $\int_{+}^{\circ}$  and therefore  $t_1t_2$  is also a feasible solution for the dual program of  $h_{\mathscr{A}_{12}}(X_1 \otimes X_2)$ . So  $h_{\mathscr{A}_{12}}(X_1 \otimes X_2) \leq$  $t_1t_2$ . As this holds for any feasible solutions  $t_1, t_2$ , we have  $h_{\mathscr{A}_{12}}(X_1 \otimes X_2) \leq h_{\mathscr{A}_1}(X_1)h_{\mathscr{A}_2}(X_2)$ . Now we prove the " $\Leftarrow$ " direction. For any  $Y_1 \in (\mathcal{A}_1)^{\circ}$  $\frac{1}{4}$  and  $Y_2 \in (\mathcal{A}_2)_+^{\circ}$  $_+^{\circ}$ , we have  $0 \leq h_{\mathscr{A}_1}(Y_1) \leq 1$  and  $0 \le h_{\mathscr{A}_2}(Y_2) \le 1$ . This implies that  $h_{\mathscr{A}_{12}}(Y_1 \otimes Y_2) \le 1$  and therefore,  $Y_1 \otimes Y_2 \in (\mathscr{A}_{12})_+^{\circ}$  $\int_{+}^{\infty}$ . This shows  $(\mathscr{A}_1)^\circ_+ \otimes (\mathscr{A}_2)^\circ_+ \subseteq (\mathscr{A}_{12})^\circ_+$ + . ⊓⊔

#### <span id="page-8-0"></span>2.3 Quantum divergences

A functional  $\mathbb{D}: \mathscr{D} \times \mathscr{H}_+ \to \mathbb{R}$  is a quantum divergence if it satisfies the data-processing inequality  $\mathbb{D}(\mathcal{E}(\rho)\|\mathcal{E}(\sigma)) \leq \mathbb{D}(\rho\|\sigma)$  for any CPTP map  $\mathcal E$  and  $(\rho,\sigma) \in \mathscr{D} \times \mathscr{H}_+$ . In the following, we will introduce several quantum divergences and their fundamental properties, which will be used throughout this work. Additionally, we will define quantum divergences between two sets of quantum states, which will be the main quantity of interest in this work.

**Definition 9.** *(Petz Rényi divergence [[Pet86\]](#page-58-7).) Let*  $\alpha \in (0,1) \cup (1,+\infty)$ *. For any*  $\rho \in \mathscr{D}$  *and*  $\sigma \in \mathscr{H}_+$ *, the Petz Renyi divergence is defined by ´*

$$
D_{P,\alpha}(\rho||\sigma) := \frac{1}{\alpha - 1} \log \text{Tr} \left[ \rho^{\alpha} \sigma^{1-\alpha} \right],\tag{17}
$$

*if* supp $(\rho) \subseteq \text{supp}(\sigma)$ *, and*  $+\infty$  *otherwise.* 

**Definition 10.** *(Sandwiched Rényi divergence [[MLDS](#page-57-6)<sup>+</sup>13, [WWY14\]](#page-59-0).) Let*  $\alpha \in (0,1) \cup (1,+\infty)$ *. For any*  $\rho \in \mathcal{D}$  *and*  $\sigma \in \mathcal{H}_+$ , the sandwiched Rényi divergence is defined by

$$
D_{\mathcal{S},\alpha}(\rho\|\sigma) := \frac{1}{\alpha - 1} \log \text{Tr} \left[ \sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}} \right]^{\alpha},\tag{18}
$$

*if* supp $(\rho) \subseteq$  supp $(\sigma)$ *, and*  $+\infty$  *otherwise.* 

**Definition 11.** *(Umegaki relative entropy [\[Ume54\]](#page-58-8).) For any*  $\rho \in \mathcal{D}$  *and*  $\sigma \in \mathcal{H}_+$ *, the Umegaki relative entropy is defined by*

$$
D(\rho \|\sigma) := \text{Tr}[\rho(\log \rho - \log \sigma)],\tag{19}
$$

*if* supp $(\rho) \subseteq \text{supp}(\sigma)$  *and*  $+\infty$  *otherwise.* 

When  $\alpha \to 1$ , both  $D_{P,Q}$  and  $D_{S,Q}$  converge to the Umegaki relative entropy [\[MLDS](#page-57-6)<sup>+</sup>13, [WWY14\]](#page-59-0),

$$
\lim_{\alpha \to 1} D_{P,\alpha}(\rho \| \sigma) = \lim_{\alpha \to 1} D_{S,\alpha}(\rho \| \sigma) = D(\rho \| \sigma).
$$
\n(20)

When  $\alpha \to 0^+$ , the Petz Rényi divergence converges to the min-relative entropy or the Petz Rényi divergence of order 0,

$$
\lim_{\alpha \to 0^+} D_{P,\alpha}(\rho \| \sigma) = D_{\min}(\rho \| \sigma) = D_{P,0}(\rho \| \sigma) := -\log \text{Tr}[\Pi_{\rho} \sigma],\tag{21}
$$

with  $\Pi_{\rho}$  the projection on the support of  $\rho$ . When  $\alpha \to \infty$ , the sandwiched Rényi divergence converges to the max-relative entropy [\[Dat09,](#page-55-6) [Ren05\]](#page-58-3),

$$
\lim_{\alpha \to \infty} D_{\mathcal{S},\alpha}(\rho \| \sigma) = D_{\max}(\rho \| \sigma) := \log \inf \left\{ t \in \mathbb{R} \; : \; \rho \le t\sigma \right\},\tag{22}
$$

if  $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$  and  $+\infty$  otherwise. Let  $F(\rho, \sigma) := \|\sqrt{\rho}\sqrt{\sigma}\|_1 + \sqrt{(1 - \text{Tr} \rho)(1 - \text{Tr} \sigma)}$  be the generalized fidelity and  $P(\rho, \sigma) := \sqrt{1 - F^2(\rho, \sigma)}$  be the purified distance. Let  $\varepsilon \in (0, 1)$ . Then the smoothed max-relative entropy is defined by

$$
D_{\max,\varepsilon}(\rho\|\sigma) := \inf_{\rho':P(\rho',\rho)\leq\varepsilon} D_{\max}(\rho'\|\sigma),\tag{23}
$$

where the infimum is taken over all subnormalized states that are  $\varepsilon$ -close to the state  $\rho$ .

**Definition 12.** *(Hypothesis testing relative entropy.) Let*  $\varepsilon \in [0,1]$ *. For any*  $\rho \in \mathscr{D}$  *and*  $\sigma \in \mathscr{H}_{+}$ *, the quantum hypothesis testing relative entropy is defined by*  $D_{H,\varepsilon}(\rho||\sigma) := -\log \beta_{\varepsilon}(\rho||\sigma)$  *where* 

$$
\beta_{\varepsilon}(\rho \| \sigma) := \min_{0 \le M \le I} \{ \text{Tr}[\sigma M] : \text{Tr}[\rho (I - M)] \le \varepsilon \}.
$$
\n(24)

The following lemma combines [\[CMW16,](#page-55-7) Lemma 5] and [\[QWW18,](#page-58-9) Proposition 3].

<span id="page-9-0"></span>**Lemma 13.** Let  $\alpha \in (0,1)$ ,  $\alpha' \in (1,+\infty)$  and  $\varepsilon \in (0,1)$ . For any  $\rho \in \mathscr{D}$  and  $\sigma \in \mathscr{H}_+$ , it holds

$$
D_{P,\alpha}(\rho\|\sigma) + \frac{\alpha}{\alpha - 1}\log\frac{1}{\varepsilon} \le D_{H,\varepsilon}(\rho\|\sigma) \le D_{S,\alpha'}(\rho\|\sigma) + \frac{\alpha'}{\alpha' - 1}\log\frac{1}{1 - \varepsilon}.\tag{25}
$$

**Definition 14.** *(Measured relative entropy [\[Don86,](#page-55-8) [HP91\]](#page-56-1).) For any*  $\rho \in \mathcal{D}$  *and*  $\sigma \in \mathcal{H}_+$ *, the measured relative entropy is defined by*

$$
D_M(\rho \|\sigma) := \sup_{(\mathcal{X}, M)} D(P_{\rho,M} \| P_{\sigma,M}), \tag{26}
$$

*where* D *is the Kullback–Leibler divergence and the optimization is over finite sets* X *and positive operator valued measures* M *on* X *such that*  $M_x \geq 0$  *and*  $\sum_{x \in \mathcal{X}} M_x = I$ ,  $P_{\rho,M}$  *is a measure on* X *defined via the relation*  $P_{\rho,M}(x) = \text{Tr}[M_x \rho]$  *for any*  $x \in \mathcal{X}$ *.* 

A variational expression for  $D_M$  is given by [\[BFT17,](#page-54-3) Lemma 1],

$$
D_{\mathcal{M}}(\rho \| \sigma) = \sup_{\omega \in \mathcal{H}_{++}} \text{Tr}[\rho \log \omega] + 1 - \text{Tr}[\sigma \omega]. \tag{27}
$$

**Definition 15.** *(Measured Rényi divergence [[BFT17\]](#page-54-3).) Let*  $\alpha \in (0,1) \cup (1,\infty)$ *. For any*  $\rho \in \mathscr{D}$  *and*  $\sigma \in \mathcal{H}_{+}$ , the measured Rényi divergence is defined as

<span id="page-10-0"></span>
$$
D_{\mathcal{M},\alpha}(\rho\|\sigma) := \sup_{(\mathcal{X},M)} D_{\alpha}(P_{\rho,M} \| P_{\sigma,M}),
$$
\n(28)

*where*  $D_{\alpha}$  *is the classical Rényi divergence.* 

A variational expression for  $D_{\text{M},\alpha}$  is given by [\[BFT17,](#page-54-3) Lemma 3],

$$
D_{\mathbf{M},\alpha}(\rho||\sigma) = \frac{1}{\alpha - 1} \log \begin{cases} \inf_{W \in \mathcal{H}_{++}} \alpha \operatorname{Tr}[\rho W] + (1 - \alpha) \operatorname{Tr} \left[ \sigma W^{\frac{\alpha}{\alpha - 1}} \right], & \text{if } \alpha \in (0, 1/2), \\ \inf_{W \in \mathcal{H}_{++}} \alpha \operatorname{Tr} \left[ \rho W^{\frac{\alpha - 1}{\alpha}} \right] + (1 - \alpha) \operatorname{Tr}[\sigma W], & \text{if } \alpha \in [1/2, 1), \\ \sup_{W \in \mathcal{H}_{++}} \alpha \operatorname{Tr} \left[ \rho W^{\frac{\alpha - 1}{\alpha}} \right] + (1 - \alpha) \operatorname{Tr}[\sigma W], & \text{if } \alpha \in (1, \infty). \end{cases} \tag{29}
$$

When  $\alpha \to 1$ , the measured Rényi divergence converges to the measured relative entropy. It is known that [\[RSB24,](#page-58-10) Lemma 2]  $D_{\text{M},\alpha}$  is jointly convex for  $\alpha \in (0,1]$  and jointly quasi-convex for  $\alpha \in (1,\infty)$ . For any  $\mathcal{E} \in \text{CPTP}$  and  $\alpha > 0$ , the data-processing inequality holds [\[RSB24,](#page-58-10) Lemma 5]

<span id="page-10-1"></span>
$$
D_{\mathcal{M},\alpha}(\mathcal{E}(\rho)\|\mathcal{E}(\sigma)) \leq D_{\mathcal{M},\alpha}(\rho\|\sigma). \tag{30}
$$

The following result shows the ordering relation among different relative entropies.

<span id="page-10-3"></span>**Lemma 16.** *Let*  $\alpha \in [1/2, 1)$ *. For any*  $\rho \in \mathcal{D}$  *and*  $\sigma \in \mathcal{H}_+$ *,* 

$$
D_{\min}(\rho \|\sigma) \le D_{\mathcal{M},\alpha}(\rho\|\sigma) \le D_{\mathcal{M}}(\rho\|\sigma) \le D(\rho\|\sigma). \tag{31}
$$

*Proof.* The last two inequalities follow from the monotonicity in  $\alpha$  of the classical Rényi divergences and the data processing inequality for D. As  $D_{\text{M},\alpha}$  is monotone increasing in  $\alpha$ , it remains to show the first inequality for  $\alpha = \frac{1}{2}$  $\frac{1}{2}$ . By the variational formula in [\[BFT17,](#page-54-3) Eq. (21)], we have

$$
D_{\mathrm{M},1/2}(\rho||\sigma) = -\log \inf_{\omega \in \mathcal{H}_{++}} \mathrm{Tr}[\rho \omega^{-1}] \mathrm{Tr}[\sigma \omega],\tag{32}
$$

which is the same as the Alberti's theorem for quantum fidelity (see e.g. [\[Wat18,](#page-59-1) Corollary 3.20]). Consider a feasible solution  $\omega_{\varepsilon} = \Pi_{\rho} + \varepsilon (I - \Pi_{\rho}) \in \mathcal{H}_{++}$  with  $\varepsilon > 0$ . It gives  $D_{\text{M},1/2}(\rho || \sigma) \ge$  $-\log \text{Tr}[\rho \omega_{\varepsilon}^{-1}] \text{Tr}[\sigma \omega_{\varepsilon}]$ . Since  $\rho$  has trace one, it gives  $\text{Tr}[\rho \omega_{\varepsilon}^{-1}] = \text{Tr}[\rho] = 1$ . Then we have

$$
D_{\text{M},1/2}(\rho||\sigma) \ge -\log \text{Tr}[\sigma\omega_{\varepsilon}] = -\log[(1-\varepsilon)\text{Tr}\,\Pi_{\rho}\sigma + \varepsilon].\tag{33}
$$

As the above holds for any  $\varepsilon > 0$ , we take  $\varepsilon \to 0^+$  and get  $D_{\text{M},1/2}(\rho||\sigma) \ge -\log \text{Tr}[\Pi_\rho \sigma] = D_{\text{min}}(\rho||\sigma)$ , which completes the proof. □

<span id="page-10-2"></span>**Lemma 17.** Let  $\alpha \in [1/2, \infty)$ *. For any*  $\rho \in \mathcal{D}$  *and*  $\sigma \in \mathcal{H}_+$ *, it hold* 

$$
D_{\mathcal{M},\alpha}(\rho||\sigma) \leq D_{\mathcal{S},\alpha}(\rho||\sigma) \leq D_{\mathcal{M},\alpha}(\rho||\sigma) + 2\log|\text{spec}(\sigma)|,\tag{34}
$$

*where*  $|\text{spec}(\sigma)|$  *is the number of mutually different eigenvalues of*  $\sigma$ *.* 

*Proof.* The first inequality is given by [\[BFT17,](#page-54-3) Theorem 6]. Now we prove the second inequality. Let  $P_{\sigma}$  be a pinching map with respect to  $\sigma$ . Then we have

$$
D_{\mathcal{S},\alpha}(\rho||\sigma) \le D_{\mathcal{S},\alpha}(\mathcal{P}_{\sigma}(\rho)||\sigma) + 2\log|\text{spec}(\sigma)| \tag{35}
$$

$$
=D_{\mathrm{M},\alpha}(\mathcal{P}_{\sigma}(\rho)\|\sigma)+2\log|\mathrm{spec}(\sigma)|\tag{36}
$$

$$
=D_{\mathcal{M},\alpha}(\mathcal{P}_{\sigma}(\rho)\|\mathcal{P}_{\sigma}(\sigma)) + 2\log|\text{spec}(\sigma)| \tag{37}
$$

$$
\leq D_{\mathrm{M},\alpha}(\rho||\sigma) + 2\log|\mathrm{spec}(\sigma)|\tag{38}
$$

where the first line follows from [\[HT16,](#page-56-5) Lemma 3], the second line follows because  $\mathcal{P}_{\sigma}(\rho)$  and  $\sigma$  commute and therefore  $D_{\text{S},\alpha}(\mathcal{P}_{\sigma}(\rho) \| \sigma) = D_{\text{M},\alpha}(\mathcal{P}_{\sigma}(\rho) \| \sigma)$ , the third line follows by  $\mathcal{P}_{\sigma}(\sigma) = \sigma$  and the last line follows from the data processing inequality. line follows from the data processing inequality. ⊓⊔

<span id="page-11-3"></span>**Lemma 18.** Let X be a permutation-invarant operator on  $\mathscr{L}(A^{\otimes n})$  with  $|A| = d$ . Then  $|\text{spec}(X)| \le$  $(n+1)^d(n+d)^{d^2} = \text{poly}(n)$ [\[FF21b,](#page-56-6) Lemma A.1].

In this work, we will focus on the study of quantum divergences between two sets of quantum states.

<span id="page-11-4"></span>Definition 19. *(Quantum divergence between two sets of states.) Let* D *be a quantum divergence between two quantum states. Then for any sets*  $\mathscr{A} \subseteq \mathscr{D}$  *and*  $\mathscr{B} \subseteq \mathscr{H}_+$ , the quantum divergence between these *two sets of quantum states is defined by*

$$
\mathbb{D}(\mathscr{A}||\mathscr{B}) := \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \mathbb{D}(\rho||\sigma). \tag{39}
$$

Note that if  $\mathbb D$  is lower semicontinuous (which is true for most quantum divergences of interest), and  $\mathscr A$  and  $\mathscr B$  are compact sets, the infimum in the above expression is always attained and can thus be replaced by a minimization [\[KZ05,](#page-57-7) Theorem 7.3.1]. From a geometric perspective, this quantity characterizes the distance between two sets  $\mathscr A$  and  $\mathscr B$  under the "distance metric"  $\mathbb D$ . In particular, if  $\mathscr{A} = \{\rho\}$  is a singleton, we write  $\mathbb{D}(\rho||\mathscr{B}) := \mathbb{D}(\{\rho\}||\mathscr{B})$ . For two sequences of sets  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$ , the regularized divergence is defined by

$$
\mathbb{D}^{\infty}(\mathscr{A}||\mathscr{B}) := \lim_{n \to \infty} \frac{1}{n} \mathbb{D}(\mathscr{A}_n||\mathscr{B}_n),\tag{40}
$$

whenever the limit on the right-hand side exists.

## <span id="page-11-0"></span>3 Variational formula and superadditivity

Unlike other quantum relative entropies, which are defined by closed-form expressions, the measured relative entropy requires maximization over all possible quantum measurements, making it inherently challenging to compute. This complexity is compounded in the case of divergences between different sets of quantum states, where additional layers of optimization transform it into a minimax problem. The following results address this issue by providing variational formulas for  $D_M(\mathscr{A} \| \mathscr{B})$  and  $D_{M,\alpha}(\mathscr{A} \| \mathscr{B})$ , reformulating them as more tractable convex optimization programs. Based on these variational formulas, we establish their superadditivity, which serves as a key technical tool for the main results of this work in the subsequent sections.

<span id="page-11-2"></span>**Lemma 20.** Let  $\mathcal{A} \subseteq \mathcal{D}$  and  $\mathcal{B} \subseteq \mathcal{H}_+$  be two compact convex sets. Then it holds

<span id="page-11-1"></span>
$$
D_M(\mathscr{A}||\mathscr{B}) = \sup_{W \in \mathscr{B}^{\circ}_{++}} -h_{\mathscr{A}}(-\log W),\tag{41}
$$

*where the objective function on the right-hand side is concave in* W*.*

*Proof.* By the variational formula of the measured relative entropy in Eq. [\(27\)](#page-10-0), we have

$$
D_M(\mathscr{A} \| \mathscr{B}) = \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \sup_{W \in \mathscr{H}_{++}} \text{Tr}[\rho \log W] + 1 - \text{Tr}[\sigma W]. \tag{42}
$$

Note that all  $\mathscr{A}, \mathscr{B}, \mathscr{H}_{++}$  are convex sets, with  $\mathscr{A}, \mathscr{B}$  being compact. Moreover, the objective function is linear in  $(\rho, \sigma)$  and concave in W. So we can apply Sion's minimax theorem [\[Sio58,](#page-58-11) Corollary 3.3] to exchange the infimum and supremum and get

$$
D_M(\mathscr{A}||\mathscr{B}) = \sup_{W \in \mathscr{H}_{++}} \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \text{Tr}[\rho \log W] + 1 - \text{Tr}[\sigma W] \tag{43}
$$

$$
= \sup_{W \in \mathcal{H}_{+}} \inf_{\rho \in \mathcal{A}} \text{Tr}[\rho \log W] + 1 - \sup_{\sigma \in \mathcal{B}} \text{Tr}[\sigma W]. \tag{44}
$$

By using the definition of the support function, we have

$$
D_M(\mathscr{A}||\mathscr{B}) = \sup_{W \in \mathscr{H}_{++}} -h_{\mathscr{A}}(-\log W) + 1 - h_{\mathscr{B}}(W). \tag{45}
$$

Next, we aim to simplify the objective function by moving the term  $1 - h\mathscr{B}(W)$  into the constraint. For this, let W be a feasible solution in the above optimization and define  $W = W/h_{\mathscr{B}}(W)$ . Then  $h_{\mathscr{B}}(W) = 1$  and we will see that W achieves an objective value no smaller than W. Indeed

$$
-h_{\mathscr{A}}(-\log \widetilde{W}) = -h_{\mathscr{A}}(-\log W) - \log h_{\mathscr{B}}(W) \ge -h_{\mathscr{A}}(-\log W) + 1 - h_{\mathscr{B}}(W) \tag{46}
$$

where the inequality follows from the fact that  $\log x \leq x - 1$ . Therefore, we can reformulate

$$
D_M(\mathscr{A} \| \mathscr{B}) = \sup_{\substack{W \in \mathscr{H}_{++} \\ h_{\mathscr{B}}(W) = 1}} -h_{\mathscr{A}}(-\log W). \tag{47}
$$

Using the same argument as above, we can relax the condition  $h_{\mathscr{B}}(W) = 1$  to  $h_{\mathscr{B}}(W) \le 1$  and get

$$
D_{\mathcal{M}}(\mathscr{A} \| \mathscr{B}) = \sup_{\substack{W \in \mathscr{H}_{++} \\ h_{\mathscr{B}}(W) \le 1}} -h_{\mathscr{A}}(-\log W). \tag{48}
$$

Finally, noting that  $h_{\mathscr{B}}(W) \leq 1$  if and only if  $W \in \mathscr{B}^{\circ}$ , we have the expression in Eq. [\(41\)](#page-11-1). Let  $f(W) \coloneqq$  $-h<sub>\mathscr{A}</sub>(-\log W) = \inf_{\rho \in \mathscr{A}} \text{Tr}[\rho \log W]$  be the objective function. Consider any convex combination  $\lambda W_1 + (1 - \lambda)W_2$  with  $\lambda \in (0, 1)$ . We have

$$
f(\lambda W_1 + (1 - \lambda)W_2) = \inf_{\rho \in \mathscr{A}} \text{Tr}[\rho \log(\lambda W_1 + (1 - \lambda)W_2)] \tag{49}
$$

$$
\geq \inf_{\rho \in \mathscr{A}} \lambda \operatorname{Tr}[\rho \log W_1] + (1 - \lambda) \operatorname{Tr}[\rho \log W_2] \tag{50}
$$

$$
\geq \lambda \inf_{\rho \in \mathscr{A}} \text{Tr}[\rho \log W_1] + (1 - \lambda) \inf_{\rho \in \mathscr{A}} \text{Tr}[\rho \log W_2] \tag{51}
$$

$$
= \lambda f(W_1) + (1 - \lambda)f(W_2)
$$
\n<sup>(52)</sup>

where the second line follows from the operator concavity of logarithm and the third line follows by putting infimum to both parts. This proves the concavity of the objective function. □

Following the same argument, we can show the variational formula for the measured Rényi divergence between two sets of quantum states. We present the result for  $\alpha \in [1/2, 1)$  and the cases with other parameters are given in Appendix [A.](#page-60-0)

<span id="page-13-1"></span>**Lemma 21.** Let  $\alpha \in [1/2, 1)$ . Let  $\mathscr{A} \subseteq \mathscr{D}$  and  $\mathscr{B} \subseteq \mathscr{H}_+$  be two compact convex sets. Then it holds

$$
D_{\mathcal{M},\alpha}(\mathscr{A}||\mathscr{B}) = \frac{\alpha}{\alpha - 1} \log \inf_{W,V} \left\{ h_{\mathscr{A}}(V) : W \in \mathscr{B}^{\circ}_{++}, W^{1 - \frac{1}{\alpha}} \le V \right\},\tag{53}
$$

*where the right-hand side is a convex program.*

*Proof.* For any fixed  $\rho \in \mathcal{A}$  and  $\sigma \in \mathcal{B}$ , we have by Eq. [\(29\)](#page-10-1) that

$$
Q_{\mathcal{M},\alpha}(\rho||\sigma) := \inf_{W \in \mathcal{H}_{++}} \alpha \operatorname{Tr} \left[ \rho W^{1-\frac{1}{\alpha}} \right] + (1-\alpha) \operatorname{Tr}[\sigma W] \tag{54}
$$

<span id="page-13-0"></span>
$$
= \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{1-\frac{1}{\alpha}} \le V}} \alpha \operatorname{Tr} \left[ \rho V \right] + (1 - \alpha) \operatorname{Tr} [\sigma W], \tag{55}
$$

where the second equality follows by introducing an additional variable  $V$ . Then we have

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{-1} \stackrel{\perp}{\alpha} \leq V}} \alpha \operatorname{Tr}[\rho V] + (1 - \alpha) \operatorname{Tr}[\sigma W]. \tag{56}
$$

Note that all  $\mathscr{A}, \mathscr{B}$  and  $\{(W, V) : W \in \mathscr{H}_{++}, W^{1-\frac{1}{\alpha}} \leq V\}$  are convex sets, with  $\mathscr{A}, \mathscr{B}$  being compact. Moreover, the objective function is linear in  $(\rho, \sigma)$ , and also linear in  $(W, V)$ . So we can apply Sion's minimax theorem [\[Sio58,](#page-58-11) Corollary 3.3] to exchange the infimum and supremum and get

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{1-\frac{1}{\alpha}} \leq V}} \sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \alpha \operatorname{Tr}[\rho V] + (1 - \alpha) \operatorname{Tr}[\sigma W] \tag{57}
$$

$$
= \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{1-\frac{1}{\alpha}} \leq V}} \alpha \sup_{\rho \in \mathcal{A}} \text{Tr}[\rho V] + (1 - \alpha) \sup_{\sigma \in \mathcal{B}} \text{Tr}[\sigma W]. \tag{58}
$$

By using the definition of the support function, we have

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{1-\frac{1}{\alpha}} \leq V}} \alpha h_{\mathscr{A}}(V) + (1 - \alpha) h_{\mathscr{B}}(W) \tag{59}
$$

$$
= \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{1-\frac{1}{\alpha}} \leq V}} h_{\mathscr{A}}(V)^{\alpha} h_{\mathscr{B}}(W)^{1-\alpha}, \tag{60}
$$

where the second line follows from the weighed arithmetic-geometric mean inequality  $\alpha x + (1 \alpha$ )y  $\leq x^{\alpha}y^{1-\alpha}$  (with equality if and only if  $x = y$ ) and the fact that  $(W, V)$  is a feasible solution implies  $(kW, k^{1-1/\alpha}V)$  is also a feasible solution for any  $k \geq 0$ . Therefore, we can choose  $k = (h_{\mathscr{A}}(V))^{\alpha}(h_{\mathscr{B}}(W))^{-\alpha}$ , which implies  $h_{\mathscr{A}}(k^{1-1/\alpha}V) = h_{\mathscr{B}}(kW)$  and therefore the equality of the weighed arithmetic-geometric mean is achieved. Similarly, for any feasible solution  $(W, V)$  we can always construct a new solution  $(W/h_{\mathscr{B}}(W), V h_{\mathscr{B}}(W)^{1/\alpha-1})$  achieves the same objective value. This implies

$$
\sup_{\substack{\rho \in \mathcal{A} \\ \sigma \in \mathcal{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{1-\frac{1}{\alpha}} \leq V \\ h_{\mathcal{B}}(W) = 1}} (h_{\mathcal{A}}(V))^{\alpha} = \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{1-\frac{1}{\alpha}} \leq V \\ h_{\mathcal{B}}(W) \leq 1}} (h_{\mathcal{A}}(V))^{\alpha},
$$
\n(61)

where the second equality follows by the same reasoning. Finally, noting that  $h\mathscr{R}(W) \leq 1$  if and only if  $W \in \mathscr{B}^{\circ}$ , we have the asserted result in Eq. [\(53\)](#page-13-0). It is easy to check that the objective function  $h_{\mathscr{A}}(V)$ is convex in V and the feasible set is also a convex set.  $□$ 

The variational formula helps establish the following superadditivity of the divergence between two sets of quantum states, which serves as a key technical tool for the applications discussed in the subsequent sections.

<span id="page-14-2"></span>**Lemma 22.** Let  $\alpha \in [1/2, 1)$  and  $\mathcal{H}_1, \mathcal{H}_2$  be finite-dimensional Hilbert spaces. Consider convex and *compact sets*  $A_1 \subseteq \mathcal{D}(\mathcal{H}_1)$ *,*  $A_2 \subseteq \mathcal{D}(\mathcal{H}_2)$ *, and*  $A_{12} \subseteq \mathcal{D}(\mathcal{H}_1 \otimes \mathcal{H}_2)$  *and also*  $B_1 \subseteq \mathcal{H}_+(\mathcal{H}_1)$ *,*  $B_2 \subseteq$  $\mathscr{H}_{+}(\mathcal{H}_{2})$ , and  $\mathscr{B}_{12} \subseteq \mathscr{H}_{+}(\mathcal{H}_{1} \otimes \mathcal{H}_{2})$ . Assume that  $(\mathscr{A}_{1})_{+}^{\circ} \otimes (\mathscr{A}_{2})_{+}^{\circ} \subseteq (\mathscr{A}_{12})_{+}^{\circ}$  $\int_{+}^{\circ}$  *and*  $(\mathscr{B}_1)_{+}^{\circ} \otimes (\mathscr{B}_2)_{+}^{\circ} \subseteq$  $(\overline{\mathscr{B}}_{12})_{+}^{\circ}$ *+ . Then,*

<span id="page-14-0"></span>
$$
D_{\mathrm{M},\alpha}(\mathscr{A}_{12}||\mathscr{B}_{12}) \ge D_{\mathrm{M},\alpha}(\mathscr{A}_1||\mathscr{B}_1) + D_{\mathrm{M},\alpha}(\mathscr{A}_2||\mathscr{B}_2). \tag{62}
$$

*Proof.* Consider the convex program in Lemma [21](#page-13-1) for  $D_{\text{M},\alpha}(\mathscr{A}_1|\mathscr{B}_1)$  and  $D_{\text{M},\alpha}(\mathscr{A}_2|\mathscr{B}_2)$ . Let  $(W_1, V_1)$ and  $(W_2, V_2)$  be any feasible solutions for these programs, respectively. Then we will see that  $(W_1 \otimes$  $W_2, V_1 \otimes V_2$ ) is also a feasible solution for  $D_{M,\alpha}(\mathscr{A}_{12}|\mathscr{B}_{12})$  with a higher objective value. To see this, it is clear that  $W_1 \in (\mathscr{B}_1)^\circ$  and  $W_2 \in (\mathscr{B}_2)^\circ$  implies that  $W_1 \otimes W_2 \in (\mathscr{B}_{12})^\circ$  by the assumption of  $(\mathscr{B}_1)^\circ_+ \otimes (\mathscr{B}_2)^\circ_+ \subseteq (\mathscr{B}_{12})^\circ_+$  $\int_{+}^{\infty}$ . Then by the multiplicativity of the power function, we have

$$
(W_1 \otimes W_2)^{\frac{\alpha-1}{\alpha}} = (W_1)^{\frac{\alpha-1}{\alpha}} \otimes (W_2)^{\frac{\alpha-1}{\alpha}} \le V_1 \otimes V_2.
$$
 (63)

This confirms that  $(W_1 \otimes W_2, V_1 \otimes V_2)$  is a feasible solution for the optimization of  $D_{\text{M},\alpha}(\mathscr{A}_{12}||\mathscr{B}_{12})$ . By the assumption of  $(\mathscr{A}_1)^\circ_+ \otimes (\mathscr{A}_2)^\circ_+ \subseteq (\mathscr{A}_{12})^\circ_+$  $_{+}^{\circ}$  and Lemma [8,](#page-8-2) we have  $h_{\mathscr{A}_{12}}(V_1 \otimes V_2) \leq h_{\mathscr{A}_{1}}(V_1)h_{\mathscr{A}_{2}}(V_2)$ , which implies

$$
\frac{\alpha}{\alpha-1}\log h_{\mathscr{A}_{12}}(V_1\otimes V_2)\geq \frac{\alpha}{\alpha-1}\log h_{\mathscr{A}_1}(V_1)+\frac{\alpha}{\alpha-1}\log h_{\mathscr{A}_2}(V_2). \hspace{1cm} (64)
$$

As the above relation holds for any feasible solutions, we have the asserted result in Eq. [\(62\)](#page-14-0). □

By using the variational formula in Appendix [A,](#page-60-0) we know that the superadditivity also holds for  $\alpha \in (0, 1/2)$ . Next, we demonstrate the superadditivity of the measured relative entropy between two sets of quantum states (i.e., the case  $\alpha = 1$ ). Instead of relying directly on the variational formula, we employ a continuity argument. This approach allows us to avoid the assumption that the set is closed under partial trace. In fact, the natural direct proof of superadditivity via the variational formula seems to require this assumption (see more details in Appendix [D\)](#page-64-0).

<span id="page-14-3"></span>**Lemma 23.** Let  $\mathcal{A} \subseteq \mathcal{D}$  and  $\mathcal{B} \subseteq \mathcal{H}$  be two compact convex sets. Then

<span id="page-14-1"></span>
$$
\lim_{\alpha \to 1^{-}} D_{\mathcal{M},\alpha}(\mathscr{A} \| \mathscr{B}) = \sup_{\alpha \in (0,1)} D_{\mathcal{M},\alpha}(\mathscr{A} \| \mathscr{B}) = D_{\mathcal{M}}(\mathscr{A} \| \mathscr{B}). \tag{65}
$$

*Proof.* Since  $D_{M,\alpha}$  is monotonically increasing in  $\alpha$ , we have for any  $\alpha > \beta$ ,

$$
D_{\mathbf{M},\alpha}(\mathscr{A}||\mathscr{B}) = \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} D_{\mathbf{M},\alpha}(\rho||\sigma) \ge \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} D_{\mathbf{M},\beta}(\rho||\sigma) = D_{\mathbf{M},\beta}(\mathscr{A}||\mathscr{B}). \tag{66}
$$

So the divergence  $D_{\text{M},\alpha}(\mathscr{A}|\mathscr{B})$  is also monotonically increasing in  $\alpha$ . This implies the first equality in Eq. [\(65\)](#page-14-1). Now we prove the second equality. Note that  $D_{\text{M},\alpha}(\rho||\sigma)$  is lower semi-continous on  $\mathcal{H}_+ \times \mathcal{H}_+$  for every  $\alpha > 0$  [\[MH23,](#page-57-8) Proposition III.11] and it is montonic increasing in  $\alpha$  for every  $(\rho, \sigma) \in \mathcal{D} \times \mathcal{D}$ . Therefore, we can apply the minimax theorem given by [\[MH11,](#page-57-9) Corollary A.2] and get

$$
\sup_{\alpha \in (0,1)} D_{\mathcal{M},\alpha}(\mathscr{A} \| \mathscr{B}) = \sup_{\alpha \in (0,1)} \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} D_{\mathcal{M},\alpha}(\rho \| \sigma)
$$
(67)

$$
= \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \sup_{\alpha \in (0,1)} D_{\mathbf{M},\alpha}(\rho \| \sigma)
$$
(68)

$$
= \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} D_{\scriptscriptstyle{\mathrm{M}}}(\rho \| \sigma) \tag{69}
$$

$$
=D_{\mathcal{M}}(\mathscr{A}||\mathscr{B}),\tag{70}
$$

where the third line follows from the continuity of  $D_{M,\alpha}$  at  $\alpha = 1$  for quantum states. □

Combining Lemma [22](#page-14-2) and Lemma [23,](#page-14-3) we have the superadditivity for measured relative entropy.

<span id="page-15-2"></span>**Lemma 24.** *Let*  $\mathcal{H}_1, \mathcal{H}_2$  *be finite-dimensional Hilbert spaces. Consider convex and compact sets*  $\mathcal{A}_1 \subseteq$  $\mathscr{D}(\mathcal{H}_1)$ *,*  $\mathscr{A}_2 \subseteq \mathscr{D}(\mathcal{H}_2)$ *, and*  $\mathscr{A}_{12} \subseteq \mathscr{D}(\mathcal{H}_1 \otimes \mathcal{H}_2)$  *and also*  $\mathscr{B}_1 \subseteq \mathscr{H}_+(\mathcal{H}_1)$ *,*  $\mathscr{B}_2 \subseteq \mathscr{H}_+(\mathcal{H}_2)$ *, and*  $\mathscr{B}_{12} \subseteq \mathscr{H}_+(\mathcal{H}_1 \otimes \mathcal{H}_2)$ . Assume that  $(\mathscr{A}_1)^{\circ}_+ \otimes (\mathscr{A}_2)^{\circ}_+ \subseteq (\mathscr{A}_{12})^{\circ}_+$  $\left(\mathscr{B}_{1}\right)^{\circ}_{+}\otimes\left(\mathscr{B}_{2}\right)^{\circ}_{+}\subseteq\left(\mathscr{B}_{12}\right)^{\circ}_{+}$ *+ . Then,*

$$
D_{\scriptscriptstyle{\mathrm{M}}}(\mathscr{A}_{12}||\mathscr{B}_{12}) \ge D_{\scriptscriptstyle{\mathrm{M}}}(\mathscr{A}_{1}||\mathscr{B}_{1}) + D_{\scriptscriptstyle{\mathrm{M}}}(\mathscr{A}_{2}||\mathscr{B}_{2}). \tag{71}
$$

Remark 1 The above superadditivity is different than the result by Piani [\[Pia09,](#page-58-6) Theorem 1] under compatible assumptions. Here we do not put constrains on the performed measurements. Therefore, the measurements and the set of quantum states may not be compatible in the case here.

## <span id="page-15-0"></span>4 Generalized quantum asymptotic equipartition property

The asymptotic equipartition property (AEP) is a fundamental concept in information theory that de-scribes the behavior of sequences of random variables as the sequence length increases [\[Cov99,](#page-55-0) [HR11\]](#page-56-7). It essentially states that, for a large number of independent and identically distributed (i.i.d.) random variables, the sequences exhibit regular and predictable behavior when considered collectively, despite individual randomness. This concept has been extended to quantum information theory, where the quan-tum version of the AEP applies to quantum states and quantum entropy (e.g. [\[TCR09,](#page-58-2) [Tom16\]](#page-58-12)). This extension is particularly useful in the study of quantum source coding and other quantum communication protocols. In this work, we further generalize the quantum AEP beyond the i.i.d. framework by considering quantum states drawn from two sets that satisfy the following assumptions.

<span id="page-15-1"></span>**Assumption 25.** *Consider a family of sets*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  *satisfying the following properties,* 

- $(A.1)$  Each  $\mathcal{A}_n$  is convex and compact;
- $(A.2)$  Each  $\mathscr{A}_n$  is permutation-invariant;
- *(A.3)*  $\mathscr{A}_m \otimes \mathscr{A}_k \subseteq \mathscr{A}_{m+k}$ , for all  $m, k \in \mathbb{N}$ ;
- $(A.4)$   $(\mathscr{A}_m)^{\circ}_{+} \otimes (\mathscr{A}_k)^{\circ}_{+} \subseteq (\mathscr{A}_{m+k})^{\circ}_{+}$  $^{\circ}_{+}$ *, for all*  $m, k \in \mathbb{N}$ .

The first three assumptions are standard in most existing literature (e.g. [\[BP10,](#page-55-4) [BHLP20,](#page-54-2) [HY24,](#page-56-3) [Lam24\]](#page-57-3)), and the new assumption (A.4) is indeed satisfied by many cases of interest. For instance, (A.4) holds when the set is a singleton of tensor product i.i.d. states, as in the existing quantum AEP; when it consists of the identity operator tensored with the set of all density operators, which is relevant for conditional quantum entropies; or when it is the image of a quantum channel or a set of quantum channels, a scenario that naturally arises in adversarial quantum channel discrimination (see Section [6\)](#page-31-0). It is also worth mentioning that even in cases where assumption (A.4) is not directly met, we can relax the original set to fulfill the required conditions. For example, we can relax the set of all separable states to the Rains set, which is widely used in entanglement theory, or relax the set of stabilizer states to those with non-positive mana, a technique used in fault-tolerant quantum computing. These applications will be discussed in detail in Section [8.](#page-45-0)

Now we proceed to present our main result in this work as follows.

<span id="page-16-0"></span>**Theorem 26.** *(Generalized AEP.) Let*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  be two sequences of sets satisfying As $sumption$  [25](#page-15-1) and  $\mathscr{A}_n \subseteq \mathscr{D}(\mathcal{H}^{\otimes n})$ ,  $\mathscr{B}_n \subseteq \mathscr{H}_+(\mathcal{H}^{\otimes n})$ . Assume moreover that  $D_{\max}(\mathscr{A}_n||\mathscr{B}_n) \le cn$ , for *all*  $n \in \mathbb{N}$  *and a constant*  $c \in \mathbb{R}_+$ *. Then for any*  $\varepsilon \in (0,1)$ *, it holds* 

$$
\lim_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = \lim_{n \to \infty} \frac{1}{n} D_{\mathrm{max},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = D^{\infty}(\mathscr{A} \| \mathscr{B}).
$$
\n(72)

*In addition,*  $D^{\infty}(\mathscr{A}|\mathscr{B})$  *can be estimated using the following bounds: for any*  $m \geq 1$ *, we have* 

<span id="page-16-6"></span><span id="page-16-5"></span><span id="page-16-4"></span><span id="page-16-2"></span>
$$
\frac{1}{m}D_{\mathcal{M}}(\mathscr{A}_m||\mathscr{B}_m) \le D^{\infty}(\mathscr{A}||\mathscr{B}) \le \frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m)
$$
\n(73)

*with explicit convergence guarantees*

$$
\frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) - \frac{1}{m}D_M(\mathscr{A}_m||\mathscr{B}_m) \le \frac{1}{m}2(d^2+d)\log(m+d). \tag{74}
$$

*where*  $d = \dim \mathcal{H}$ *.* 

*If there is a constant* C *such that for any*  $n \ge 1$  *and*  $\rho_n \in \mathscr{A}_n$ ,  $\sigma_n \in \mathscr{B}_n$ ,  $D_{\max}(\rho_n || \sigma_n) \le Cn/4$ *and*  $\log \text{Tr}(\sigma_n) \leq Cn/4$ <sup>[3](#page-16-1)</sup>*we can obtain explicit bounds of the form:* 

<span id="page-16-7"></span>
$$
-C'f(n,\varepsilon) \le D_{H,\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) - nD^{\infty}(\mathscr{A} \| \mathscr{B}) \le C'f(n,1-\varepsilon),\tag{75}
$$

*and*

$$
-C'f(n, 1-2\varepsilon) - \log \frac{1}{\varepsilon} \le D_{\max,\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) - nD^{\infty}(\mathscr{A} \| \mathscr{B}) \le C'f(n, \varepsilon) + \log \frac{2}{\varepsilon^2},\tag{76}
$$

where  $f(n,\varepsilon) = n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon}$  and C' only depends on C and the local dimension d.

<span id="page-16-8"></span>**Remark 2** (Approximating  $D^{\infty}(\mathscr{A}|\mathscr{B})$  efficiently.) Eq. [\(73\)](#page-16-2) provides converging upper and lower bounds on  $D^{\infty}(\mathscr{A}|\mathscr{B})$  in terms of convex optimization programs as both D and  $D_M$  are convex. In addition, if  $\mathcal{A}_m$  and  $\mathcal{B}_m$  are semidefinite representable (i.e., can be written as the feasible sets of a semidefinite program), then the upper bound  $\frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m)$  is a quantum relative entropy program [\[CS17,](#page-55-9) [FF18\]](#page-55-10) that can be solved using interior point methods [\[FS23\]](#page-56-8). For the lower bound  $\frac{1}{m}D_M(\mathscr{A}_m||\mathscr{B}_m)$ , we use Lemma [20](#page-11-2) to write

$$
D_M(\mathscr{A}_m \|\mathscr{B}_m) = \sup_{W,X} \left\{ -h_{\mathscr{A}_m}(X) : W \in (\mathscr{B}_m)_{++}^\circ, X \ge -\log W \right\} \tag{77}
$$

<span id="page-16-3"></span>
$$
= \sup_{W,X,t,s} \left\{ -t+s : t > 0, \ W \in (\mathcal{B}_m)_{++}^\circ, X+sI \in t(\mathcal{A}_m)_{+}^\circ, X \ge -\log W \right\} \tag{78}
$$

<span id="page-16-1"></span>A slightly weaker condition is sufficient, see  $(*)$ .

where we used the fact that  $h_{\mathscr{A}_m}$  is monotone (i.e.,  $h_{\mathscr{A}_m}(X) \leq h_{\mathscr{A}_m}(Y)$  if  $X \leq Y$ ) since  $\mathscr{A}_m \subseteq$  $\mathscr{H}_+$ , and that  $h_{\mathscr{A}_m}(X + sI) = h_{\mathscr{A}_m}(X) + s$  as  $\mathscr{A}_m \subseteq \mathscr{D}$  so that  $h_{\mathscr{A}_m}(X) = \inf_s \{h_{\mathscr{A}_m}(X + sI) - sI\}$  $s: X + sI \geq 0$ } = inf{ $t - s: t > 0$  and  $X + sI \in t\mathcal{A}_{m}^{\circ}, X + sI \geq 0$ }. If  $\mathcal{A}_{m}$  and  $\mathcal{B}_{m}$  are semidefinite representable, then, using semidefinite programming duality, we know that the sets  $(\mathscr{B}_m)_{+}^{\circ}$ and  $\{(t, s, X) : t > 0 \text{ and } X + sI \in t(\mathcal{A}_m)\}^{\circ}$  $_{+}^{\circ}$  are semidefinite representable. The last constraint in [\(78\)](#page-16-3) is convex since the matrix logarithm is operator concave, and the whole maximization problem [\(78\)](#page-16-3) can also be solved using interior-point methods [\[FS23\]](#page-56-8).

Assuming the semidefinite programs describing  $\mathscr{A}_m$ ,  $\mathscr{B}_m$  are of size  $d^{O(m)}$  (polynomial in the Hilbert space dimension), then we can approximate  $D^{\infty}(\mathscr{A}||\mathscr{B})$  within additive  $\delta$  by a relative entropy program of size  $d^{O(d^2/\delta)}$  (using [\(74\)](#page-16-4)). Furthermore as  $\mathscr{A}_m, \mathscr{B}_m$  are invariant under permutations, we have  $\frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) = \frac{1}{m}D(\mathscr{A}_m \cap \mathscr{I}_m||\mathscr{B}_m \cap \mathscr{I}_m)$ , where  $\mathscr{I}_m$  is the set of positive operators on  $\mathcal{H}^{\otimes m}$ that are invariant under permutations. In many settings  $\mathscr{A}_m \cap \mathscr{I}_m$ ,  $\mathscr{B}_m \cap \mathscr{I}_m$  can be described in size polynomial in m (see e.g., [\[FST22\]](#page-56-9)) which leads to programs of size polynomial in  $1/\delta$ .

The proof is provided in the following section.

#### <span id="page-17-0"></span>4.1 Lemmas required to prove the generalized AEP

The proof of Theorem [26](#page-16-0) requires the following lemmas.

<span id="page-17-1"></span>Lemma 27. *(Subadditivity.) Let* D *be a quantum divergence that is additive or subaddtive under tensor product of quantum states. Let*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  *and*  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  *be two sequences of sets satisfying (A.1) and (A.3) in Assumption [25](#page-15-1) and*  $\mathscr{A}_n$  ⊆  $\mathscr{D}(\mathcal{H}^{\otimes n})$ *,*  $\mathscr{B}_n$  ⊆  $\mathscr{H}_+(\mathcal{H}^{\otimes n})$ *, and*  $\mathbb{D}(\mathscr{A}_n|\mathscr{B}_n)$  ≤  $cn$  *for all*  $n \in \mathbb{N}$  *and a constant*  $c \in \mathbb{R}_+$ . Then for any  $m, k \in \mathbb{N}$ , it holds

$$
\mathbb{D}(\mathscr{A}_{m+k}||\mathscr{B}_{m+k}) \le \mathbb{D}(\mathscr{A}_m||\mathscr{B}_m) + \mathbb{D}(\mathscr{A}_k||\mathscr{B}_k). \tag{79}
$$

*Proof.* Note that the assumption  $\mathbb{D}(\mathscr{A}_n|\mathscr{B}_n) \le cn$  ensures that all divergences considered here are finite. For any  $\rho_m \in \mathscr{A}_m$ ,  $\sigma_m \in \mathscr{B}_m$ ,  $\rho_k \in \mathscr{A}_k$  and  $\sigma_k \in \mathscr{B}_k$ , we know that  $\rho_m \otimes \rho_k \in \mathscr{A}_{m+k}$  and  $\sigma_m \otimes \sigma_k \in \mathcal{B}_{m+k}$  by the assumption (A.3). This gives

$$
\mathbb{D}(\mathscr{A}_{m+k}||\mathscr{B}_{m+k}) \le \mathbb{D}(\rho_m \otimes \rho_k || \sigma_m \otimes \sigma_k) \le \mathbb{D}(\rho_m || \sigma_m) + \mathbb{D}(\rho_k || \sigma_k),
$$
\n(80)

where the second inequality follows by the additivity or subadditivity assumption of  $\mathbb{D}$ . As the above holds for any  $\rho_m$ ,  $\sigma_m$ ,  $\rho_k$ ,  $\sigma_k$ , we have

$$
\mathbb{D}(\mathscr{A}_{m+k}||\mathscr{B}_{m+k}) \leq \inf_{\substack{\rho_m \in \mathscr{A}_m \\ \sigma_m \in \mathscr{B}_m}} \mathbb{D}(\rho_m || \sigma_m) + \inf_{\substack{\rho_k \in \mathscr{A}_k \\ \sigma_k \in \mathscr{B}_k}} \mathbb{D}(\rho_k || \sigma_k) = \mathbb{D}(\mathscr{A}_m || \mathscr{B}_m) + \mathbb{D}(\mathscr{A}_k || \mathscr{B}_k), \quad (81)
$$

which completes the proof. □

Note that if a sequence of numbers  $\{a_n\}_{n\in\mathbb{N}}$  satisfies  $a_{n+m} \le a_n + a_m$ , then  $a_n/n$  is convergent [ $Dav07$ , Lemma 4.1.2]. Therefore, if  $D$  satisifies the above assumptions, we have

$$
\mathbb{D}^{\infty}(\mathscr{A}||\mathscr{B}) = \inf_{n>1} \frac{1}{n} \mathbb{D}(\mathscr{A}_n||\mathscr{B}_n).
$$
 (82)

<span id="page-18-2"></span>**Lemma 28.** *(Continuity.)* Let  $\{\mathcal{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathcal{B}_n\}_{n\in\mathbb{N}}$  be two sequences of sets such that  $\mathcal{A}_n \subseteq$  $\mathscr{D}(\mathcal{H}^{\otimes n})$ ,  $\mathscr{B}_n \subseteq \mathscr{H}_+(\mathcal{H}^{\otimes n})$  and  $D_{\max}(\mathscr{A}_n \| \mathscr{B}_n) \leq cn$ , for all  $n \in \mathbb{N}$  and a constant  $c \in \mathbb{R}_+$ . If *these sets satisfy (A.1), (A.3) in Assumption [25,](#page-15-1) it holds*

$$
\inf_{\alpha>1} D_{s,\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}) = D^{\infty}(\mathscr{A} \| \mathscr{B}).
$$
\n(83)

*If these sets satisfy (A.1), (A.4) in Assumption [25,](#page-15-1) it holds*

$$
\sup_{\alpha \in (0,1)} D_{\mathcal{M},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}) = D_{\mathcal{M}}^{\infty}(\mathscr{A} \| \mathscr{B}).
$$
\n(84)

*Proof.* Note that the assumption on  $D_{\text{max}}(\mathscr{A}_n|\mathscr{B}_n)$  ensures that all the divergences considered here are finite. Then the following chain of relations hold

$$
\inf_{\alpha>1} D_{\mathcal{S},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}) = \inf_{\alpha>1} \inf_{n>1} \frac{1}{n} D_{\mathcal{S},\alpha}(\mathscr{A}_n \| \mathscr{B}_n)
$$
\n(85)

$$
= \inf_{\alpha>1} \inf_{n>1} \inf_{\substack{\rho_n \in \mathscr{A}_n \\ \sigma_n \in \mathscr{B}_n}} \frac{1}{n} D_{\mathcal{S},\alpha}(\rho_n \| \sigma_n)
$$
(86)

$$
= \inf_{n>1} \inf_{\substack{\rho_n \in \mathscr{A}_n \\ \sigma_n \in \mathscr{B}_n}} \inf_{\alpha > 1} \frac{1}{n} D_{s,\alpha}(\rho_n \| \sigma_n) \tag{87}
$$

$$
= \inf_{n>1} \inf_{\substack{\rho_n \in \mathscr{A}_n \\ \sigma_n \in \mathscr{B}_n}} \frac{1}{n} D(\rho_n \| \sigma_n)
$$
\n(88)

$$
= \inf_{n>1} \frac{1}{n} D(\mathscr{A}_n | \mathscr{B}_n)
$$
\n(89)

$$
=D^{\infty}(\mathscr{A}||\mathscr{B})\tag{90}
$$

where the first and the last equalities follow from the assumptions  $(A.1)$  and  $(A.3)$  and therefore the subadditivity of  $D_{s,a}(\mathscr{A}_n|\mathscr{B}_n)$  and  $D(\mathscr{A}_n|\mathscr{B}_n)$  from Lemma [27,](#page-17-1) and the fourth equality uses the continuity of the sandwiched Rényi divergence at  $\alpha = 1$  for quantum states.

Similarly, we have the following

$$
\sup_{\alpha \in (0,1)} D_{\mathcal{M},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}) = \sup_{\alpha \in (0,1)} \sup_{n>1} \frac{1}{n} D_{\mathcal{M},\alpha}(\mathscr{A}_n \| \mathscr{B}_n)
$$
(91)

$$
= \sup_{n>1} \sup_{\alpha \in (0,1)} \frac{1}{n} D_{\mathbf{M},\alpha}(\mathscr{A}_n \| \mathscr{B}_n)
$$
(92)

$$
= \sup_{n>1} \frac{1}{n} D_M(\mathscr{A}_n \| \mathscr{B}_n)
$$
\n(93)

<span id="page-18-0"></span>
$$
=D_{\mathcal{M}}^{\infty}(\mathscr{A}||\mathscr{B})\tag{94}
$$

where the first and the last equalities follow from the assumptions  $(A.1)$  and  $(A.4)$  and therefore the su-peradditivity from Lemma [22](#page-14-2) and Lemma [24,](#page-15-2) and the third equality uses the continuity of  $D_{M,\alpha}(\mathscr{A}_n||\mathscr{B}_n)$ <br>at  $\alpha = 1$  from Lemma 23. This completes the proof at  $\alpha = 1$  from Lemma [23.](#page-14-3) This completes the proof.

<span id="page-18-1"></span>**Lemma 29.** *(Asymptotic equivalence.) Let*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  *and*  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  *be two sequences of sets satisfying*  $(A.1)$  and  $(A.2)$  in Assumption [25](#page-15-1) and  $\mathscr{A}_n \subseteq \mathscr{D}(\mathcal{H}^{\otimes n})$ ,  $\mathscr{B}_n \subseteq \mathscr{H}_+(\mathcal{H}^{\otimes n})$  and  $D_{\max}(\mathscr{A}_n||\mathscr{B}_n) \le cn$ , for *all*  $n \in \mathbb{N}$  *and a* constant  $c \in \mathbb{R}_+$ . Let  $d = \dim(\mathcal{H})$  and  $\alpha \in [1/2, \infty)$ *. Then it holds* 

$$
\frac{1}{n}D_{\mathcal{M},\alpha}(\mathscr{A}_n||\mathscr{B}_n) \le \frac{1}{n}D_{\mathcal{S},\alpha}(\mathscr{A}_n||\mathscr{B}_n) \le \frac{1}{n}D_{\mathcal{M},\alpha}(\mathscr{A}_n||\mathscr{B}_n) + \frac{1}{n}2(d^2+d)\log(n+d). \tag{95}
$$

*As a consequence, it holds*

<span id="page-19-0"></span>
$$
D_{\mathbf{M},\alpha}^{\infty}(\mathscr{A}|\mathscr{B})=D_{\mathbf{S},\alpha}^{\infty}(\mathscr{A}|\mathscr{B}).
$$
\n(96)

*Proof.* Since  $D_{\text{M},\alpha}(\rho||\sigma) \leq D_{\text{S},\alpha}(\rho||\sigma)$ , it is clear that

$$
\frac{1}{n}D_{\mathbf{M},\alpha}(\mathscr{A}_n||\mathscr{B}_n) \le \frac{1}{n}D_{\mathbf{S},\alpha}(\mathscr{A}_n||\mathscr{B}_n). \tag{97}
$$

Now we prove the second inequality. Let  $T$  be the twirling operation under permutation group. Then for any  $\rho_n \in \mathscr{A}_n$  and  $\sigma_n \in \mathscr{B}_n$ , we have  $\mathcal{T}(\rho_n) \in \mathscr{A}_n$  and  $\mathcal{T}(\sigma_n) \in \mathscr{B}_n$  by the permutation-invariant assumption (A.2) of  $\{\mathcal{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathcal{B}_n\}_{n\in\mathbb{N}}$ . This gives

$$
D_{\mathcal{S},\alpha}(\mathscr{A}_n|\mathscr{B}_n) \le D_{\mathcal{S},\alpha}(\mathcal{T}(\rho_n)\|\mathcal{T}(\sigma_n))\tag{98}
$$

$$
\leq D_{\mathrm{M},\alpha}(\mathcal{T}(\rho_n) \| \mathcal{T}(\sigma_n)) + 2 \log |\mathrm{spec}(\mathcal{T}(\sigma_n))| \tag{99}
$$

$$
\leq D_{\mathcal{M},\alpha}(\mathcal{T}(\rho_n) \| \mathcal{T}(\sigma_n)) + 2 \log \left[ (n+1)^d (n+d)^{d^2} \right] \tag{100}
$$

$$
\leq D_{\mathcal{M},\alpha}(\rho_n \|\sigma_n) + 2\log\left[ (n+1)^d (n+d)^{d^2} \right],\tag{101}
$$

where the second inequality follows from Lemma [17,](#page-10-2) the third inequality follows from Lemma [18](#page-11-3) and the fact that  $\mathcal{T}(\sigma_n)$  is permutation invariant, the last inequality follows from the data-processing inequality of  $D_M$ . As the above holds for any  $\rho_n \in \mathscr{A}_n$  and  $\sigma_n \in \mathscr{B}_n$ , we have

$$
\frac{1}{n}D_{\mathrm{S},\alpha}(\mathscr{A}_n||\mathscr{B}_n) \le \frac{1}{n}D_{\mathrm{M},\alpha}(\mathscr{A}_n||\mathscr{B}_n) + \frac{2}{n}\log\left[(n+1)^d(n+d)^{d^2}\right].\tag{102}
$$

Finally, noting that  $\log[(n+1)^d(n+d)^{d^2}] \le (d^2+d)\log(n+d)$ , we have the result in Eq. [\(95\)](#page-18-0). Taking lim<sub>n→∞</sub> in Eq. [\(95\)](#page-18-0), we have the relation in Eq. [\(96\)](#page-19-0). □

<span id="page-19-1"></span>**Lemma 30.** *(Finite estimation.)* Let  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  be two sequences of sets satisfying As- $sumption 25$  $sumption 25$  and  $\mathscr{A}_n \subseteq \mathscr{D}(\mathcal{H}^{\otimes n})$ ,  $\mathscr{B}_n \subseteq \mathscr{H}_+(\mathcal{H}^{\otimes n})$  and  $D_{\max}(\mathscr{A}_n \| \mathscr{B}_n) \le cn$ , for all  $n \in \mathbb{N}$  and a *constant*  $c \in \mathbb{R}_+$ *. Let*  $d = \dim(\mathcal{H})$ *. Then it holds* 

$$
\frac{1}{m}D_{\mathcal{M}}(\mathscr{A}_m||\mathscr{B}_m) \le D^{\infty}(\mathscr{A}||\mathscr{B}) \le \frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m). \tag{103}
$$

*Proof.* By the superadditivity of the measured relative entropy in Lemma [24](#page-15-2) and the subadditivity of the Umegaki relative entropy in Lemma [27](#page-17-1) (by choosing  $\alpha = 1$ ), we have

$$
D_M^{\infty}(\mathscr{A}||\mathscr{B}) = \sup_{m>1} \frac{1}{m} D_M(\mathscr{A}_m||\mathscr{B}_m), \qquad (104)
$$

$$
D^{\infty}(\mathscr{A}||\mathscr{B}) = \inf_{m \ge 1} \frac{1}{m} D(\mathscr{A}_m||\mathscr{B}_m).
$$
 (105)

This gives

$$
\frac{1}{m}D_{\mathcal{M}}(\mathscr{A}_{m}||\mathscr{B}_{m}) \leq D_{\mathcal{M}}^{\infty}(\mathscr{A}||\mathscr{B}) = D^{\infty}(\mathscr{A}||\mathscr{B}) \leq \frac{1}{m}D(\mathscr{A}_{m}||\mathscr{B}_{m})
$$
\n(106)

where the equality follows from the asymptotic equivalence in Eq. [\(96\)](#page-19-0) by choosing  $\alpha = 1$ . □

In order to obtain the explicit error bound, we use the following lemma.

<span id="page-20-0"></span>**Lemma 31.** Let  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  and  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  be two sequences of sets satisfying Assumption [25](#page-15-1) and  $\mathscr{A}_n \subseteq$  $\mathscr{D}(\mathcal{H}^{\otimes n})$ ,  $\mathscr{B}_n \subseteq \mathscr{H}_+(\mathcal{H}^{\otimes n})$ . Let  $m \geq 2$  and assume there is a constant  $C > 0$  such that

$$
\text{For all } \alpha \in [1/2, 1]: \ D_{P, 3/2}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) \le \frac{C}{4}m \quad \text{and} \quad \text{for all } \sigma_m \in \mathscr{B}_m: \ \log \text{Tr}(\sigma_m) \le \frac{C}{4}m,
$$

where  $\rho_m^{(\alpha)} \in \mathscr{A}_m, \sigma_m^{(\alpha)} \in \mathscr{B}_m$  satisfy  $D_{P,\alpha}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) = D_{P,\alpha}(\mathscr{A}_m \| \mathscr{B}_m)$ . *For*  $1 - \frac{1}{(2+C)m} < \alpha < 1$ *, we have* 

$$
0 \le D^{\infty}(\mathscr{A}||\mathscr{B}) - \frac{1}{m}D_{\mathrm{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \le (1-\alpha)(2+C)^2m + \frac{2(d^2+d)\log(m+d)}{m},\qquad(107)
$$

and for  $1 < \alpha < 1 + \frac{1}{(2+C)m}$ , we have

$$
0 \leq \frac{1}{m}D_{\mathcal{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m) - D^{\infty}(\mathscr{A}||\mathscr{B}) \leq (\alpha - 1)(2 + C)^2 m + \frac{2(d^2 + d)\log(m + d)}{m}.
$$
 (108)

*Proof.* We start by establishing [\(108\)](#page-20-1). For that, we use the continuity of  $D_{s,\alpha}$  as  $\alpha \to 1^+$  as stated in Lemma [51.](#page-66-0) More specifically,

$$
\frac{1}{m}D_{\mathcal{S},\alpha}(\mathcal{A}_m||\mathcal{B}_m) = \frac{1}{m} \inf_{\substack{\rho_m \in \mathcal{A}_m \\ \sigma_m \in \mathcal{B}_m}} D_{\mathcal{S},\alpha}(\rho_m||\sigma_m)
$$
\n(109)

<span id="page-20-2"></span><span id="page-20-1"></span>
$$
\leq \frac{1}{m} \inf_{\substack{\rho_m \in \mathscr{A}_m \\ \sigma_m \in \mathscr{B}_m}} D(\rho_m \| \sigma_m) + (\alpha - 1)(\log \eta_m)^2 \tag{110}
$$

$$
\leq \frac{1}{m} D(\rho_m^{(1)} \| \sigma_m^{(1)}) + (\alpha - 1)(\log \eta_m^{(1)})^2 \tag{111}
$$

for  $\alpha \in (1, 1 + 1/\log \eta_m)$  where  $\sqrt{\eta_m} = \max(4, 2^{2D_{\text{P},3/2}(\rho_m || \sigma_m)} + 2^{-2D_{\text{P},1/2}(\rho_m || \sigma_m)} + 1)$  and  $\sqrt{\eta_m^{(1)}} =$  $\max(4, 2^{2D_{\text{P},3/2}(\rho_m^{(1)} \| \sigma_m^{(1)})} + 2^{-2D_{\text{P},1/2}(\rho_m^{(1)} \| \sigma_m^{(1)})} + 1).$  The assumption  $(*)$  gives  $D_{\text{P},3/2}(\rho_m^{(1)} \| \sigma_m^{(1)}) \leq \frac{C}{4}m$ and  $-D_{P,1/2}(\rho_m^{(1)} \| \sigma_m^{(1)}) \leq \log \text{Tr}(\sigma_m^{(1)}) \leq \frac{C}{4}m$ . As a result,  $\eta_m^{(1)} \leq (3 \cdot 2^{Cm/2})^2$ , which implies  $\log \eta_m^{(1)} \leq 2\log 3 + Cm \leq (2+C)m$ . Finally, for  $\alpha \in \left(1,1+\frac{1}{(2+C)m}\right)$ , we have

$$
\frac{1}{m}D_{s,\alpha}(\mathscr{A}_m||\mathscr{B}_m) \le \frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) + (\alpha - 1)(2 + C)^2 m
$$
\n(112)

$$
\leq \frac{1}{m}D_M(\mathscr{A}||\mathscr{B}) + \frac{1}{m}2(d^2 + d)\log(m + d) + (\alpha - 1)(2 + C)^2m \tag{113}
$$

$$
\leq D^{\infty}(\mathscr{A}||\mathscr{B}) + \frac{1}{m}2(d^2 + d)\log(m + d) + (\alpha - 1)(2 + C)^2 m \tag{114}
$$

using Lemma [29](#page-18-1) then Lemma [30.](#page-19-1)

Now, we move to the approximation from below [\(107\)](#page-20-2). We have

$$
D^{\infty}(\mathscr{A}||\mathscr{B}) - \frac{1}{m}D_{\mathcal{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \le \frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) - \frac{1}{m}D_{\mathcal{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m)
$$
(115)

$$
+\frac{1}{m}D_{\mathrm{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m)-\frac{1}{m}D_{\mathrm{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m).
$$
 (116)

For the term  $\frac{1}{m}D_{\text{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m) - \frac{1}{m}D_{\text{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m)$ , we use Lemma [29](#page-18-1) and obtain the bound

$$
\frac{1}{m}D_{\mathrm{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m) - \frac{1}{m}D_{\mathrm{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \le 2\frac{(d^2+d)\log(m+d)}{m}.\tag{117}
$$

For the term  $\frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) - \frac{1}{m}D_{\mathrm{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m)$ , we use the continuity bound for  $\alpha \to 1^-$  in Lemma [50.](#page-65-0) For  $\alpha \in (1/2, 1)$ , we use the relation between the sandwiched and Petz Rényi divergences

$$
D_{\mathcal{S},\alpha}(\mathcal{A}_m||\mathcal{B}_m) \ge \inf_{\rho_m \in \mathcal{A}_m, \sigma_m \in \mathcal{B}_m} \alpha D_{\mathcal{P},\alpha}(\rho_m||\sigma_m) - (1-\alpha)\log \operatorname{Tr} \sigma_m \tag{118}
$$

$$
\geq D_{\mathcal{P},\alpha}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)})) - (1 - \alpha)(D_{\mathcal{P},\alpha}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)})) + \frac{C}{4}m) \tag{119}
$$

$$
\geq D(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) - (1 - \alpha) \left( (\log \eta_m^{(\alpha)})^2 + D(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) + \frac{C}{4} m \right),\tag{120}
$$

provided  $\alpha \in (1 - 1/\log \eta_m^{(\alpha)}, 1)$  with  $\sqrt{\eta_m^{(\alpha)}} = \max(4, 1 + 2^{D_{\text{P},3/2}(\rho_m^{(\alpha)} || \sigma_m^{(\alpha)})} + 2^{-D_{\text{P},1/2}(\rho_m^{(\alpha)} || \sigma_m^{(\alpha)})})$ . Again we have  $\log \eta_m^{(\alpha)} \leq 2 \log 3 + Cm$ . As a result, we get

$$
\frac{1}{m}D_{\mathcal{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \ge \frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) - (1-\alpha)\left((2\log 3 + Cm)^2/m + C/2\right) \tag{121}
$$

$$
\geq \frac{1}{m}D(\mathscr{A}_m||\mathscr{B}_m) - (1-\alpha)(2+C)^2m,\tag{122}
$$

for  $m \ge 2$  and where we used Lemma [30.](#page-19-1) □

Now we are ready to prove the generalized AEP in Theorem [26.](#page-16-0)

#### <span id="page-21-0"></span>4.2 Proof of the generalized AEP

We prove the results first for  $D_{H,\varepsilon}$ . The results for  $D_{\max,\varepsilon}$  will follow easily at the end. By using the relation of hypothesis testing relative entropy and the sandwiched Rényi divergence (see Lemma [13\)](#page-9-0), it holds for any  $\varepsilon \in (0,1)$  and  $\alpha > 1$ ,

$$
D_{\mathrm{H},\varepsilon}(\rho\|\sigma) \le D_{\mathrm{S},\alpha}(\rho\|\sigma) + \frac{\alpha}{\alpha - 1}\log\frac{1}{1 - \varepsilon}.\tag{123}
$$

This implies

$$
\frac{1}{n}D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \le \frac{1}{n}D_{\mathrm{S},\alpha}(\mathscr{A}_n||\mathscr{B}_n) + \frac{1}{n}\frac{\alpha}{\alpha-1}\log\frac{1}{1-\varepsilon}.\tag{124}
$$

We start by establishing the upper bound on the asymptotic statement [\(72\)](#page-16-5) for  $D_{\text{H,c}}$ . Taking n go to infinity, we have

$$
\limsup_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \le \limsup_{n \to \infty} \frac{1}{n} D_{\mathrm{S},\alpha}(\mathscr{A}_n \| \mathscr{B}_n) = D_{\mathrm{S},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}).
$$
\n(125)

As the above relation holds for any  $\alpha > 1$ , we can take the infimum of  $\alpha$  on the right-hand side and get

$$
\limsup_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \le \inf_{\alpha > 1} D_{\mathrm{S},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}) = D^{\infty}(\mathscr{A} \| \mathscr{B}),\tag{126}
$$

where the equality follows from the continuity in Lemma  $28$ .

We now establish the explicit upper bound [\(75\)](#page-16-6). We start from [\(124\)](#page-21-1) and first use the subadditivity of  $D_{s,\alpha}$  (Lemma [27\)](#page-17-1) and write for some  $m \geq 2$  to be chosen later and  $n = m\lfloor n/m \rfloor + r$  with  $0 \leq r < m$ :

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \leq \lfloor n/m \rfloor D_{\mathrm{S},\alpha}(\mathscr{A}_m \| \mathscr{B}_m) + D_{\mathrm{S},\alpha}(\mathscr{A}_r \| \mathscr{B}_r) + \frac{\alpha}{\alpha-1} \log \frac{1}{1-\varepsilon}
$$
(127)

<span id="page-21-2"></span>
$$
\leq \lfloor n/m \rfloor D_{\mathrm{s},\alpha}(\mathscr{A}_m \|\mathscr{B}_m) + Cm + \frac{\alpha}{\alpha - 1} \log \frac{1}{1 - \varepsilon}.\tag{128}
$$

<span id="page-21-3"></span><span id="page-21-1"></span>

We now use Lemma [31](#page-20-0) for  $m \ge 2$  and  $\alpha \le 1 + \frac{1}{(2+C)m}$  to get

$$
\frac{1}{m}D_{\mathcal{S},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \le D^\infty(\mathscr{A}||\mathscr{B}) + (\alpha - 1)(2 + C)^2 m + \frac{2(d^2 + d)\log(m + d)}{m}.\tag{129}
$$

Let us now choose  $\alpha - 1 = \frac{8d^2 \log m}{(2+C)^2 m^2}$  $\frac{8d^2 \log m}{(2+C)^2 m^2}$  and assume that  $m \ge \max\left(d, \left(\frac{16d^2}{2+C}\right)^2\right)$  $\left(\frac{16d^2}{2+C}\right)^2$  so that the condition  $\alpha \leq 1 + \frac{1}{(2+C)m}$  is satisfied and  $\log(m+d) \leq 2 \log m$ . With this choice, we have

$$
D_{\mathcal{S},\alpha}(\mathcal{A}_m||\mathcal{B}_m) \le mD^{\infty}(\mathcal{A}||\mathcal{B}) + 16d^2 \log m. \tag{130}
$$

Getting back to  $(128)$ , we get

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \le \lfloor n/m \rfloor m D^{\infty}(\mathscr{A} \| \mathscr{B}) + \lfloor n/m \rfloor 16d^2 \log m + Cm + \frac{2(2+C)^2 m^2}{8d^2 \log m} \log \frac{1}{1-\varepsilon}.
$$
\n(131)

Note that if  $D^{\infty}(\mathscr{A}||\mathscr{B}) \geq 0$ , we have

$$
\lfloor n/m \rfloor m D^{\infty}(\mathscr{A} \|\mathscr{B}) \le n D^{\infty}(\mathscr{A} \|\mathscr{B})
$$
\n(132)

and otherwise,

$$
\lfloor n/m \rfloor m D^{\infty}(\mathscr{A} \|\mathscr{B}) \le n D^{\infty}(\mathscr{A} \|\mathscr{B}) - m D^{\infty}(\mathscr{A} \|\mathscr{B}) \le n D^{\infty}(\mathscr{A} \|\mathscr{B}) + Cm,
$$
 (133)

using the condition  $\frac{1}{n} \log \text{Tr}(\sigma_n) \leq C$  for any  $n \geq 1$  and  $\sigma_n \in \mathcal{B}_n$ . As a result,

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n|\mathscr{B}_n) \le nD^\infty(\mathscr{A}|\mathscr{B}) + \frac{n16d^2}{m}\log m + 2Cm + \frac{2(2+C)^2m^2}{8d^2\log m}\log\frac{1}{1-\varepsilon}.\tag{134}
$$

We now choose  $m = \left(\frac{64d^4n}{(2+C)^2}\right)$  $\frac{1}{(2+C)^2 \log \frac{1}{1-\varepsilon}}$  $\bigg\}$ <sup>1/3</sup> and get  $D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \leq nD^\infty(\mathscr{A}||\mathscr{B}) + C'n^{2/3}\log n\left(\log\frac{1}{1-\varepsilon}\right)$  $1-\varepsilon$  $\setminus$ <sup>1/3</sup>  $(135)$ 

for some constant  $C'$  that only depends on  $d$  and  $C$ .

Now we prove the other direction starting first with the asymptotic statement [\(72\)](#page-16-5). Using the relation of hypothesis testing relative entropy and the Petz Rényi divergence (see Lemma [13\)](#page-9-0), it holds for any  $\varepsilon \in (0,1)$  and  $\alpha \in (0,1)$ ,

$$
D_{\mathrm{H},\varepsilon}(\rho\|\sigma) \ge D_{\mathrm{P},\alpha}(\rho\|\sigma) + \frac{\alpha}{\alpha - 1}\log\frac{1}{\varepsilon} \ge D_{\mathrm{M},\alpha}(\rho\|\sigma) + \frac{\alpha}{\alpha - 1}\log\frac{1}{\varepsilon},\tag{136}
$$

where the second inequality follows from the data-processing inequality of the Petz Rényi divergence. Applying this to two sets of quantum states, we get

<span id="page-22-1"></span><span id="page-22-0"></span>
$$
\frac{1}{n}D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \ge \frac{1}{n}D_{\mathrm{M},\alpha}(\mathscr{A}_n||\mathscr{B}_n) + \frac{1}{n}\frac{\alpha}{\alpha-1}\log\frac{1}{\varepsilon}.\tag{137}
$$

Taking  $n$  to infinity, we get

$$
\liminf_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \ge \liminf_{n \to \infty} \frac{1}{n} D_{\mathrm{M},\alpha}(\mathscr{A}_n \| \mathscr{B}_n) = D_{\mathrm{M},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}).
$$
\n(138)

As the above relation holds for any  $\alpha \in (0, 1)$ , we take the supremum of  $\alpha$  and get

$$
\liminf_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \ge \sup_{\alpha \in (0,1)} D_{\mathrm{M},\alpha}^{\infty}(\mathscr{A} \| \mathscr{B}) = D_{\mathrm{M}}^{\infty}(\mathscr{A} \| \mathscr{B}) = D^{\infty}(\mathscr{A} \| \mathscr{B}) \tag{139}
$$

where the first equality follows from the continuity in Lemma [28](#page-18-2) and the second equality follows from the asymptotic equivalence in Lemma [29.](#page-18-1) Putting Eqs. [\(126\)](#page-21-3) and [\(139\)](#page-22-0) together, we have the asserted result in Eq. [\(72\)](#page-16-5) about the hypothesis testing relative entropy.

We now prove the explicit lower bound statement in [\(75\)](#page-16-6). Starting from [\(137\)](#page-22-1) and using the super-additivity statement in Lemma [24](#page-15-2) with  $n = m/n/m + r$ , we have

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \ge \lfloor n/m \rfloor D_{\mathrm{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m) + D_{\mathrm{M},\alpha}(\mathscr{A}_r||\mathscr{B}_r) + \frac{\alpha}{\alpha-1}\log\frac{1}{\varepsilon} \tag{140}
$$

<span id="page-23-0"></span>
$$
\geq \lfloor n/m \rfloor D_{\mathcal{M},\alpha}(\mathscr{A}_m \|\mathscr{B}_m) - Cm + \frac{\alpha}{\alpha - 1} \log \frac{1}{\varepsilon} \tag{141}
$$

where we used the fact that  $D_{M,\alpha}(\mathscr{A}_r|\mathscr{B}_r) \geq D_{P,0}(\mathscr{A}_r|\mathscr{B}_r) \geq \inf_{\sigma_r \in \mathscr{B}_r} -\log \text{Tr} \sigma_r \geq -Cr$  as per Lemma [16.](#page-10-3)

We now apply Lemma [31](#page-20-0) for  $m \ge 2$  and  $\alpha \ge 1 - \frac{1}{(2+\epsilon)}$  $\frac{1}{(2+C)m}$  to get

$$
\frac{1}{m}D_{\mathcal{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \ge D^{\infty}(\mathscr{A}||\mathscr{B}) - (1-\alpha)(2+C)^2m + \frac{2(d^2+d)\log(m+d)}{m}.
$$
 (142)

Let us now choose  $1 - \alpha = \frac{8d^2 \log m}{(2+C)^2 m^2}$  $\frac{8d^2 \log m}{(2+C)^2 m^2}$  and assume that  $m \ge \max\left(d, \left(\frac{16d^2}{2+C}\right)$  $\left(\frac{16d^2}{2+C}\right)^2$  so that the condition  $\alpha \geq 1 - \frac{1}{(2+\epsilon)}$  $\frac{1}{(2+C)m}$  is satisfied and  $\log(m+d) \leq 2 \log m$ . With this choice, we have

$$
D_{\mathcal{M},\alpha}(\mathscr{A}_m||\mathscr{B}_m) \ge mD^{\infty}(\mathscr{A}||\mathscr{B}) - 16d^2 \log m. \tag{143}
$$

Getting back to [\(140\)](#page-23-0), we get

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n|\mathscr{B}_n) \ge \lfloor n/m \rfloor m D^{\infty}(\mathscr{A}|\mathscr{B}) - \lfloor n/m \rfloor 16d^2 \log m + Cm - \frac{2(2+C)^2 m^2}{8d^2 \log m} \log \frac{1}{\varepsilon}.
$$
 (144)

Note that if  $D^{\infty}(\mathscr{A}|\mathscr{B}) \geq 0$ , we have

$$
\lfloor n/m \rfloor m D^{\infty}(\mathscr{A} \|\mathscr{B}) \ge n D^{\infty}(\mathscr{A} \|\mathscr{B}) - m D^{\infty}(\mathscr{A} \|\mathscr{B}) \ge n D^{\infty}(\mathscr{A} \|\mathscr{B}) - mC \tag{145}
$$

because for all  $n \geq 1$ ,  $D(\mathscr{A}_n||\mathscr{B}_n) \geq D_{P,0}(\mathscr{A}_n||\mathscr{B}_n) \geq \inf_{\sigma_n \in \mathscr{B}_n} -\log \text{Tr}(\sigma_n) \geq -Cn$ . Otherwise, if  $D^{\infty}(\mathscr{A}||\mathscr{B}) < 0$ , then

$$
\lfloor n/m \rfloor m D^{\infty}(\mathscr{A} \|\mathscr{B}) \ge n D^{\infty}(\mathscr{A} \|\mathscr{B})
$$
\n(146)

As a result,

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \ge nD^{\infty}(\mathscr{A} \| \mathscr{B}) - \frac{n16d^2}{m} \log m - 2Cm - \frac{2(2+C)^2 m^2}{8d^2 \log m} \log \frac{1}{\varepsilon}.
$$
 (147)

We now choose  $m = \left(\frac{64d^4n}{(3+{\cal C})^2}\right)^2$  $\sqrt{(2+C)^2 \log \frac{1}{\varepsilon}}$  $\int^{1/3}$  and get

$$
D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \ge nD^{\infty}(\mathscr{A}||\mathscr{B}) - C'n^{2/3}\log n\left(\log\frac{1}{\varepsilon}\right)^{1/3}.\tag{148}
$$

for some constant  $C'$  that only depends on  $d$  and  $C$ .

The analogous statements for  $D_{\text{max},\varepsilon}$  follow directly from known relations between  $D_{\text{max},\varepsilon}$  and  $D_{H,\varepsilon}$ . In fact, we have for any  $\varepsilon \in (0,1)$  and  $\varepsilon' \in (0, 1-\varepsilon)$ ,

$$
D_{\mathrm{H},\varepsilon'}(\rho\|\sigma) + \log\left(1 - \varepsilon - \varepsilon'\right) \le D_{\mathrm{max},\varepsilon}(\rho\|\sigma) \le D_{\mathrm{H},1-\frac{1}{2}\varepsilon^2}(\rho\|\sigma) + \log\left(\frac{2}{\varepsilon^2}\right). \tag{149}
$$

The upper bound is from  $[DKF^+14,$  $[DKF^+14,$  Proposition 4.1] and the lower bound from  $[DMHB13,$  Theorem 11]. Taking the infimum over  $\rho \in \mathscr{A}_n$  and  $\sigma_n \in \mathscr{B}_n$ , and then the limit as  $n \to \infty$ , we get the  $D_{\max,\varepsilon}$ statement in [\(72\)](#page-16-5). Moreover, combining this bound with  $\varepsilon' = 1 - 2\varepsilon$  and [\(75\)](#page-16-6) gives [\(76\)](#page-16-7).

## <span id="page-24-0"></span>5 Application 1: another generalized quantum Stein's lemma

Given many i.i.d. copies of a quantum system described by either the state  $\rho$  or  $\sigma$  (referred to as the null and alternative hypotheses, respectively), what is the optimal measurement to determine the true state? In asymmetric hypothesis testing, the objective is to minimize the probability of mistakenly identifying  $\rho$ as  $\sigma$ , while keeping the probability of identifying  $\sigma$  as  $\rho$  below a small, fixed threshold. The well-known quantum Stein's lemma [\[HP91,](#page-56-1) [NO00\]](#page-57-1) establishes that the asymptotic exponential rate at which this error probability tends to zero is given by the quantum relative entropy between  $\rho$  and  $\sigma$ . This result was later extended to distinguish between an i.i.d. state and a set of quantum states by Brandão and Plenio [[BP10\]](#page-55-4). However, a gap in the original proof was discovered [\[FGW21,](#page-56-10) [BBG](#page-54-5)<sup>+</sup>23, BBG<sup>+</sup>24], which was recently resolved by [\[HY24\]](#page-56-3), with an alternative proof provided by [\[Lam24\]](#page-57-3).

In this section, we propose a new generalization of the quantum Stein's lemma that distinguishes between two sets of quantum states beyond the i.i.d. framework. Our results allows both a composite null hypothesis and composite alternate hypothesis. Moreover, we introduce a new framework of quantum resource theory with partial information and prove that such a theory is reversible under asymptotically resource non-generating operations, identifying the regularized quantum relative entropy between two sets as the "unique" measure of resource in this new framework.

#### <span id="page-24-1"></span>5.1 Operational setting: quantum hypothesis testing between two sets of states

Let us first introduce the operational setting for quantum hypothesis testing between two sets of quantum states. In this setting, a tester draws samples from two sets of quantum states,  $\mathscr{A}_n$  and  $\mathscr{B}_n$ , and performs measurements to determine which set the sample belongs to. The null hypothesis assumes that the sample comes from  $\mathcal{A}_n$ , while the alternative hypothesis assumes it is drawn from  $\mathcal{B}_n$ . An illustration of this task is provided in Figure [1.](#page-24-2) Such a setting provides a very general framework. For instance, it includes the standard quantum Stein's lemma when  $\mathcal{A}_n$  and  $\mathcal{B}_n$  are two singletons of i.i.d. states [\[HP91,](#page-56-1) [NO00\]](#page-57-1). It also encompasses the generalized quantum Stein's lemma [\[HY24,](#page-56-3) [Lam24\]](#page-57-3), where the null hypothesis  $\mathcal{A}_n$  is a singleton of i.i.d. states and  $\mathcal{B}_n$  is a set of quantum states. Additionally, it covers the composite quantum Stein's lemma setting, as explored in [\[BBH21\]](#page-54-1), which deals with mixtures of i.i.d. states in both  $\mathscr{A}_n$  and  $\mathscr{B}_n$ . This framework can further accommodate the correlated states setting, as studied in [\[HMO08,](#page-56-11) [MO15\]](#page-57-10), where  $\mathcal{A}_n$  and  $\mathcal{B}_n$  are particular sets of correlated states on a finite spin chain.

<span id="page-24-2"></span>

Figure 1: An illustration of the quantum hypothesis testing between two sets of quantum states  $\mathscr{A}_n$  and  $\mathscr{B}_n$ . A tester draws samples from the sets and performs quantum measurement to determine which class the sample belongs to. The null hypothesis is that the sample is from  $\mathcal{A}_n$ , which typically represents the bad case, and the alternative hypothesis is that the sample is from  $\mathscr{B}_n$ .

As in standard hypothesis testing, the tester will make two types of errors:

- Type-I error: The sample is from  $\mathcal{A}_n$ , but the tester incorrectly classifies it as coming from  $\mathcal{B}_n$ ,
- Type-II error: The sample is from  $\mathcal{B}_n$ , but the tester incorrectly classifies it as coming from  $\mathcal{A}_n$ .

In general, we take the null hypothesis as the more challenging case and aim to control the type-I error rate within a constant threshold. Under this constraint, the goal is to minimize the type-II error as much as possible. To distinguish between two sets  $\mathscr{A}_n$  and  $\mathscr{B}_n$ , we use a quantum measurement  ${M_n, I - M_n}$ . Since we aim to control the discrimination errors for any state within the given sets, regardless of which one is drawn, the type-I error is defined by

$$
\alpha(\mathscr{A}_n, M_n) := \sup_{\rho_n \in \mathscr{A}_n} \text{Tr}[\rho_n (I - M_n)],\tag{150}
$$

and the type-II error is defined by

<span id="page-25-2"></span>
$$
\beta(\mathcal{B}_n, M_n) := \sup_{\sigma_n \in \mathcal{B}_n} \text{Tr}[\sigma_n M_n]. \tag{151}
$$

The asymmetric setting then seeks to determine the optimal exponent at which the type-II error probability decays, known as the *Stein's exponent*, while keeping the type-I error within a fixed threshold  $\varepsilon$ . Specifically, the goal is to evaluate:

$$
\beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) := \inf_{0 \le M_n \le I} \left\{ \beta(\mathscr{B}_n, M_n) : \alpha(\mathscr{A}_n, M_n) \le \varepsilon \right\}.
$$
 (152)

#### <span id="page-25-0"></span>5.2 Generalized quantum Stein's lemma

Before presenting our generalized quantum Stein's lemma, we first show that the operational quantity  $\beta_{\varepsilon}(\mathscr{A}_n|\mathscr{B}_n)$  in Eq. [\(152\)](#page-25-2) is given by the hypothesis testing relative entropy between two sets of quantum states  $\mathcal{A}_n$  and  $\mathcal{B}_n$ , as mathematically defined in Definition [19.](#page-11-4)

<span id="page-25-1"></span>**Lemma 32.** Let  $\mathscr{A} \subseteq \mathscr{D}$  and  $\mathscr{B} \subseteq \mathscr{H}_+$  be two convex sets. For any  $\varepsilon \in (0,1)$ , it holds

$$
-\log \beta_{\varepsilon}(\mathscr{A} \| \mathscr{B}) = D_{\mathrm{H},\varepsilon}(\mathscr{A} \| \mathscr{B}). \tag{153}
$$

*Proof.* Note that, unless both  $\mathscr A$  and  $\mathscr B$  are singletons, the stated result does not follow trivially from their definitions. This is because the left-hand side represents an operational quantity, while the righthand side is a mathematical generalization. To prove this result, it is sufficient to show that

$$
\beta_{\varepsilon}(\mathscr{A}||\mathscr{B}) = \inf_{0 \le M \le I} \left\{ \sup_{\sigma \in \mathscr{B}} \text{Tr}[\sigma M] : \sup_{\rho \in \mathscr{A}} \text{Tr}[\rho (I - M)] \le \varepsilon \right\} = \sup_{\rho \in \mathscr{A}} \sup_{\sigma \in \mathscr{B}} \beta_{\varepsilon}(\rho || \sigma). \tag{154}
$$

This requires pulling the supremum over  $\mathscr A$  in the condition and the supremum over  $\mathscr B$  in the objective function to the left-hand side of  $inf_{0\leq M\leq I}$ , which is a non-trivial task. First observe that we can write:

$$
\beta_{\varepsilon}(\mathscr{A}, \mathscr{B}) := \inf_{0 \le M \le I} \left\{ \sup_{\sigma \in \mathscr{B}} \text{Tr}[\sigma M] : \sup_{\rho \in \mathscr{A}} \text{Tr}[\rho((1 - \varepsilon)I - M)] \le 0 \right\} \tag{155}
$$

$$
= \inf_{0 \le M \le I} \sup_{\sigma \in \mathcal{B}} \sup_{\rho \in \mathcal{A}} \sup_{z > 0} \text{Tr}[\sigma M] + z \text{Tr}[\rho((1 - \varepsilon) - M)] \tag{156}
$$

since  $\sup_{z>0} \text{Tr}[\sigma M] + z \text{Tr}[\rho((1-\varepsilon)I - M)] = 0$  if  $\text{Tr}[\rho((1-\varepsilon)I - M)] \leq 0$  and  $+\infty$  otherwise. By doing the change of variables  $\rho' = z\rho \in Cone(\mathcal{A})$  where  $Cone(\mathcal{A})$  is the convex cone generated by  $\mathscr A$ , we can write:

$$
\beta_{\varepsilon}(\mathscr{A}, \mathscr{B}) = \inf_{\substack{0 \le M \le I \\ \sigma \in \mathscr{B}}} \sup_{\substack{\rho' \in \text{Cone}(\mathscr{A}) \\ \sigma \in \mathscr{B}}} \text{Tr}[\sigma M] + \text{Tr}[\rho'((1 - \varepsilon)I - M)] \tag{157}
$$

Define

$$
f(M, \rho', \sigma) = \text{Tr}[\sigma M] + z \text{Tr}[\rho'((1 - \varepsilon)I - M)] \tag{158}
$$

which is linear in M for  $(\rho', \sigma)$  fixed, and vice-versa. Furthermore the set  $\{M : 0 \le M \le I\}$  is convex compact, and Cone( $\mathscr A$ )  $\times$   $\mathscr B$  is convex. So Sion's minimax theorem [\[Sio58,](#page-58-11) Corollary 3.3] applies and we can write

$$
\beta_{\varepsilon}(\mathscr{A}, \mathscr{B}) = \sup_{\substack{\rho' \in \text{Cone}(\mathscr{A}) \\ \sigma \in \mathscr{B}}} \inf_{0 \le M \le I} \text{Tr}[\sigma M] + \text{Tr}[\rho'((1 - \varepsilon)I - M)] \tag{159}
$$

$$
= \sup_{\substack{\rho \in \mathcal{A} \\ \sigma \in \mathcal{B}}} \sup_{z > 0} \inf_{0 \le M \le I} \text{Tr}[\sigma M] + z \text{Tr}[\rho((1 - \varepsilon) - M)] \tag{160}
$$

$$
= \sup_{\substack{\rho \in \mathcal{A} \\ \sigma \in \mathcal{B}}} \inf_{0 \le M \le I} \sup_{z > 0} \text{Tr}[\sigma M] + z \text{Tr}[\rho((1 - \varepsilon)I - M)] \tag{161}
$$

$$
= \sup_{\substack{\rho \in \mathcal{A} \\ \sigma \in \mathcal{B}}} \inf_{0 \le M \le I} \{ \text{Tr}[\sigma M] : \text{Tr}[\rho((1 - \varepsilon)I - M)] \le 0 \}
$$
(162)

$$
=\sup_{\substack{\rho\in\mathcal{A}\\ \sigma\in\mathcal{B}}} \beta_{\varepsilon}(\rho,\sigma) \tag{163}
$$

where in the third line we applied Sion's minimax theorem again to interchange the  $\sup_{z>0}$  with the inf<sub>0≤M≤1</sub>. This completes the proof. We can also refer to Appendix [B](#page-62-0) for an alternative proof. □

Due to the above relation between the optimal type-II error  $\beta_{\varepsilon}(\mathscr{A}||\mathscr{B})$  and the hypothesis testing relative entropy  $D_{H,\varepsilon}(\mathscr{A}|\mathscr{B})$ , we can now apply the generalized AEP in Theorem [26](#page-16-0) to derive a generalized quantum Stein's lemma as follows.

<span id="page-26-1"></span>**Theorem 33.** *(Generalized quantum Stein's lemma.) Let*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  *and*  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  *be two sequences of*  $sets$  satisfying Assumption [25](#page-15-1) and  $\mathscr{A}_n \subseteq \mathscr{D}(\mathcal{H}^{\otimes n})$ ,  $\mathscr{B}_n \subseteq \mathscr{H}_+(\mathcal{H}^{\otimes n})$  and  $D_{\max}(\mathscr{A}_n \| \mathscr{B}_n) \leq cn$ , for all  $n \in \mathbb{N}$  *and a constant*  $c \in \mathbb{R}_+$ *. Then for any*  $\varepsilon \in (0,1)$ *, it holds* 

$$
\lim_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = D^{\infty}(\mathscr{A} \| \mathscr{B}). \tag{164}
$$

*Proof.* This is a direct consequence of Theorem [26](#page-16-0) and Lemma [32.](#page-25-1) □

Note that Theorem [33](#page-26-1) is incomparable to the generalized quantum Stein lemma of [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3). It is weaker in the sense that we have the additional assumption on the stability of the polar sets under tensor product (see (A.4) on Assumption [25\)](#page-15-1) for the alternate hypothesis  $\mathcal{B}_n$ , but it is stronger in that it allows for a composite null hypothesis and in addition we can obtain efficient and controlled approximations of the Stein exponent as presented in Theorem [26.](#page-16-0)

#### <span id="page-26-0"></span>5.3 Quantum resource theory with partial information and its reversibility

Quantum resource theory is a framework that finds great success in studying different quantum resource features in recent years (see e.g. [\[CG19\]](#page-55-13) for an introduction). A standard resource theory is built upon a set of *free operations* Ω and a set of *free states*  $\mathcal F$  (in contrast to resource states). Take the entanglement theory as an example,  $\mathscr F$  consists of the separable (unentangled) states, and local operationa and classical communication (LOCC) is a standard choice of  $\Omega$ . In principle,  $\mathscr F$  and  $\Omega$  can be adaptively defined, which give rise to a wide variety of meaningful resource theories, as long as they follow a *golden rule*: any free operation can only map a free state to another free state, i.e.  $\Lambda(\rho) \in \mathscr{F}, \forall \rho \in \mathscr{F}, \forall \Lambda \in \Omega$ .

Whether some resource state can be (approximately) converted to another by certain free operations is a fundamental type of problem in quantum information.

Existing studies on quantum resource theory require precise characterization of the source and the target states, and impose an i.i.d. structure, that is, considering transformation  $\rho^{\otimes n}$  to  $\sigma^{\otimes m}$  and evaluating the rate of transformation  $m/n$ . A standard approach to obtain a precise characterization of the state to be manipulated is quantum tomography, which is a resource-intensive task. In practical scenarios, full knowledge of the source state may be unavailable, and different copies of the sources can exhibit correlation by nature. For instance, when establishing entangled photons over a remote distance in a quantum network, the noise affecting the states can vary over time for different copies of the entangled state, making the precise description of the shared state unclear.

Motivated by this, we propose a new framework of quantum resource theory in this section that works with resource manipulation with partial information. More specifically, we consider the transformation of a quantum state  $\rho_n \in \mathscr{D}(A^{\otimes n})$  to another quantum state  $\sigma_m \in \mathscr{D}(B^{\otimes m})$ . In this case, we do not have complete information about the specific states  $\rho_n$  and  $\sigma_m$ . All we know is that the source state lies within a certain set of quantum states  $\mathcal{A}_n$  (for example, we may know the range of noise affecting the source). Our goal is to design a resource manipulation protocol  $\Lambda$  such that regardless of the specific state in  $\mathscr{A}_n$ , it universally transforms this state into a target state that falls within the target range  $\mathcal{B}_n$ . A comparison of this setting with the standard resource manipulation is depicted in Figure [2.](#page-27-0)

<span id="page-27-0"></span>

Figure 2: Illustration of the standard quantum resource manipulation (left) that transforms an i.i.d. state  $\rho^{\otimes n}$  to another i.i.d. state  $\sigma^{\otimes m}$  and the quantum resource manipulation with partial information (right) that transforms an unknown quantum state in  $\mathcal{A}_n$  to a quantum state in the target range  $\mathcal{B}_m$ .

Let  $\Delta(\rho,\sigma) := \frac{1}{2}$  $\frac{1}{2} \|\rho - \sigma\|_1$  be the trace distance between two quantum states. Consider quantum state transformation from  $\mathscr{A}_n$  to  $\mathscr{B}_m$  via free operation class  $\Omega_n$ , and define the transformation error by

$$
\Delta\left(\mathcal{A}_n \xrightarrow{\Omega_n} \mathcal{B}_m\right) := \inf_{\Lambda_n \in \Omega_n} \sup_{\rho_n \in \mathcal{A}_n} \inf_{\sigma_m \in \mathcal{B}_m} \Delta(\Lambda_n(\rho_n), \sigma_m). \tag{165}
$$

That is, there exists a free operation  $\Lambda_n \in \Omega_n$  such that it universally transforms any state in  $\rho_n \in \mathcal{A}_n$ to a quantum state  $\sigma_m\in\mathscr{B}_m$  within the error  $\Delta\left(\mathscr{A}_n\xrightarrow{\Omega_n}\mathscr{B}_m\right)$ . Based on this notion of error measure, we can introduce the asymptotic rate of transformation by

$$
r\left(\mathscr{A}\xrightarrow{\Omega}\mathscr{B}\right):=\sup\left\{\limsup_{n\to\infty}\frac{m_n}{n}:\lim_{n\to\infty}\Delta\left(\mathscr{A}_n\xrightarrow{\Omega_n}\mathscr{B}_{m_n}\right)=0\right\},\qquad(166)
$$

which is the optimal achievable rate of transformation with vanishing transformation error.

As the previous generalized quantum Stein's lemma [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3) is deeply related to the reversibility of quantum resource manipulation, here we show an analogy result in this new framework. We say a quantum resource theory with partial information is reversible if the asymptotic rate satisfies,

$$
r\left(\mathscr{A}\xrightarrow{\Omega}\mathscr{B}\right)\cdot r\left(\mathscr{B}\xrightarrow{\Omega}\mathscr{A}\right)=1,\tag{167}
$$

which means we can transform from  $\mathscr A$  to  $\mathscr B$  and then back from  $\mathscr B$  to  $\mathscr A$  without compromising any resources. For this, we consider the asymptotically resource non-generating operations RNG introduced in [\[BP10\]](#page-55-4). A quantum operation  $\Lambda$  is called  $\delta$ -resource nongenerating operation if for every free state  $\omega \in \mathscr{F}$ , we have  $\mathcal{R}(\Lambda(\omega)) \leq \delta$ , where

$$
\mathcal{R}(\rho) := \min_{\pi \in \mathcal{D}} \left\{ s \ge 0 : \frac{\rho + s\pi}{1 + s} \in \mathcal{F} \right\}
$$
\n(168)

is the global robustness of  $\rho$ . Simple calculation tells that,  $\log(\mathcal{R}(\rho) + 1) = D_{\max}(\rho||\mathcal{F})$ . Then we denote the set of all  $\delta$ -resource nongenerating operations by  $RNG(\delta)$ . An asymptotically resource nongenerating operation is a sequence of operations  $\{\Lambda_n\}_{n\in\mathbb{N}}$  such that  $\Lambda_n \in \text{RNG}(\delta_n)$  and  $\lim_{n\to\infty} \delta_n =$ 0. With this, the asymptotic rate of transformation under RNG operations is given by

$$
r\left(\mathscr{A}\xrightarrow{\text{RNG}}\mathscr{B}\right) = \sup\left\{\limsup_{n\to\infty}\frac{m_n}{n} : \lim_{n\to\infty}\Delta\left(\mathscr{A}_n\xrightarrow{\text{RNG}(\delta_n)}\mathscr{B}_{m_n}\right) = 0, \lim_{n\to\infty}\delta_n = 0\right\}.
$$
 (169)

The following result gives a precise characterization of this asymptotic rate, which identifies the regularized quantum relative entropy between two sets as the "unique" measure of a resource in the asymptotic limit. It also implies the reversibility (also referred to as the second law) of the resource manipulation under RNG.

**Theorem 34.** Let  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$ ,  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  and  $\{\mathscr{F}_n\}_{n\in\mathbb{N}}$  be threes sequences of sets satisfying Assump-*tion [25](#page-15-1) where*  $\mathscr{A}_n\subseteq\mathscr{D}(\mathcal{H}^{\otimes n})$  and  $\mathscr{B}_n\subseteq\mathscr{D}(\mathcal{H}^{\otimes n})$  are the sets of source states and the sets of target states, respectively, and  $\mathscr{F}_n\subseteq\mathscr{D}(\mathcal{H}^{\otimes n})$  are the sets of free states, which defines the operation class RNG*. Let*  $\kappa := \limsup_{n\to\infty} \frac{1}{n} \max_{\omega \in \mathscr{D}} D(\omega || \mathscr{F}_n) < \infty$  and  $D^{\infty}(\mathscr{B} || \mathscr{F}) > 0$ *. Then it holds* 

$$
r\left(\mathscr{A}\xrightarrow{\text{RNG}}\mathscr{B}\right) = \frac{D^{\infty}(\mathscr{A}||\mathscr{F})}{D^{\infty}(\mathscr{B}||\mathscr{F})}.
$$
\n(170)

*Proof.* The proof applies the generalized AEP in Theorem [26](#page-16-0) and follows a similar argument in [\[BG15\]](#page-54-6) and also a more detailed version in [\[Gou24,](#page-56-12) Chapter 11]. This proof contains the converse and the achievable parts, and we will prove the converse part first.

*1) Proof of the converse part:* Suppose by contradiction that

$$
r\left(\mathscr{A}\xrightarrow{\text{RNG}}\mathscr{B}\right) > \frac{D^{\infty}(\mathscr{A}||\mathscr{F})}{D^{\infty}(\mathscr{B}||\mathscr{F})} + 2\delta,\tag{171}
$$

for some small positive  $\delta$ . By definition, this means in particular that for sufficiently small  $\varepsilon \in (0,1)$ , there exists an achievable rate  $r$  such that

$$
r > \frac{D^{\infty}(\mathscr{A}||\mathscr{F})}{D^{\infty}(\mathscr{B}||\mathscr{F})} + \delta.
$$
 (172)

That is, there exists a sequence  $\{m_n\}_{n\in\mathbb{N}}\subseteq\mathbb{N}$  such that  $r=\lim_{n\to\infty}\frac{m_n}{n}$  and there exists another sequence  $\{\delta_n\}_{n\in\mathbb{N}}\subseteq\mathbb{R}_+$  with a limit of zero such that for every  $n\in\mathbb{N}$ ,

$$
\Delta\left(\mathcal{A}_n \xrightarrow{\text{RNG}(\delta_n)} \mathcal{B}_{m_n}\right) \leq \varepsilon. \tag{173}
$$

This means that there exists  $\Lambda_n \in \text{RNG}(\delta_n)$  such that for any  $\rho_n \in \mathcal{A}_n$ 

$$
\Delta(\Lambda_n(\rho_n), \sigma_{m_n}) \le \varepsilon,\tag{174}
$$

for some  $\sigma_{m_n} \in \mathcal{B}_{m_n}$ . By using the continuity in [\[Gou24,](#page-56-12) Theorem 10.2.2], we have

$$
|D(\Lambda_n(\rho_n)||\mathscr{F}_{m_n}) - D(\sigma_{m_n}||\mathscr{F}_{m_n})| \leq c_{m_n}\varepsilon + (1+\varepsilon)h\left(\frac{\varepsilon}{1+\varepsilon}\right),\tag{175}
$$

where  $c_{m_n} := \max_{\omega \in \mathcal{D}} D(\omega || \mathcal{F}_{m_n})$  and  $h(x)$  is the binary Shannon entropy. This gives

$$
D(\sigma_{m_n} \|\mathscr{F}_{m_n}) \le D(\Lambda_n(\rho_n) \|\mathscr{F}_{m_n}) + c_{m_n} \varepsilon + (1+\varepsilon) h\left(\frac{\varepsilon}{1+\varepsilon}\right). \tag{176}
$$

Since  $D(\mathscr{B}_{m_n}||\mathscr{F}_{m_n}) \leq D(\sigma_{m_n}||\mathscr{F}_{m_n})$  by definition, we get

$$
D(\mathscr{B}_{m_n}||\mathscr{F}_{m_n}) \leq D(\Lambda_n(\rho_n)||\mathscr{F}_{m_n}) + c_{m_n}\varepsilon + (1+\varepsilon)h\left(\frac{\varepsilon}{1+\varepsilon}\right). \tag{177}
$$

Now, let  $\omega_n \in \mathscr{F}_n$  be the optimizer satisfying  $D(\rho_n || \omega_n) = D(\rho_n || \mathscr{F}_n)$ . Then we have

$$
D(\Lambda_n(\rho_n) \| \mathcal{F}_{m_n}) = \min_{\tau_{m_n} \in \mathcal{F}_{m_n}} D(\Lambda_n(\rho_n) \| \tau_{m_n})
$$
\n(178)

<span id="page-29-0"></span>
$$
\leq D(\Lambda_n(\rho_n) \|\Lambda_n(\omega_n)) + \min_{\tau_{m_n} \in \mathscr{F}_{m_n}} D_{\max}(\Lambda_n(\omega_n) \|\tau_{m_n}) \tag{179}
$$

$$
\leq D(\rho_n \| \omega_n) + D_{\max}(\Lambda_n(\omega_n) \| \mathcal{F}_{m_n}) \tag{180}
$$

where the first inequality follows from the triangle inequality  $D(\rho||\sigma) \le D(\rho||\omega) + D_{\text{max}}(\omega||\sigma)$  (see e.g. [\[Gou24,](#page-56-12) Theorem 6.3.3]), and the second inequality follows from the data-processing inequality of quantum relative entropy. Since  $\omega_n$  is a free state and  $\Lambda_n \in \text{RNG}(\delta_n)$ , the global robustness of  $\Lambda_n(\omega_n)$ cannot exceed  $\delta_n$  by definition and in particular  $D_{\max}(\Lambda_n(\omega_n) || \mathscr{F}_{m_n}) \leq \log(1+\delta_n)$ . Therefore,

$$
D(\Lambda_n(\rho_n) \|\mathscr{F}_{m_n}) \le D(\rho_n \|\mathscr{F}_n) + \log(1 + \delta_n), \tag{181}
$$

where we use the optimality assumption of  $\omega_n$  here. Taking this into Eq. [\(177\)](#page-29-0), we get

$$
D(\mathscr{B}_{m_n}||\mathscr{F}_{m_n}) \le D(\rho_n||\mathscr{F}_n) + \log(1+\delta_n) + c_{m_n}\varepsilon + (1+\varepsilon)h\left(\frac{\varepsilon}{1+\varepsilon}\right). \tag{182}
$$

As this holds for any  $\rho_n \in \mathcal{A}_n$ , we have

$$
D(\mathscr{B}_{m_n}||\mathscr{F}_{m_n}) \le D(\mathscr{A}_n||\mathscr{F}_n) + \log(1+\delta_n) + c_{m_n}\varepsilon + (1+\varepsilon)h\left(\frac{\varepsilon}{1+\varepsilon}\right). \tag{183}
$$

Noting that  $\lim_{n\to\infty} m_n/n = r$  and  $\lim_{n\to\infty} \frac{1}{m}$  $\frac{1}{m_n}c_{m_n} \leq \kappa$ , we have

$$
\lim_{n \to \infty} \frac{1}{m_n} D(\mathcal{B}_{m_n} \| \mathcal{F}_{m_n}) \le \lim_{n \to \infty} \frac{n}{m_n} \frac{1}{n} D(\mathcal{A}_n \| \mathcal{F}_n) + \kappa \varepsilon. \tag{184}
$$

That is,

$$
D^{\infty}(\mathscr{B}||\mathscr{F}) \le \frac{1}{r}D^{\infty}(\mathscr{A}||\mathscr{F}) + \kappa \varepsilon. \tag{185}
$$

However, since  $r > \frac{D^{\infty}(\mathscr{A}||\mathscr{F})}{D^{\infty}(\mathscr{B}||\mathscr{F})} + \delta$  for sufficiently small  $\varepsilon \in (0, 1)$ , we get a contradiction that

$$
D^{\infty}(\mathscr{B}||\mathscr{F}) \le \frac{1}{r}D^{\infty}(\mathscr{A}||\mathscr{F}) + \kappa \varepsilon < D^{\infty}(\mathscr{B}||\mathscr{F}),\tag{186}
$$

by taking a small enough  $\varepsilon \in (-\infty, \delta D^{\infty}(\mathscr{B}||\mathscr{F})/\kappa) \cap (0, 1)$ . This concludes the converse part.

*2) Proof of the achievable part:* For any

<span id="page-30-2"></span><span id="page-30-1"></span>
$$
r < \frac{D^{\infty}(\mathscr{A} \|\mathscr{F})}{D^{\infty}(\mathscr{B} \|\mathscr{F})},\tag{187}
$$

we aim to show that r is an achievable rate. For this, we fix  $\varepsilon \in (0,1)$  and denote  $m_n := \lceil nr \rceil$  so that  $\lim_{n\to\infty}m_n/n = r$ . Let  $M_n$  be the optimal test for  $\beta_{\varepsilon}(\mathscr{A}_n||\mathscr{F}_n)$ . That is, for any  $\delta > 0$  (will be determined later) and sufficiently large  $n$ , we have from the generalized AEP in Theorem [26,](#page-16-0)

$$
\sup_{\rho_n \in \mathscr{A}_n} \text{Tr}[\rho_n (I - M_n)] \le \frac{\varepsilon}{2}, \quad \text{and} \quad \sup_{\omega_n \in \mathscr{F}_n} \text{Tr}[\omega_n M_n] \le 2^{-n(D^\infty(\mathscr{A}||\mathscr{F}) - \delta)}.
$$
 (188)

Let  $\sigma_{m_n}^*$  and  $\sigma_{m_n}'$  be the optimizers satisfying

$$
D_{\max,\varepsilon/2}(\mathscr{B}_{m_n}||\mathscr{F}_{m_n}) = D_{\max,\varepsilon/2}(\sigma_{m_n}^*||\mathscr{F}_{m_n}) = D_{\max}(\sigma_{m_n}'||\mathscr{F}_{m_n})
$$
(189)

where  $\Delta(\sigma'_m, \sigma^*_m) \leq \varepsilon/2^4$  $\Delta(\sigma'_m, \sigma^*_m) \leq \varepsilon/2^4$ . Define the sequence of maps

$$
\Lambda_n(X) := \text{Tr}[M_n X] \sigma'_{m_n} + \text{Tr}[(I - M_n)X] \omega_{m_n},\tag{190}
$$

where  $\omega_{m_n} \in \mathscr{F}_{m_n}$  is a free state. Then we can check that these operations give the expected transformation within the error threshold, and they are indeed asymptotically resource nongenerating operations. For this, we can first check that for any  $\rho_n \in \mathcal{A}_n$ ,

$$
\Delta(\Lambda_n(\rho_n), \sigma_{m_n}^*) \le \Delta(\Lambda_n(\rho_n), \sigma_{m_n}') + \Delta(\sigma_{m_n}', \sigma_{m_n}^*)
$$
\n(191)

$$
= (1 - \text{Tr}[M_n \rho_n]) \Delta(\omega_{m_n}, \sigma'_{m_n}) + \Delta(\sigma'_{m_n}, \sigma^*_{m_n})
$$
\n(192)

$$
\leq \varepsilon,\tag{193}
$$

where the first inequality follows from the triangle inequality of the trace distance, the equality follows by evaluating  $\Lambda_n(\rho_n)$  using the definition and the second inequality follows by Eq. [\(188\)](#page-30-1) and the fact that  $\Delta(\omega_{m_n}, \sigma'_{m_n}) \leq 1$ . As this holds for any  $\rho_n \in \mathscr{A}_n$ , we have

$$
\sup_{\rho_n \in \mathscr{A}_n} \inf_{\sigma_{m_n} \in \mathscr{B}_{m_n}} \Delta(\Lambda_n(\rho_n), \sigma_{m_n}) \le \varepsilon. \tag{194}
$$

Then we check  $\Lambda_n \in \text{RNG}(\delta_n)$  with  $\delta_n = 2^{-n\delta}$  and therefore  $\lim_{n\to\infty} \delta_n = 0$  (note that  $\delta$  will be determined later). We have the following

$$
D^{\infty}(\mathscr{B}||\mathscr{F}) = \lim_{n \to \infty} \frac{1}{m_n} D_{\max,\varepsilon/2}(\mathscr{B}_{m_n}||\mathscr{F}_{m_n})
$$
(195)

$$
= \lim_{n \to \infty} \frac{1}{m_n} D_{\max}(\sigma'_{m_n} \| \mathcal{F}_{m_n})
$$
\n(196)

$$
= \lim_{n \to \infty} \frac{n}{m_n} \frac{1}{n} D_{\max}(\sigma'_{m_n} \| \mathcal{F}_{m_n})
$$
\n(197)

<span id="page-30-3"></span>
$$
=\frac{1}{r}\lim_{n\to\infty}\frac{1}{n}D_{\max}(\sigma'_{m_n}||\mathscr{F}_{m_n}),
$$
\n(198)

where the first line follows from the generalized AEP in Theorem [26,](#page-16-0) the second line follows by the optimality assumption of  $\sigma'_{m_n}$  in Eq. [\(189\)](#page-30-2) and the last equality follows since  $\lim_{n\to\infty}m_n/n=r$ . Since  $r < \frac{D^{\infty}(\mathscr{A} \| \mathscr{F})}{D^{\infty}(\mathscr{B} \| \mathscr{F})}$ , there exists  $\delta > 0$  such that  $r < (D^{\infty}(\mathscr{A} \| \mathscr{F}) - 2\delta)/D^{\infty}(\mathscr{B} \| \mathscr{F})$ , or equivalently,

$$
rD^{\infty}(\mathscr{B}||\mathscr{F}) < D^{\infty}(\mathscr{A}||\mathscr{F}) - 2\delta. \tag{199}
$$

<span id="page-30-0"></span>Here we use trace distance for the definition of the smoothed max-relative entropy for convenience.

Taking this into Eq.  $(198)$ , we have for sufficiently large n,

$$
D_{\max}(\sigma'_{m_n} \|\mathscr{F}_{m_n}) \le n(D^{\infty}(\mathscr{A} \|\mathscr{F}) - 2\delta). \tag{200}
$$

That is, the global robustness of  $\sigma'_{m_n}$  satisfies

$$
\mathcal{R}(\sigma'_{m_n}) \le 2^{n(D^{\infty}(\mathscr{A}||\mathscr{F}) - 2\delta)} - 1.
$$
\n(201)

To show that  $\Lambda_n \in \text{RNG}(\delta_n)$ , let  $\eta_n \in \mathscr{F}_n$  and denote by  $t_n := \text{Tr}[M_n \eta_n]$  and  $r_{m_n} := \mathcal{R}(\sigma'_{m_n})$ . From the convexity of global robustness we get

$$
\mathcal{R}(\Lambda_n(\eta_n)) \le t_n R(\sigma'_{m_n}) + (1 - t_n)R(\omega_{m_n})
$$
\n(202)

<span id="page-31-2"></span>
$$
=t_n r_{m_n} \tag{203}
$$

$$
\leq t_n(r_{m_n}+1) \tag{204}
$$

$$
\leq t_n 2^{n(D^{\infty}(\mathscr{A}||\mathscr{F}) - 2\delta)} \tag{205}
$$

$$
\leq 2^{-n\delta} \tag{206}
$$

$$
=\delta_n\tag{207}
$$

where the second line follows because  $\omega_{m_n} \in \mathscr{F}_{m_n}$  ant therefore  $R(\omega_{m_n}) = 0$ , the fourth line follows by Eq. [\(201\)](#page-31-2) and the fifth line follows as  $t_n \leq 2^{-n(D^{\infty}(\mathscr{A}||\mathscr{F})-\delta)}$  by Eq. [\(188\)](#page-30-1). This concludes that  $\Lambda_n \in \text{RNG}(\delta_n)$  and therefore concludes the proof of the achievable part. □

As special cases, the conventional entanglement distillation task—where the set of free states is given by the Rains set—and magic state distillation—where the free states have non-positive mana—both fit within this new framework. Further discussion, including explanations of why these cases satisfy the assumptions of the theorem, can be found in Section [8.](#page-45-0)

## <span id="page-31-0"></span>6 Application 2: adversarial quantum channel discrimination

In this section, we explore adversarial quantum channel discrimination using both adaptive and nonadaptive strategies, deriving the corresponding quantum Stein's lemma for channels. In particular, we will see that the Stein's exponent is determined by a new notion of quantum channel divergence.

#### <span id="page-31-1"></span>6.1 Operational setting: adversarial quantum channel hypothesis testing

Consider a scenario in which a tester is working with an untrusted quantum device that generates a quantum state upon request. This device guarantees that the state is produced by either the quantum channel  $\mathcal N$  (the bad case) or the quantum channel  $\mathcal M$  (the good case). The tester is permitted to request multiple samples from the device and perform measurements to determine which channel is being used. However, because the device is untrusted, it may have access to the environmental system of the channel and also possess an internal memory that correlates with the generated samples, actively misleading the tester and undermining their ability to correctly identify the channel.

The key question is: *How effectively can the tester distinguish between the good and bad cases while playing against the adversary?* Such a setting has been studied by [\[BHLP20,](#page-54-2) Theorem 2] in the classical case.<sup>[5](#page-31-3)</sup> Here we investigate the general quantum case.

More formally, let  $\mathcal{N}_{A\to B}$  or  $\mathcal{M}_{A\to B}$  be two quantum channels and  $U_{A\to BE}$  and  $V_{A\to BE}$  be their Stinspring dialations, respectively, with  $E$  being the environmental system. Denote

$$
\mathcal{U}_{A\to BE}(\cdot) = U_{A\to BE}(\cdot)U_{A\to BE}^{\dagger}, \quad \text{and} \quad \mathcal{V}_{A\to BE}(\cdot) = V_{A\to BE}(\cdot)V_{A\to BE}^{\dagger}.
$$
 (208)

<span id="page-31-3"></span><sup>5</sup> In  $[BHLP20]$ , the problem is formulated in terms of convex sets  $P, Q$  of probability distributions. We find it convenient for a quantum generalization (and operationally well-motivated) to define the sets  $P$  and  $Q$  as the images of all input probability distributions of some physical channels that we denote  $\mathcal N$  and  $\mathcal M$ . This means that the channel  $\mathcal N$  takes as input a description of some  $p \in P$  and outputs a sample from p.

Then we have

<span id="page-32-5"></span>
$$
\mathcal{N}_{A \to B} = \text{Tr}_E \circ \mathcal{U}_{A \to BE}, \quad \text{and} \quad \mathcal{M}_{A \to B} = \text{Tr}_E \circ \mathcal{V}_{A \to BE}.
$$
 (209)

The adversarial quantum channel discrimination proceeds as follows. Suppose the channel is given by U. First, the adversary prepares a quantum state using a channel  $\mathcal{P}^1 \in \text{CPTP}(R_0E_0 : A_1R_1)$ with  $|R_0| = |E_0| = 1$ , and sends system  $A_1$  to the channel U to generate an output state  $U \circ \mathcal{P}^1$ . Then the adversary returns system  $B_1$  to the tester. After this, the adversary performs an internal update  $\mathcal{P}^2\in\mathrm{CPTP}(E_1R_1:A_2R_2)$  and sends system  $A_2$  to the channel to generate another output  $\mathcal{U}\circ\mathcal{P}^2\circ\mathcal{U}\circ\mathcal{P}^1$ and returns system  $B_2$  to the tester. These processes can be repeated n times. Finally, the tester performs a quantum measurement  $\{M_n, I - M_n\}$  on the systems  $B_{[n]} := B_1 \cdots B_n$  in their possession to decide which channel was used to prepare the states. An illustation of this process is depicted in Figure [3.](#page-32-0)

<span id="page-32-0"></span>

Figure 3: Adversarial quantum channel hypothesis testing for three uses of the channel, where  $U$  and  $V$ are the Stinspring dilations of quantum channels N and M respectively,  $P_i$  and  $Q_i$  are internal operations controlled by the adversary,  $\{M_n, I - M_n\}$  is a quantum measurement performed by the tester.

After *n* uses of the channel, the tester obtains an overall state on the systems  $B_{[n]}$  as:

<span id="page-32-1"></span>
$$
\rho[\{\mathcal{P}^i\}_{i=1}^n] := \text{Tr}_{R_nE_n} \prod_{i=1}^n \mathcal{U}_{A_i \to B_iE_i} \circ \mathcal{P}^i_{R_{i-1}E_{i-1} \to A_iR_i}.
$$
\n(210)

Similarly, if the channel is given by  $V$  and the internal operations by the adversary are given by  $Q^i$ , then the overall state on system  $B_{[n]}$  is given by

<span id="page-32-2"></span>
$$
\sigma[\{\mathcal{Q}^i\}_{i=1}^n] := \text{Tr}_{R_n E_n} \prod_{i=1}^n \mathcal{V}_{A_i \to B_i E_i} \circ \mathcal{Q}^i_{R_{i-1} E_{i-1} \to A_i R_i}.
$$
\n(211)

Note that in both cases, the adversary is allowed to perform different updates  $\mathcal{P}^i$  and  $\mathcal{Q}^i$  at each step depending on whether the channel is  $U$  or  $V$ .

Denote the set of all possible output states from Eq. [\(210\)](#page-32-1) by

<span id="page-32-4"></span><span id="page-32-3"></span>
$$
\mathscr{A}_n := \{ \rho[\{\mathcal{P}^i\}_{i=1}^n] : \mathcal{P}^i \in \text{CPTP}(R_{i-1}E_{i-1} : A_iR_i), \forall R_i \}
$$
(212)

and the set of all possible output states from Eq.  $(211)$  by

$$
\mathscr{B}_n := \{ \sigma[\{\mathcal{Q}^i\}_{i=1}^n] : \mathcal{Q}^i \in \text{CPTP}(R_{i-1}E_{i-1} : A_iR_i), \forall R_i \}. \tag{213}
$$

Note that the dimension of the reference systems, which serve as the adversary's internal memory, can be arbitrarily large.

There are specific types of strategies that do not utilize the information from the environmental systems  $E_i$ , and for each update operation  $\mathcal{P}_i$ ,  $\mathcal{Q}_i$  ( $i \geq 2$ ), these strategies simply apply identity operations with  $R_i = A_{i+1} \cdots A_n$ . An illustration of such a strategy is depicted in Figure [4.](#page-33-0) We refer to these simpler strategies as non-adaptive ones and denote the sets of all generated outputs as  $\mathscr{A}'_n$  and  $\mathscr{B}'_n$ . It is clear to see that these two sets are the images of the tensor product channels,

<span id="page-33-1"></span>
$$
\mathscr{A}'_n = \{ \mathcal{N}^{\otimes n}(\rho_n) : \rho_n \in \mathscr{D} \},\tag{214}
$$

<span id="page-33-2"></span>
$$
\mathscr{B}'_n = \{ \mathcal{M}^{\otimes n}(\sigma_n) : \sigma_n \in \mathscr{D} \}. \tag{215}
$$

<span id="page-33-0"></span>

Figure 4: Non-adaptive strategy for the quantum hypothesis testing for three uses of the channel, where U and V are the Stinspring dialations of quantum channels N and M respectively, P and Q are any operations controlled by the adversary, Id is the identity channel,  $R_i = A_{i+1} \cdots A_n$ , and  $\{M_n, I - M_n\}$ is a quantum measurement performed by the tester.

The adversarial quantum channel hypothesis testing is essentially to discriminate between these two sets  $\mathcal{A}_n$  and  $\mathcal{B}_n$ . Define the type-I error and type-II error as follows:

$$
\alpha(\mathscr{A}_n, M_n) := \sup_{\rho_n \in \mathscr{A}_n} \text{Tr}[\rho_n (I - M_n)],\tag{216}
$$

$$
\beta(\mathcal{B}_n, M_n) := \sup_{\sigma_n \in \mathcal{B}_n} \text{Tr}[\sigma_n M_n].
$$
\n(217)

The asymmetric hypothesis testing investigates the optimal exponent of the type-II error probability decays when the type-I error is within a constant threshold  $\varepsilon$ , that is, to evaluate:

$$
\beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) := \inf_{0 \le M_n \le I} \left\{ \beta(\mathscr{B}_n, M_n) : \alpha(\mathscr{A}_n, M_n) \le \varepsilon \right\}.
$$
 (218)

Since non-adaptive strategies are a specific type of adaptive strategy, we have the following inclusions

$$
\mathscr{A}'_n \subseteq \mathscr{A}_n \quad \text{and} \quad \mathscr{B}'_n \subseteq \mathscr{B}_n. \tag{219}
$$

This gives the relations for the type-I and type-II errros by

$$
\alpha(\mathscr{A}'_n, M_n) \le \alpha(\mathscr{A}_n, M_n) \quad \text{and} \quad \beta(\mathscr{B}'_n, M_n) \le \beta(\mathscr{B}_n, M_n). \tag{220}
$$

So we have the general relation that

$$
\beta_{\varepsilon}(\mathscr{A}'_n \| \mathscr{B}'_n) = \inf_{\substack{0 \le M_n \le I \\ \alpha(\mathscr{A}'_n, M_n) \le \varepsilon}} \beta(\mathscr{B}'_n, M_n) \le \inf_{\substack{0 \le M_n \le I \\ \alpha(\mathscr{A}_n, M_n) \le \varepsilon}} \beta(\mathscr{B}_n, M_n) = \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n),\tag{221}
$$

where the inequality holds because the right-hand side represents the infimum of a larger objective value over a smaller feasible set.

#### <span id="page-34-0"></span>6.2 Adversarial quantum Stein's lemma

We will see in this section that the optimal exponent at which  $\beta_{\epsilon}(\mathscr{A}_n||\mathscr{B}_n)$  decays is precisely characterized by a new variant of quantum channel divergence. For this, we introduce the following notions.

Definition 35. *(Minimum output quantum channel divergence.) Let* D *be a quantum divergence between quantum states.* Let  $\mathcal{N} \in \text{CPTP}(A : B)$  *and*  $\mathcal{M} \in \text{CP}(A : B)$ *. Define the corresponding minimum output channel divergence by*

$$
\mathbb{D}^{\text{inf}}(\mathcal{N}||\mathcal{M}) := \inf_{\substack{\rho \in \mathcal{D}(A) \\ \sigma \in \mathcal{D}(A)}} \mathbb{D}(\mathcal{N}_{A \to B}(\rho_A) || \mathcal{M}_{A \to B}(\sigma_A)).
$$
\n(222)

*Define its regularized channel divergence by*

$$
\mathbb{D}^{\inf,\infty}(\mathcal{N}|\mathcal{M}) := \lim_{n \to \infty} \frac{1}{n} \mathbb{D}^{\inf}(\mathcal{N}^{\otimes n}|\mathcal{M}^{\otimes n}). \tag{223}
$$

The minimum output channel divergence captures a worst-case scenario where different test states are chosen to minimize the distinguishability between the given channels. This concept has been previously examined in the context of quantum fidelity in [\[Wat18,](#page-59-1) Definition 3.57]. However, the more general notion and its broader applications have yet to be thoroughly explored in the literature.

<span id="page-34-1"></span>Remark 3 The minimum output channel divergence can be seen as a special case of the divergence between two sets of quantum states. Let  $\mathcal{L}(\mathcal{D}) := \{ \mathcal{L}(\rho) : \rho \in \mathcal{D} \}$  be the image of the set of density operators under the linear map  $\mathcal{L}$ . Then, by definition, we have the following

<span id="page-34-2"></span>
$$
\mathbb{D}^{\inf}(\mathcal{N}|\mathcal{M}) = \mathbb{D}(\mathcal{N}(\mathcal{D})|\mathcal{M}(\mathcal{D})).\tag{224}
$$

**Remark 4** Note that the measured Rényi divergence coincides with the sandwiched Rényi divergence at  $\alpha \in \{1/2, +\infty\}$ . Thus, by applying the superadditivity in Lemma [22](#page-14-2) and the subadditivity in Lemma [27,](#page-17-1) we conclude that the minimum output channel divergence is additive for these cases,

$$
D_{s,1/2}^{\text{inf}}(\mathcal{N}_1 \otimes \mathcal{N}_2 \| \mathcal{M}_1 \otimes \mathcal{M}_2) = D_{s,1/2}^{\text{inf}}(\mathcal{N}_1 \| \mathcal{M}_1) + D_{s,1/2}^{\text{inf}}(\mathcal{N}_2 \| \mathcal{M}_2)
$$
(225)

$$
D_{\max}^{\text{inf}}(\mathcal{N}_1 \otimes \mathcal{N}_2 || \mathcal{M}_1 \otimes \mathcal{M}_2) = D_{\max}^{\text{inf}}(\mathcal{N}_1 || \mathcal{M}_1) + D_{\max}^{\text{inf}}(\mathcal{N}_2 || \mathcal{M}_2), \tag{226}
$$

where the first equality recovers Watrous' result [\[Wat18,](#page-59-1) Corollary 3.60].

With the notion of minimum output channel divergence, we can now present our main theorem for adversarial quantum channel hypothesis testing as follows.

<span id="page-34-3"></span>**Theorem 36.** *(Adversarial quantum Stein's lemma.) Let*  $\mathcal{N} \in \text{CPTP}$  *and*  $\mathcal{M} \in \text{CP}$ *. Let*  $\mathcal{A}_n$  *and*  $\mathcal{B}_n$  *be the sets of quantum states generated by adaptive strategies in Eqs.* [\(212\)](#page-32-3) *and* [\(213\)](#page-32-4). *Similarly, let*  $\mathscr{A}'_n$  *and* 

B′ <sup>n</sup> *be the sets of quantum states generated by non-adaptive strategies in Eqs.* [\(214\)](#page-33-1) *and* [\(215\)](#page-33-2)*. Then for*  $any \varepsilon \in (0,1)$ *, it holds* 

$$
\lim_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = \lim_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}'_n \| \mathscr{B}'_n) = D^{\inf, \infty}(\mathcal{N} \| \mathcal{M}),\tag{227}
$$

*where the Stein's exponent on the right-hand side can be efficiently computed as per Remark [2.](#page-16-8) Consequently, adaptive strategies used by the adversary offer no advantage over non-adaptive ones in adversarial quantum channel hypothesis testing.*

This result is proved by two parts. In the converse part, we prove that

$$
\liminf_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \ge D^{\inf, \infty}(\mathcal{N} \| \mathcal{M}),\tag{228}
$$

by using a new chain rule property of quantum relative entropy. In the achievable part, we prove that

$$
\limsup_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}'_n \| \mathscr{B}'_n) \le D^{\inf, \infty}(\mathcal{N} \| \mathcal{M}),\tag{229}
$$

by using the generalized AEP in Theorem [26.](#page-16-0) The complete proof is presented below.

#### <span id="page-35-0"></span>6.2.1 Proof of the converse part

The proof of the converse part requires the following lemmas.

<span id="page-35-4"></span>**Lemma 37.** *(Convexity.) The sets*  $\{\mathscr{A}_n\}_{n\in\mathbb{N}}$  *and*  $\{\mathscr{B}_n\}_{n\in\mathbb{N}}$  *in Eqs.* [\(212\)](#page-32-3) *and* [\(213\)](#page-32-4) *are convex.* 

*Proof.* We prove the assumptions for  $\{\mathcal{A}_n\}_{n\in\mathbb{N}}$  and the same reasoning works for  $\{\mathcal{B}_n\}_{n\in\mathbb{N}}$  as well. Let  $\{\mathcal{P}^i\}_{i=1}^n$  with systems  $R_1,\ldots,R_n$  and  $\{\bar{\mathcal{P}}^i\}_{i=1}^n$  with systems  $R_1,\ldots,R_n$  be two strategies and  $\lambda \in [0, 1]$ . Note that we may assume both strategies have the same systems  $R_i$  as we can always increase the dimension of the systems  $R_i$  by extending the action of the channel in an arbitrary way without affecting the output. Let us now define another strategy  $\{\bar{\bar{\mathcal{P}}}^i\}_{i=1}^n$  as follows. Let  $\bar{R}_i = R_i \otimes C$  for all  $i = 1, \ldots, n$  where C is a two-dimensional system. Then define

$$
\bar{\bar{\mathcal{P}}}^{1}(\cdot) = \lambda \mathcal{P}^{1}(\cdot) \otimes |0\rangle\langle 0|_{C} + (1 - \lambda)\bar{\mathcal{P}}^{1}(\cdot) \otimes |1\rangle\langle 1|_{C}
$$
\n(230)

and for  $i \geq 2$ , define

$$
\bar{\bar{\mathcal{P}}}^i(X) = (\mathcal{P}^i \circ \mathcal{C}_0(X)) \otimes |0\rangle\langle 0|_C + (\bar{\mathcal{P}}^i \circ \mathcal{C}_1(X)) \otimes |1\rangle\langle 1|_C \tag{231}
$$

where  $C_0(X) = \langle 0 | C X | 0 \rangle_C$  and  $C_1(X) = \langle 1 | C X | 1 \rangle_C$ . Since  $C_0, C_1, \mathcal{P}_2^i \overline{\mathcal{P}}^i$  are all CP maps, we know that  $\bar{\bar{\mathcal{P}}}^i$  is also a CP map. Moreover, if  $\mathcal{P}^i$ ,  $\bar{\mathcal{P}}^i$  are trace-preserving, then  $\bar{\bar{\mathcal{P}}}^i$  is also trace-preserving. It is easy to check the following relations,

<span id="page-35-1"></span>
$$
\mathcal{C}_0 \circ \bar{\bar{\mathcal{P}}}^1 = \lambda \mathcal{P}^1 \qquad \text{and} \quad \mathcal{C}_1 \circ \bar{\bar{\mathcal{P}}}^1 = (1 - \lambda) \bar{\mathcal{P}}^1,\tag{232}
$$

<span id="page-35-3"></span><span id="page-35-2"></span>
$$
\mathcal{C}_0 \circ \bar{\bar{\mathcal{P}}}^i = \mathcal{P}^i \circ \mathcal{C}_0 \quad \text{and} \quad \mathcal{C}_1 \circ \bar{\bar{\mathcal{P}}}^i = \bar{\mathcal{P}}^i \circ \mathcal{C}_1, \quad \forall i \ge 2. \tag{233}
$$

Noting that  $\text{Tr}_C$  commutes with U as they are acting on different systems, we have

$$
\operatorname{Tr}_C \circ \prod_{i=1}^n \mathcal{U} \circ \bar{\bar{\mathcal{P}}}^i = \mathcal{U} \circ \mathcal{P}^n \circ \mathcal{C}_0 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\bar{\mathcal{P}}}^i + \mathcal{U} \circ \bar{\mathcal{P}}^n \circ \mathcal{C}_1 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\bar{\mathcal{P}}}^i. \tag{234}
$$

Also noting that  $C_0$  and  $C_1$  both commute with U as they are acting on different systems and using the relations in Eqs.  $(232)$  and  $(233)$ , we have

$$
\mathcal{U} \circ \mathcal{P}^n \circ \mathcal{C}_0 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i = \lambda \mathcal{U} \circ \mathcal{P}^n \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \mathcal{P}^i = \lambda \prod_{i=1}^n \mathcal{U} \circ \mathcal{P}^i,
$$
\n(235)

$$
\mathcal{U} \circ \bar{\mathcal{P}}^n \circ \mathcal{C}_1 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\bar{\mathcal{P}}}^i = (1 - \lambda) \mathcal{U} \circ \mathcal{P}^n \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i = (1 - \lambda) \prod_{i=1}^n \mathcal{U} \circ \bar{\mathcal{P}}^i.
$$
 (236)

Taking these into Eq. [\(234\)](#page-35-3), we have

$$
\operatorname{Tr}_{\bar{R}_n E_n} \circ \prod_{i=1}^n \mathcal{U} \circ \bar{\bar{\mathcal{P}}}^i = \lambda \operatorname{Tr}_{R_n E_n} \circ \prod_{i=1}^n \mathcal{U} \circ \mathcal{P}^i + (1-\lambda) \operatorname{Tr}_{R_n E_n} \circ \prod_{i=1}^n \mathcal{U} \circ \bar{\mathcal{P}}^i \tag{237}
$$

This shows that any mixture of the reduced states on  $B_{[n]}$  by the strategies  $\{\mathcal{P}^i\}_{i=1}^n$  and  $\{\bar{\mathcal{P}}^i\}_{i=1}^n$  is also given by the reduced state of another strategy  $\{\bar{\bar{\mathcal{P}}}^i\}_{i=1}^n$  which proves the convexity of  $\mathscr{A}_n$ . □

The key technical tool for the proof of the converse part makes use of new chain rule properties of quantum relative entropy. For classical probability distributions  $P_{XY}$  and  $Q_{XY}$  we have the chain rule [\[Cov99,](#page-55-0) Theorem 2.5.3]

$$
D(P_{XY}||Q_{XY}) = D(P_X||Q_X) + \sum_x P_X(x)D(P_{Y|X=x}||Q_{Y|X=x}),
$$
\n(238)

where  $D$  is the Kullback-Leibler divergence. There is no known quantum analogue that exactly corresponds to this chain rule. However, by relaxing the equality to the following inequality

$$
D(P_{XY} \| Q_{XY}) \le D(P_X \| Q_X) + \max_x D(P_{Y|X=x} \| Q_{Y|X=x}),
$$
\n(239)

quantum analogs for the Umegaki relative entropy were found in [\[FFRS20\]](#page-56-4) and for the Belavkin-Staszewski relative entropy in [\[FF21a\]](#page-56-13). We now demonstrate that the chain rule property also holds in the reverse direction, providing the first quantum analog for

$$
D(P_{XY}||Q_{XY}) \ge D(P_X||Q_X) + \min_x D(P_{Y|X=x}||Q_{Y|X=x}).
$$
\n(240)

More specifically, by using the superadditivity in Lemma [24](#page-15-2) we have the following chain rules, which can also be seen as an enhancement of the data processing inequality under partial trace.

<span id="page-36-2"></span>**Lemma 38.** *(Chain rule.) Let*  $\alpha \in [1/2, 1)$ *. For any*  $\rho \in \mathcal{D}$ ,  $\sigma \in \mathcal{H}_+$ ,  $\mathcal{N} \in$  CPTP and  $\mathcal{M} \in$  CP, it holds

$$
D_{\scriptscriptstyle{\mathrm{M}}}(\mathcal{N}_{A\to B}(\rho_{RA}) \| \mathcal{M}_{A\to B}(\sigma_{RA})) \ge D_{\scriptscriptstyle{\mathrm{M}}}(\rho_R \| \sigma_R) + D_{\scriptscriptstyle{\mathrm{M}}}^{\scriptscriptstyle{\mathrm{inf}}}(\mathcal{N} \| \mathcal{M}), \tag{241}
$$

$$
D_{\mathcal{M},\alpha}(\mathcal{N}_{A\to B}(\rho_{RA})||\mathcal{M}_{A\to B}(\sigma_{RA})) \geq D_{\mathcal{M},\alpha}(\rho_R||\sigma_R) + D_{\mathcal{M},\alpha}^{\text{inf}}(\mathcal{N}||\mathcal{M}).
$$
\n(242)

*Proof.* The proof utilizes the superadditivity of the divergence between two sets of quantum states, as established in Lemma [24](#page-15-2) and Lemma [22.](#page-14-2) To this end, we consider the following sets:

<span id="page-36-1"></span><span id="page-36-0"></span>
$$
\mathscr{A}_1 = \{ \rho_R \}, \quad \mathscr{A}_2 = \mathcal{N}(\mathscr{D}), \quad \mathscr{A}_3 = \{ \mathcal{N}_{A \to B}(\rho_{RA}) \}, \tag{243}
$$

$$
\mathscr{B}_1 = \{\sigma_R\}, \quad \mathscr{B}_2 = \mathcal{M}(\mathscr{D}), \quad \mathscr{B}_3 = \{\mathcal{M}_{A \to B}(\sigma_{RA})\},\tag{244}
$$

and verify that they meet the required assumptions. We do this for  $\{\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3\}$  and the same argument works for  $\{\mathscr{B}_1, \mathscr{B}_2, \mathscr{B}_3\}$ . For any  $Y_B \in (\mathscr{A}_2)^\circ_+$  $_{+}^{\circ}$ , we have  $\text{Tr}[Y_B \mathcal{N}(\rho)] \leq 1$  for any  $\rho \in \mathscr{D}(A)$ . This

implies that  $\mathcal{N}^{\dagger}(Y_B) \leq I_A$ , with  $\mathcal{N}^{\dagger}$  being the adjoint map of  $\mathcal{N}$ . Therefore, for any  $X_R \in (\mathcal{A}_1)^{\circ}_{+}$  $\int_{+}^{\circ}$  and  $Y_B \in (\mathcal{A}_2)^\circ_+$  $\int_{+}^{\infty}$ , we have the following equations,

$$
\operatorname{Tr}[(X_R \otimes Y_B) \mathcal{N}_{A \to B}(\rho_{RA})] = \operatorname{Tr}[(X_R \otimes \mathcal{N}^{\dagger}(Y_B)) \rho_{RA}]
$$
\n(245)

$$
\leq \operatorname{Tr}[(X_R \otimes I_A)(\rho_{RA})] \tag{246}
$$

$$
= \text{Tr}[X_R \rho_R] \tag{247}
$$

$$
\leq 1. \tag{248}
$$

This implies that  $X_R \otimes Y_B \in (\mathcal{A}_3)^{\circ}_+$  $\frac{\infty}{4}$  and therefore  $(\mathscr{A}_1)^{\circ}_{+} \otimes (\mathscr{A}_2)^{\circ}_{+} \subseteq (\mathscr{A}_3)^{\circ}_{+}$  $\int_{+}^{\infty}$ . Applying the superadditivity in Lemma [24](#page-15-2) and noting that  $D_M^{\text{inf}}(\mathcal{N}||\mathcal{M}) = D_M(\mathcal{N}(\mathcal{D})||\mathcal{M}(\mathcal{D}))$  (see Remark [3\)](#page-34-1), we have the asserted result in Eq. [\(241\)](#page-36-0). The proof for Eq. [\(242\)](#page-36-1) follows the same by the superadditivity in Lemma [22.](#page-14-2)

Considering the multicopy case, we can get the chain rules for the Umegaki relative entropy and the sandwiched Rényi divergence.

<span id="page-37-0"></span>**Lemma 39.** *(Chain rule.) Let*  $\alpha \in [1/2, 1)$ *. For any*  $\rho \in \mathcal{D}$ ,  $\sigma \in \mathcal{H}_+$ ,  $\mathcal{N} \in$  CPTP and  $\mathcal{M} \in$  CP, it holds

$$
D(\mathcal{N}_{A\to B}(\rho_{RA})||\mathcal{M}_{A\to B}(\sigma_{RA})) \ge D(\rho_R || \sigma_R) + D^{\inf,\infty}(\mathcal{N} || \mathcal{M}),
$$
\n(249)

$$
D_{\mathcal{S},\alpha}(\mathcal{N}_{A\to B}(\rho_{RA})||\mathcal{M}_{A\to B}(\sigma_{RA})) \ge D_{\mathcal{S},\alpha}(\rho_R||\sigma_R) + D_{\mathcal{S},\alpha}^{\text{inf},\infty}(\mathcal{N}||\mathcal{M}).
$$
\n(250)

*Proof.* The result is derived by applying Lemma [38](#page-36-2) to the multicopy channels and states, then taking the regularization on both sides, and finally using Lemma [29.](#page-18-1) ⊓⊔

In the following, we introduce the notion of the amortized minimum output channel divergence and show that it coincides with the regularized divergence, being an analog result for the best-case channel divergence [\[FFRS20\]](#page-56-4). This, in turn, demonstrates the tightness of our chain rule properties.

Similar to the amortized channel divergence used in the existing literature [\[WBHK20\]](#page-59-2), we can define the minimum output version as follows.

**Definition 40.** Let  $\mathbb{D}$  be a quantum divergence between states. Let  $\mathcal{N} \in \text{CPTP}(A : B)$  and  $\mathcal{M} \in \text{CP}(A : B)$ B)*. Then the amortized minimum output channel divergence is defined by*

$$
\mathbb{D}^{\text{inf,amo}}(\mathcal{N}||\mathcal{M}) := \inf_{\substack{\rho \in \mathscr{D}(RA) \\ \sigma \in \mathscr{D}(RA)}} \mathbb{D}(\mathcal{N}_{A \to B}(\rho_{RA}) || \mathcal{M}_{A \to B}(\sigma_{RA})) - \mathbb{D}(\rho_R || \sigma_R). \tag{251}
$$

**Lemma 41.** *Let*  $\alpha \in [1/2, 1)$ *. For any*  $\mathcal{N} \in \text{CPTP}(A : B)$ *,*  $\mathcal{M} \in \text{CP}(A : B)$ *, it holds* 

$$
D^{\inf,\text{amo}}(\mathcal{N}|\mathcal{M}) = D^{\inf,\infty}(\mathcal{N}|\mathcal{M}),\tag{252}
$$

<span id="page-37-1"></span>
$$
D_{\mathbf{S},\alpha}^{\text{inf},\text{amo}}(\mathcal{N}||\mathcal{M}) = D_{\mathbf{S},\alpha}^{\text{inf},\infty}(\mathcal{N}||\mathcal{M}).\tag{253}
$$

*Proof.* We prove the result for the quantum relative entropy and the same argument works for the sandwiched Rényi divergence as well. Note that the chain rule property in Lemma  $39$  is equivalent to  $D^{\inf, \text{amo}}(\mathcal{N}||\mathcal{M}) > D^{\inf,\infty}(\mathcal{N}||\mathcal{M})$ . Now we prove the other direction. For this, we will first show the superadditivity of the amortized divergence under tensor product. That is,

$$
D^{\inf,\text{amo}}(\mathcal{N}_1 \otimes \mathcal{N}_2 \|\mathcal{M}_1 \otimes \mathcal{M}_2) \ge D^{\inf,\text{amo}}(\mathcal{N}_1 \|\mathcal{M}_1) + D^{\inf,\text{amo}}(\mathcal{N}_2 \|\mathcal{M}_2),\tag{254}
$$

for any quantum channels  $\mathcal{N}_1 \in \text{CPTP}(A_1 : B_1), \mathcal{N}_2 \in \text{CPTP}(A_2 : B_2)$ , CP maps  $\mathcal{M}_1 \in \text{CP}(A_1 : B_1)$ ,  $M_2 \in \text{CP}(A_2 : B_2)$ . To see this, let  $(\rho_{RA_1A_2}, \sigma_{RA_1A_2})$  be any feasible solution to the optimization on the left-hand side of Eq. [\(254\)](#page-37-1) and denote its corresponding objective value by

$$
\delta_{12} := D(\mathcal{N}_1 \otimes \mathcal{N}_2(\rho_{RA_1A_2}) \| \mathcal{M}_1 \otimes \mathcal{M}_2(\sigma_{RA_1A_2})) - D(\rho_R \| \sigma_R). \tag{255}
$$

Let  $\omega_{RA_2B_1} = \mathcal{N}_1(\rho_{RA_1A_2})$  and  $\gamma_{RA_2B_1} = \mathcal{M}_1(\sigma_{RA_1A_2})$ . We can check that  $(\rho_{RA_1}, \sigma_{RA_1})$  and  $(\omega_{RA_2B_1}, \gamma_{RA_2B_1})$  are feasible solutions to the amortized divergence on the right-hand side of Eq. [\(254\)](#page-37-1), respectively, with the corresponding objective values by

$$
\delta_1 := D(\mathcal{N}_1(\rho_{RA_1}) \| \mathcal{M}_1(\sigma_{RA_1})) - D(\rho_R \| \sigma_R) \ge D^{\inf, \text{amo}}(\mathcal{N}_1 \| \mathcal{M}_1)
$$
\n(256)

$$
\delta_2 := D(\mathcal{N}_2(\omega_{RA_2B_1}) \| \mathcal{M}_2(\gamma_{RA_2B_1})) - D(\omega_{RB_1} \| \gamma_{RB_1}) \ge D^{\inf, \text{amo}}(\mathcal{N}_2 \| \mathcal{M}_2). \tag{257}
$$

Noting that  $\omega_{RB_1} = \mathcal{N}_1(\rho_{RA_1})$  and  $\gamma_{RB_1} = \mathcal{M}_1(\sigma_{RA_1})$ , we have  $\delta_{12} = \delta_1 + \delta_2$ . This implies

$$
\delta_{12} \ge D^{\inf,\text{amo}}(\mathcal{N}_1 \| \mathcal{M}_1) + D^{\inf,\text{amo}}(\mathcal{N}_2 \| \mathcal{M}_2). \tag{258}
$$

As this holds for any feasible solution  $(\rho_{RA_1A_2}, \sigma_{RA_1A_2})$ , we have the asserted result in Eq. [\(254\)](#page-37-1).

By trivilizing the reference system in the amortized divergence, we get  $D^{\inf, \text{amo}}(\mathcal{N} \mid \mathcal{M}) \leq D^{\inf}(\mathcal{N} \mid \mathcal{M})$ . Then we have

$$
D^{\inf,\text{amo}}(\mathcal{N} \| \mathcal{M}) \le \frac{1}{n} D^{\inf,\text{amo}}(\mathcal{N}^{\otimes n} \| \mathcal{M}^{\otimes n}) \le \frac{1}{n} D^{\inf}(\mathcal{N}^{\otimes n} \| \mathcal{M}^{\otimes n}),\tag{259}
$$

where the first inequality follows from Eq. [\(254\)](#page-37-1). As the above holds for any n, we can take  $n \to \infty$  on the right-hand side and conclude that  $D^{\inf, \text{amo}}(\mathcal{N} \| \mathcal{M}) \leq D^{\inf, \infty}(\mathcal{N} \| \mathcal{M})$ . This completes the proof. □

#### Proof of the converse part

With the above lemmas, we are now ready to prove the converse part, that is, to show

<span id="page-38-0"></span>
$$
\liminf_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \ge D^{\inf, \infty}(\mathcal{N} \| \mathcal{M}). \tag{260}
$$

Using the relation of hypothesis testing relative entropy and the Petz Rényi divergence in Lemma [13](#page-9-0) and the fact that  $D_{S,\alpha}(\rho||\sigma) \leq D_{P,\alpha}(\rho||\sigma)$  [\[Tom16,](#page-58-12) Eq.(4.88)], we have for any  $\alpha \in [1/2, 1)$  and  $\varepsilon \in (0, 1)$ ,

$$
\frac{1}{n}D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \ge \frac{1}{n}D_{\mathrm{S},\alpha}(\mathscr{A}_n||\mathscr{B}_n) + \frac{1}{n}\frac{\alpha}{\alpha-1}\log\frac{1}{\varepsilon}.\tag{261}
$$

Denote the joint states before the  $n$ -th use of the channel by

$$
\rho'_{n} := \mathcal{P}_{R_{n-1}E_{n-1}\to A_{n}R_{n}}^{n} \circ \prod_{i=1}^{n-1} \mathcal{U}_{A_{i}\to B_{i}E_{i}} \circ \mathcal{P}_{R_{i-1}E_{i-1}\to A_{i}R_{i}}^{i},
$$
\n(262)

$$
\sigma'_{n} := \mathcal{Q}_{R_{n-1}E_{n-1}\to A_nR_n}^{n} \circ \prod_{i=1}^{n-1} \mathcal{V}_{A_i\to B_iE_i} \circ \mathcal{Q}_{R_{i-1}E_{i-1}\to A_iR_i}^{i}.
$$
 (263)

Then we have

$$
\rho[\{\mathcal{P}^i\}_{i=1}^n] = \text{Tr}_{R_nE_n} \circ \mathcal{U}_{A_n \to B_nE_n}(\rho'_n),\tag{264}
$$

$$
\sigma[\{\mathcal{Q}^i\}_{i=1}^n] = \text{Tr}_{R_nE_n} \circ \mathcal{V}_{A_n \to B_nE_n}(\sigma'_n). \tag{265}
$$

By the relation in Eqs.  $(209)$ , we have

$$
D_{s,\alpha}\left(\rho[\{\mathcal{P}^i\}_{i=1}^n]||\sigma[\{\mathcal{Q}^i\}_{i=1}^n]\right) = D_{s,\alpha}(\mathcal{N}_{A_n \to B_n}(\text{Tr}_{R_n}(\rho'_n))||\mathcal{M}_{A_n \to B_n}(\text{Tr}_{R_n}(\sigma'_n))).
$$
 (266)

By using the chain rule in Lemma [39,](#page-37-0) we get

$$
D_{\mathcal{S},\alpha}\left(\rho[\{\mathcal{P}^i\}_{i=1}^n] \|\sigma[\{\mathcal{Q}^i\}_{i=1}^n]\right) \ge D_{\mathcal{S},\alpha}(\mathrm{Tr}_{A_nR_n}(\rho'_n) \|\mathrm{Tr}_{A_nR_n}(\sigma'_n)) + D_{\mathcal{S},\alpha}^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M}).\tag{267}
$$

Note that for any  $\mathcal{P}^n$  and  $\mathcal{Q}^n$  we always have

<span id="page-39-1"></span>
$$
\text{Tr}_{A_n R_n} \circ \mathcal{P}_{R_{n-1} E_{n-1} \to A_n R_n}^n = \text{Tr}_{E_{n-1} R_{n-1}},\tag{268}
$$

<span id="page-39-2"></span>
$$
\text{Tr}_{A_n R_n} \circ \mathcal{Q}_{R_{n-1} E_{n-1} \to A_n R_n}^n = \text{Tr}_{E_{n-1} R_{n-1}}.
$$
\n(269)

This gives

$$
D_{s,\alpha}(\text{Tr}_{A_nR_n}(\rho'_n) \| \text{Tr}_{A_nR_n}(\sigma'_n)) = D_{s,\alpha} \left( \rho [\{\mathcal{P}^i\}_{i=1}^{n-1}] \| \sigma [\{\mathcal{Q}^i\}_{i=1}^{n-1}] \right). \tag{270}
$$

Combining Eqs.  $(267)$  and  $(270)$ , we have

$$
D_{s,\alpha}\left(\rho[\{\mathcal{P}^i\}_{i=1}^n] \| \sigma[\{\mathcal{Q}^i\}_{i=1}^n] \right) \ge D_{s,\alpha}\left(\rho[\{\mathcal{P}^i\}_{i=1}^{n-1}] \| \sigma[\{\mathcal{Q}^i\}_{i=1}^{n-1}] \right) + D_{s,\alpha}^{\text{inf},\infty}(\mathcal{N} \| \mathcal{M}).\tag{271}
$$

Using this relation  $n$  times, we get

$$
D_{\mathbf{S},\alpha}\left(\rho[\{\mathcal{P}^i\}_{i=1}^n] \|\sigma[\{\mathcal{Q}^i\}_{i=1}^n]\right) \ge n D_{\mathbf{S},\alpha}^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M}).\tag{272}
$$

As this relation holds for any update operations  $\mathcal{P}^i$  and  $\mathcal{Q}^i$ , we have

$$
D_{\mathcal{S},\alpha}(\mathcal{A}_n||\mathcal{B}_n) \ge n D_{\mathcal{S},\alpha}^{\inf,\infty}(\mathcal{N}||\mathcal{M}).
$$
\n(273)

Taking this into Eq.  $(261)$ , we have

$$
\frac{1}{n}D_{\mathrm{H},\varepsilon}(\mathscr{A}_n||\mathscr{B}_n) \ge D_{\mathrm{S},\alpha}^{\mathrm{inf},\infty}(\mathcal{N}||\mathcal{M}) + \frac{1}{n}\frac{\alpha}{\alpha-1}\log\frac{1}{\varepsilon}.\tag{274}
$$

By the convexity in Lemma [37](#page-35-4) and the operational interpretation in Lemma [32,](#page-25-1) we have

$$
\liminf_{n \to \infty} -\log \frac{1}{n} \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) = \liminf_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n)
$$
\n(275)

$$
\geq D_{\mathbf{S},\alpha}^{\text{inf},\infty}(\mathcal{N}||\mathcal{M})\tag{276}
$$

$$
\geq D_{\mathrm{M},\alpha}^{\mathrm{inf},\infty}(\mathcal{N}||\mathcal{M}),\tag{277}
$$

where the last inequality follows since  $D_{\text{M},\alpha}(\rho||\sigma) \leq D_{\text{S},\alpha}(\rho||\sigma)$  for  $\alpha \in [1/2,1)$ . As this holds for any  $\alpha \in [1/2, 1)$ , we get

$$
\liminf_{n \to \infty} -\log \frac{1}{n} \beta_{\varepsilon}(\mathscr{A}_n \| \mathscr{B}_n) \ge \sup_{\alpha \in [1/2, 1)} D_{\mathcal{M}, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M})
$$
\n(278)

$$
=D_{\mathcal{M}}^{\inf,\infty}(\mathcal{N}||\mathcal{M})\tag{279}
$$

$$
=D^{\inf,\infty}(\mathcal{N}||\mathcal{M})\tag{280}
$$

where the first equality follows from the continuitiy in Lemma [28](#page-18-2) and the second equality follows from the asymptotic equivalence in Lemma [29](#page-18-1) by considering the image sets of the channels (note that checking Assumption [25](#page-15-1) for the image sets can be found in the following Lemma [42\)](#page-40-1). This completes the proof of the converse part.

## <span id="page-39-0"></span>6.2.2 Proof of the achievable part

<span id="page-40-1"></span>**Lemma 42.** *The sets*  $\{\mathscr{A}'_n\}_{n\in\mathbb{N}}$  *and*  $\{\mathscr{B}'_n\}_{n\in\mathbb{N}}$  *in Eqs.* [\(214\)](#page-33-1) *and* [\(215\)](#page-33-2) *satisfy Assumption [25.](#page-15-1)* 

*Proof.* First, the set of all density matrices  $\mathscr{D}$  is convex and compact, so  $\mathscr{A}'_n$  is also convex and compact. Since  $\mathcal{N}^{\otimes n}$  and  $\mathscr{D}$  are permutation invariant, we know that  $\mathscr{A}'_n$  is also permutation invariant. For any  $\mathcal{N}^{\otimes m}(\rho_m) \in \mathscr{A}'_m$  and  $\mathcal{N}^{\otimes k}(\rho_k) \in \mathscr{A}'_k$ , we have  $\mathcal{N}^{\otimes m}(\rho_m) \otimes \mathcal{N}^{\otimes k}(\rho_k) = \mathcal{N}^{\otimes (m+k)}(\rho_m \otimes \rho_k) \in \mathscr{A}'_{m+k}$ . This implies  $\mathscr{A}'_m \otimes \mathscr{A}'_k \subseteq \mathscr{A}'_{m+k}$ . The support function of  $\mathscr{A}'_n$  is given by

$$
h_{\mathscr{A}'_n}(X_n) = \sup_{\rho_n \in \mathscr{D}} \text{Tr}\left[X_n \mathcal{N}^{\otimes n}(\rho_n)\right] = \sup_{\rho_n \in \mathscr{D}} \text{Tr}\left[(\mathcal{N}^{\otimes n})^{\dagger}(X_n)\rho_n\right] = \lambda_{\max}\left((\mathcal{N}^{\otimes n})^{\dagger}(X_n)\right). \tag{281}
$$

So for any  $X_m \in \mathcal{H}_+$  and  $X_k \in \mathcal{H}_+$ , we have

$$
h_{\mathscr{A}'_{m+k}}(X_m \otimes X_k) = \lambda_{\max}\left( (\mathcal{N}^{\otimes (m+k)})^{\dagger} (X_m \otimes X_k) \right) \tag{282}
$$

$$
= \lambda_{\max}\left( (\mathcal{N}^{\otimes m})^{\dagger}(X_m) \otimes (\mathcal{N}^{\otimes k})^{\dagger}(X_k) \right) \tag{283}
$$

$$
= \lambda_{\max}\left( (\mathcal{N}^{\otimes m})^{\dagger}(X_m) \right) \lambda_{\max}\left( (\mathcal{N}^{\otimes k})^{\dagger}(X_k) \right) \tag{284}
$$

$$
=h_{\mathscr{A}'_m}(X_m)h_{\mathscr{A}'_k}(X_k),\tag{285}
$$

where the third line follows by the multiplicativity of the maximum eigenvalue of tensor product oper-ators. Finally, using Lemma [8,](#page-8-2) we know that  $(\mathscr{A}'_m)^{\circ}$   $\otimes$   $(\mathscr{A}'_k)^{\circ}$   $\subseteq$   $(\mathscr{A}'_{m+k})^{\circ}$ <sup>o</sup>. This proves that  $\{\mathscr{A}'_n\}_{n\in\mathbb{N}}$ satisfy all assumptions in Assumption [25.](#page-15-1) The same argument works for  $\{\mathscr{B}'_n\}_{n\in\mathbb{N}}$ . □

#### Proof of the achievable part

Since the sets  $\{\mathscr{A}_n'\}_{n\in\mathbb{N}}$  and  $\{\mathscr{B}_n'\}_{n\in\mathbb{N}}$  satisfy Assumption [25,](#page-15-1) we can apply the generalized AEP in Theorem [26](#page-16-0) and get

$$
\limsup_{n \to \infty} -\frac{1}{n} \log \beta_{\varepsilon}(\mathscr{A}'_n \| \mathscr{B}'_n) = \limsup_{n \to \infty} \frac{1}{n} D_{\mathrm{H},\varepsilon}(\mathscr{A}'_n \| \mathscr{B}'_n)
$$
\n(286)

$$
= \lim_{n \to \infty} \frac{1}{n} D(\mathscr{A}'_n | \mathscr{B}'_n)
$$
\n(287)

$$
= \lim_{n \to \infty} \frac{1}{n} D^{\inf} (\mathcal{N}^{\otimes n} \| \mathcal{M}^{\otimes n})
$$
\n(288)

$$
=D^{\inf,\infty}(\mathcal{N}||\mathcal{M}),\tag{289}
$$

where the first equality follows from Lemma [32,](#page-25-1) the second equality follows by applying the generalized AEP in Theorem [26,](#page-16-0) the third equality follows from the relation in Eq. [\(224\)](#page-34-2) and the last equality follows by definition. This completes the proof of the achievable part.

#### <span id="page-40-0"></span>6.3 Justification of the regularization in the Stein's exponent

We provide an explicit example to demonstrate that the minimum output Umegaki channel divergence can be strictly subadditive under tensor product channels. This shows that the regularization in the adversarial quantum Stein's lemma, as stated in Theorem [36,](#page-34-3) is necessary in general. Nevertheless, our example shows that the regularized quantity can be approximated by  $D^{\inf}(\mathcal{N}^{\otimes m}||\mathcal{M}^{\otimes m})/m$  from above and  $D_M^{\inf}(\mathcal{N}^{\otimes m}||\mathcal{M}^{\otimes m})/m$  from below and the approximation gets improved by increasing m.

This example is given by two qutrit quantum channels. Let  $\mathcal{N}(\cdot) = \text{Tr}[\cdot || \rho \rangle \langle \rho |$  to be the replacer channel with  $|\rho\rangle = (2|0\rangle + |1\rangle + 2|2\rangle)/3$ . Let M be the platypus channel [\[LLS](#page-57-11)<sup>+</sup>23, Eq. (170)],  $\mathcal{M}(X) =$   $M_0 X M_0^\dagger + M_1 X M_1^\dagger$  with Kraus operators

$$
M_0 = \begin{bmatrix} \sqrt{s} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \qquad M_1 = \begin{bmatrix} 0 & 0 & 0 \\ \sqrt{1 - s} & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.
$$
 (290)

Since  $\mathcal{M}(I/3) = (s|0\rangle\langle 0| + (1-s)|1\rangle\langle 1| + 2|2\rangle\langle 2|)/3$  is a full rank state for all  $s \in (0, 1)$ , the following Since  $\mathcal{M}(1/3) = (s|0\rangle\langle 0| + (1-s)|1\rangle\langle 1| + 2|2\rangle\langle 2|)/3$  is a full failt state for a quantities are finite and can be evaluated by the QICS package [\[HSF24\]](#page-56-14),

$$
D^{\inf}(\mathcal{N}^{\otimes m}||\mathcal{M}^{\otimes m}) = \inf_{\sigma_m \in \mathcal{D}} D(\rho^{\otimes m}||\mathcal{M}^{\otimes m}(\sigma_m)),\tag{291}
$$

$$
D_{\mathcal{M}}^{\inf}(\mathcal{N}^{\otimes m}||\mathcal{M}^{\otimes m}) = \inf_{\sigma_m \in \mathcal{D}} D_{\mathcal{M}}(\rho^{\otimes m}||\mathcal{M}^{\otimes m}(\sigma_m)).
$$
\n(292)

The numerical result is given in Figure [5.](#page-41-1) It shows a clear separation between  $D^{\inf}(\mathcal{N}^{\otimes m}||\mathcal{M}^{\otimes m})/m$ with  $m = 1, 2, 3$  and  $s \in [0.01, 0.1]$  and also  $D_M^{\text{inf}}(\mathcal{N}^{\otimes m}||\mathcal{M}^{\otimes m})/m$  with the same parameter range, confirming the strict subadditivity of the minimum output Umegaki channel divergence and the strict superadditivity of the minimum output measured channel divergence. Moreover, as we increase the number of m, the lower and upper bounds provide better approximation to  $D^{\infty}(N||\mathcal{M})$ .

Hierarchy for evaluating  $D^{\infty}(\mathcal{N}||\mathcal{M})$ 

<span id="page-41-1"></span>

Figure 5: Nonadditivity of the minimum output channel divergences.

As a remark, if the adversary uses tensor product states as channel inputs, we can similarly prove that the Stein's exponent is given by  $D^{\text{inf}}(\mathcal{N} \| \mathcal{M})$ . Therefore, the strict subadditivity implies that correlations in the adversary's input can indeed help to reduce distinguishability against the tester. However, as stated in Theorem [36,](#page-34-3) such an advantage cannot be further enhanced if the adversary uses more complicated adaptive strategies.

## <span id="page-41-0"></span>7 Application 3: a relative entropy accumulation theorem

Consider two states  $\rho_{R_0}$  and  $\sigma_{R_0}$  and quantum operations  $\mathcal{N}_i \in \text{CPTP}(R_{i-1} : R_i A_i)$  and  $\mathcal{M}_i \in$  $\mathbb{CP}(R_{i-1} : R_i A_i)$  that are applied sequentially from  $i = 1$  to  $i = n$  and generating the systems  $A_i$ . The systems R<sup>i</sup> should be seen as an internal memory system that we do not control. *Can we bound the operationally relevant divergence between the obtained states as the sum of the contributions of each* *step?* Here, more specifically, we consider the hypothesis testing divergence, which can be seen as a smoothed  $D_{\text{min}}$  divergence, and we are looking for a lower bound of the form:

$$
D_{\mathrm{H},\varepsilon}\left(\mathrm{Tr}_{R_n}\circ\prod_{i=1}^n\mathcal{N}_i(\rho_{R_0})\middle\|\mathrm{Tr}_{R_n}\circ\prod_{i=1}^n\mathcal{M}_i(\sigma_{R_0})\right)\geq\sum_{i=1}^n\mathbb{D}(\mathrm{Tr}_{R_i}\circ\mathcal{N}_i\Vert\mathrm{Tr}_{R_i}\circ\mathcal{M}_i)-o(n),\quad(293)
$$

for some appropriate divergence  $D$  between quantum operations. An illustration of this setting is depicted in Figure [6.](#page-42-0) A dual question for smoothed  $D_{\text{max}}$  was raised as an open question in [\[MFSR22\]](#page-57-5).

<span id="page-42-0"></span>

Figure 6: Illustration of the setting for the relative entropy accumulation theorem.

<span id="page-42-2"></span>**Theorem 43.** *(Relative entropy accumulation.) Let*  $\rho_{R_0} \in \mathcal{D}(R_0)$ ,  $\sigma_{R_0} \in \mathcal{D}(R_0)$  *and quantum operations*  $\mathcal{N}_i \in \text{CPTP}(R_{i-1} : R_iA_i)$  *and*  $\mathcal{M}_i \in \text{CP}(R_{i-1} : R_iA_i)$  *for*  $i \in \{1, \ldots, n\}$ *. Let*  $\varepsilon \in (0,1)$  *and assume for all*  $i \in \{1, \ldots, n\}$  *and*  $m \ge 2$ 

$$
\forall \alpha \in [1/2, 1]: \ D_{P, 3/2}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) \le \frac{C}{4}m \text{ and } \forall \sigma \in \mathscr{D}(R_{i-1}): \ \log \text{Tr}(\mathcal{M}_i(\sigma)) \le \frac{C}{4},\tag{294}
$$

where  $\rho_m^{(\alpha)},\sigma_m^{(\alpha)}$  are outputs of the channels  $(\text{Tr}_{R_i}\circ \mathcal{N}_i)^{\otimes m},(\text{Tr}_{R_i}\circ \mathcal{M}_i)^{\otimes m}$  (respectively) and achieve the minimum  $D_{P,\alpha}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) = D_{P,\alpha}^{\text{inf}}((\text{Tr}_{R_i} \circ \mathcal{N}_i)^{\otimes m} \|(\text{Tr}_{R_i} \circ \mathcal{M}_i)^{\otimes m}).$ 

*Then we have*

<span id="page-42-3"></span><span id="page-42-1"></span>
$$
D_{\mathrm{H},\varepsilon} \left( \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{R_0}) \middle\| \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{R_0}) \right) \tag{295}
$$

$$
\geq \sum_{i=1}^{n} D^{\inf,\infty}(\operatorname{Tr}_{R_i} \circ \mathcal{N}_i || \operatorname{Tr}_{R_i} \circ \mathcal{M}_i) - C'n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon},\tag{296}
$$

where  $C'$  is a constant that only depends on  $d = \max \dim A_i$  and  $C$ .

*Proof.* We start as usual by bounding the hypothesis testing relative entropy with a Rényi divergence of order  $\alpha \in (0,1)$ :

$$
D_{\mathrm{H},\varepsilon} \left( \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{R_0}) \middle\| \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{R_0}) \right) \tag{297}
$$

$$
\geq D_{\mathcal{S},\alpha} \left( \operatorname{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{R_0}) \middle\| \operatorname{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{R_0}) \right) + \frac{\alpha}{\alpha - 1} \log(1/\varepsilon). \tag{298}
$$

We can now use the chain rule in Lemma [39](#page-37-0) to bound

$$
D_{\mathrm{S},\alpha}\left(\mathrm{Tr}_{R_n}\circ\prod_{i=1}^n\mathcal{N}_i(\rho_{R_0})\middle\|\mathrm{Tr}_{R_n}\circ\prod_{i=1}^n\mathcal{M}_i(\sigma_{R_0})\right) \tag{299}
$$

$$
\geq D_{\mathbf{S},\alpha} \left( \operatorname{Tr}_{R_{n-1}} \circ \prod_{i=1}^{n-1} \mathcal{N}_i(\rho_{R_0}) \middle\| \operatorname{Tr}_{R_{n-1}} \circ \prod_{i=1}^{n-1} \mathcal{M}_i(\sigma_{R_0}) \right) + D_{\mathbf{S},\alpha}^{\text{inf},\infty} (\operatorname{Tr}_{R_n} \circ \mathcal{N}_n \Vert \operatorname{Tr}_{R_n} \circ \mathcal{M}_n) (300)
$$

$$
\geq \sum_{i=1}^{n} D_{\mathbf{S},\alpha}^{\inf,\infty}(\mathrm{Tr}_{R_i} \circ \mathcal{N}_i || \mathrm{Tr}_{R_i} \circ \mathcal{M}_i)
$$
\n(301)

$$
=\sum_{i=1}^{n}D_{\mathrm{M},\alpha}^{\mathrm{inf},\infty}(\mathrm{Tr}_{R_{i}}\circ\mathcal{N}_{i}\|\mathrm{Tr}_{R_{i}}\circ\mathcal{M}_{i})
$$
\n(302)

$$
\geq \sum_{i=1}^{n} \frac{1}{m} D_{\mathbf{M},\alpha}^{\mathrm{inf}} \left( (\mathrm{Tr}_{R_i} \circ \mathcal{N}_i)^{\otimes m} \middle\| (\mathrm{Tr}_{R_i} \circ \mathcal{M}_i)^{\otimes m} \right) \tag{303}
$$

where  $m \ge 2$  and we used Eq. [\(96\)](#page-19-0) for the equality and the superadditivity of  $D_{\text{M},\alpha}$  in Lemma [24](#page-15-2) for the last inequality. Note that both of these results were applied for the family of states  $\mathscr{A}_m$  =  $(\text{Tr}_{R_i} \circ \mathcal{N}_i)^{\otimes m}(\mathscr{D})$  and  $\mathscr{B}_m = (\text{Tr}_{R_i} \circ \mathcal{M}_i)^{\otimes m}(\mathscr{D})$  which satisfies Assumption [25](#page-15-1) as proved in Lemma [42.](#page-40-1)

Observe that assumption [\(294\)](#page-42-1) implies assumption (\*) as  $\log \text{Tr}(\mathcal{M}_i^{\otimes m}(\sigma)) \leq m \log \lambda_{\max}(\mathcal{M}_i^\dagger(I)) \leq$  $\frac{C}{4}m$ . As a result, Lemma [31](#page-20-0) gives

$$
\frac{1}{m}D_{\mathbf{M},\alpha}^{\text{inf}}((\text{Tr}_{R_i}\circ\mathcal{N}_i)^{\otimes m}\|(\text{Tr}_{R_i}\circ\mathcal{M}_i)^{\otimes m})\tag{304}
$$

$$
\geq D^{\inf,\infty}(\text{Tr}_{R_i}\circ\mathcal{N}_i \| \text{Tr}_{R_i}\circ\mathcal{M}_i) - (1-\alpha)(2+C)^2m - \frac{2(d^2+d)\log(m+d)}{m},\tag{305}
$$

for  $1 - \frac{1}{(2+C)m} < \alpha < 1$ . Let us now choose  $1-\alpha = \frac{8d^2 \log m}{(2+C)^2 m}$  $\frac{8d^2\log m}{(2+C)^2m^2}$  and assume that  $m\geq \max\left(d,\left(\frac{16d^2}{2+C}\right)$  $\frac{16d^2}{2+C}\right)^2$ so that the condition  $\alpha \geq 1 - \frac{1}{(2+\epsilon)}$  $\frac{1}{(2+C)m}$  is satisfied and  $\log(m+d) \leq 2 \log m$ .

Putting everything together, we get

$$
D_{\mathrm{H},\varepsilon} \left( \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{R_0}) \middle\| \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{R_0}) \right)
$$
\n(306)

$$
\geq \left(\sum_{i=1}^{n} D^{\inf,\infty}(\operatorname{Tr}_{R_i} \circ \mathcal{N}_i || \operatorname{Tr}_{R_i} \circ \mathcal{M}_i)\right) - n \frac{16d^2 \log m}{m} - \frac{(2+C)^2 m^2}{8d^2 \log m} \log \frac{1}{\varepsilon} \tag{307}
$$

We now choose  $m = \left(\frac{64d^4n}{(2+\sqrt{32}}\right)^2$  $\sqrt{(2+C)^2 \log \frac{1}{\varepsilon}}$  $\int^{1/3}$ . With this choice

$$
D_{\mathrm{H},\varepsilon} \left( \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{R_0}) \middle\| \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{R_0}) \right)
$$
(308)

$$
\geq \left(\sum_{i=1}^{n} D^{\inf,\infty}(\text{Tr}_{R_i} \circ \mathcal{N}_i \| \text{Tr}_{R_i} \circ \mathcal{M}_i)\right) - C' n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon},\tag{309}
$$

for a constant  $C'$  that only depends on C and d. □

As a corollary, we get a weak form of the entropy accumulation statement for the smoothed maxentropy obtained in [\[MFSR22\]](#page-57-5). We recall that the smoothed max-entropy is defined as

$$
H_{\max}^{\varepsilon}(B|C)_{\rho} = \log \inf_{\substack{\tilde{\rho}_{BC} \in \mathcal{H}_{+}(BC) \\ \text{Tr}(\tilde{\rho}) \le 1 \\ F(\rho, \tilde{\rho}) \ge 1-\varepsilon}} \sup_{\sigma_C \in \mathcal{D}(C)} \left\| \tilde{\rho}_{BC}^{\frac{1}{2}} \text{id}_{B} \otimes \sigma_C^{\frac{1}{2}} \right\|_{1}^{2}.
$$
 (310)

**Corollary 44.** *(H<sub>max</sub>-entropy accumulation.)* Let  $\mathcal{N}_i \in \text{CPTP}(R'_{i-1} : B_i C_i R'_i)$  be quantum channels and  $\rho_{R'_0} \in \mathscr{D}(R_0)$  *be a quantum state. Define the state*  $\rho_{B_1...B_nC_1...C_nR'_n} = \prod_{i=1}^n \mathcal{N}_i(\rho_{R'_0})$ *. We have for*  $\varepsilon \in [0,1/2]$ ,

$$
H_{\max}^{\varepsilon}(B_1 \dots B_n | C_1 \dots C_n)_{\rho} \le \sum_{i=1}^n \sup_{\omega \in \mathscr{D}(R'_{i-1})} H(B_i | C_i)_{\mathcal{N}(\omega)} + K n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon},
$$
 (311)

where  $K$  is a constant only depending on  $\max_i \dim B_iC_i$ .

Remark 5 The second order term we achieve with our proof technique is worse than the one achieved in [\[DFR20\]](#page-55-5) and [\[MFSR22\]](#page-57-5). In addition, the statement of [\[MFSR22\]](#page-57-5) is stronger in that it also includes conditioning on the system  $R'_i$  provided a non-signalling assumption is satisfied. We still believe that our new proof technique, which is more naturally adapted to find upper bounds on  $H_{\text{max}}$  (as opposed to the techniques of [\[MFSR22\]](#page-57-5) which naturally apply to  $H_{\text{min}}$ ) could lead to insights and improvements for the applications of entropy accumulation. However, this is outside the scope of this paper and we leave it for future work.

*Proof.* We apply Theorem [43](#page-42-2) with the following replacements: letting  $C_i'$  be isomorphic to  $C_i$ , we set  $R_i \leftarrow R'_i C'_{i+1} \dots C'_n$  for  $i = 0$  to  $n, A_i \leftarrow B_i C_i$  for  $i = 1$  to  $n, \mathcal{N}_i \leftarrow \mathcal{N}_i \otimes \text{Tr}_{C'_i} \otimes \mathcal{I}_{C'_{i+1} \dots C'_n}$  (here, *I* refers to the identity map) and finally  $M_i$  is defined as  $M_i(X_{R'_{i-1}C'_i...C'_n}) = id_{B_i} \otimes X_{C_iC'_{i+1}...C'_n} \otimes$  $|0\rangle\langle 0|_{R'_i}$ , where  $|0\rangle\langle 0|_{R'_i}$  is an arbitrary fixed state in  $\mathscr{D}(R'_i)$ .

Theorem [43](#page-42-2) gives for the left hand side of [\(295\)](#page-42-3) and any  $\sigma \in \mathscr{D}(C_1 \dots C_n)$ :

$$
D_{\mathrm{H},\varepsilon}\left(\mathrm{Tr}_{R_n}\circ\prod_{i=1}^n\mathcal{N}_i(\rho_{R'_0}\otimes\sigma_{C'_1\ldots C'_n})\middle\|\mathrm{Tr}_{R_n}\circ\prod_{i=1}^n\mathcal{M}_i(|0\rangle\langle0|_{R'_0}\otimes\sigma_{C'_1\ldots C'_n})\right) \tag{312}
$$

$$
=D_{\mathrm{H},\varepsilon}(\rho_{B_{1}...B_{n}C_{1}...C_{n}}||\mathrm{id}_{B_{1}...B_{n}}\otimes\sigma_{C_{1}...C_{n}}).
$$
\n(313)

As the right hand side of [\(295\)](#page-42-3) does not depend on  $\sigma$ , we can take an infimum over  $\sigma$  and use Proposition [53](#page-67-0) to get the following

$$
\inf_{\sigma} D_{\mathrm{H},\varepsilon} \left( \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{N}_i (\rho_{R'_0} \otimes \sigma_{C'_1...C'_n}) \middle\| \mathrm{Tr}_{R_n} \circ \prod_{i=1}^n \mathcal{M}_i (|0\rangle\langle 0|_{R'_0} \otimes \sigma_{C'_1...C'_n}) \right)
$$
(314)

$$
\leq -H_{\max}^{\sqrt{2\varepsilon}}(B_1 \dots B_n | C_1 \dots C_n)_{\rho}.
$$
\n(315)

On the right hand side of [\(295\)](#page-42-3), we have terms of the form  $D^{\inf,\infty}(\text{Tr}_{R_i} \circ \mathcal{N}_i || \text{Tr}_{R_i} \circ \mathcal{M}_i)$ . Note that for  $\omega \in \mathscr{D}(R'_{i-1}C'_{i} \dots C'_{n})$ , we have  $(\text{Tr}_{R_i} \circ \mathcal{N}_i)(\omega) = \mathcal{N}_i(\omega_{R'_{i-1}})$  and  $(\text{Tr}_{R_i} \circ \mathcal{M}_i)(\omega) = id_{B_i} \otimes \omega_{C_i}$ . As a result,  $D^{\inf}(\text{Tr}_{R_i} \circ \mathcal{N}_i) | \text{Tr}_{R_i} \circ \mathcal{M}_i) = \inf_{\omega \in \mathscr{D}(R'_{i-1})} -H(B_i|C_i)_{\mathcal{N}_i(\omega)}$ , where we used the fact that  $-H(B|C)_{\rho} = \inf_{\sigma \in \mathscr{D}(C)} D(\rho_{BC} || \mathrm{id}_{B} \otimes \sigma_C)$ . We now need to evaluate the regularization:

$$
D^{\inf,\infty}(\mathrm{Tr}_{R_i}\circ\mathcal{N}_i||\mathrm{Tr}_{R_i}\circ\mathcal{M}_i) \tag{316}
$$

$$
= \inf_{m \ge 1} \frac{1}{m} \inf_{\substack{\omega \in \mathcal{D}((R'_{i-1})^{\otimes m}) \\ \sigma \in \mathcal{D}((C_i...C_n)^{\otimes m})}} D(\mathcal{N}_i^{\otimes m}(\omega) \|\mathrm{id}_{B_{i,1}...B_{1,m}} \otimes \sigma_{C_{i,1}...C_{i,m}}) \tag{317}
$$

$$
= -\sup_{m\geq 1} \sup_{\omega} \frac{1}{m} H(B_{i,1} \dots B_{i,m} | C_{i,1} \dots C_{i,m})_{\mathcal{N}_i^{\otimes m}(\omega)}
$$
(318)

$$
\geq -\sup_{m\geq 1} \sup_{\omega} \frac{1}{m} \sum_{j=1}^{m} H(B_{i,j}|C_{i,j})_{\mathcal{N}_i(\omega_{R'_{i,j}})}
$$
(319)

<span id="page-45-1"></span>
$$
= - \sup_{\omega \in \mathcal{D}(R'_{i-1})} H(B_i|C_i)_{\mathcal{N}_i(\omega)},
$$
\n(320)

which shows that in this case, the regularization is not needed. In order to apply Theorem [43,](#page-42-2) we need to check condition [\(294\)](#page-42-1). First, we have for any  $\omega \in \mathcal{D}(R'_i) \text{Tr}(\mathcal{M}_i(\omega)) = \dim B_i$ . In addition, as in the previous calculation, for  $\alpha \in [1/2, 1]$ , we have

$$
D_{\mathcal{P},\alpha}^{\text{inf}}((\text{Tr}_{R_i} \circ \mathcal{N}_i)^{\otimes m} \| (\text{Tr}_{R_i} \circ \mathcal{M}_i)^{\otimes m}) \tag{321}
$$

$$
= \inf_{\substack{\omega \in \mathscr{D}((R'_{i-1})^{\otimes m}) \\ \sigma \in \mathscr{D}((C_i...C_n)^{\otimes m})}} D_{P,\alpha}(\mathcal{N}_i^{\otimes m}(\omega_{R'_{i-1,1}...R'_{i-1,m}}) \|\mathrm{id}_{B_{i,1}...B_{1,m}} \otimes \sigma_{C_{i,1}...C_{i,m}})
$$
(322)

$$
= -\sup_{\omega} H_{P,\alpha}^{\uparrow}(B_{i,1}\dots B_{i,m}|C_{i,1}\dots C_{i,m})_{\mathcal{N}_i^{\otimes m}(\omega)},\tag{323}
$$

using the notation  $H_{P,\alpha}^{\uparrow}(D|E)_{\rho} = -\inf_{\sigma \in \mathscr{D}(E)} D_{P,\alpha}(\rho_{DE} || id_D \otimes \sigma_E)$  from [\[TBH14\]](#page-58-13). It is shown in [\[TBH14,](#page-58-13) Lemma 1], that an explicit choice of  $\sigma$  achieves this infimum namely  $\sigma_E^{(\alpha)} = \frac{(\text{Tr}_D \rho_{DE}^{\alpha})^{1/\alpha}}{\text{Tr}(\text{Tr}_D \rho_{DE}^{\alpha})^{1/2}}$  $\frac{(\Pi D \nu_{DE})}{(\text{Tr}(\text{Tr}_D \rho_{DE}^{\alpha}))^{1/\alpha}}.$ But in Lemma [52](#page-66-1) (Appendix [E\)](#page-64-1), we showed that for this choice  $D_{P,3/2}(\rho_{DE} \| \mathrm{id}_D \otimes \sigma_E^{(\alpha)}$  $E^{(\alpha)}(E) \leq 4 \log \dim D.$ Applying this to the state  $\rho = (\text{Tr}_{R_i} \circ \mathcal{N}_i)^{\otimes m}(\omega)$ , an optimal choice for  $\sigma$  in [\(322\)](#page-45-1) is given by  $\sigma_m^{(\alpha)} =$  $(\text{Tr}_{B_{i}^{\otimes m}} \rho^{\alpha})^{1/\alpha}$  $\frac{\prod_{i=1}^{n} \sum_{j=1}^{n} (T_{\text{max}} - \rho^{\alpha})^{1/\alpha}}{\text{Tr}(\text{Tr}_{B_i^{\otimes m}} \rho^{\alpha})^{1/\alpha}}$ . We get that for any  $\omega \in \mathscr{D}(R_{i-1}^{\otimes m})$ ,

$$
D_{\mathcal{P},3/2}((\text{Tr}_{R_i} \circ \mathcal{N}_i)^{\otimes m}(\omega) \|\text{id}_{B_i}^{\otimes m} \otimes \sigma_m^{(\alpha)}) \le 4m \log \dim B_i. \tag{324}
$$

This means that choosing  $C = 16 \max_i \log \dim B_i$  satisfies condition [\(294\)](#page-42-1). □

# <span id="page-45-0"></span>8 Application 4: efficient bounds for quantum resource theory

The generalized quantum Stein's lemma [\[BP10,](#page-55-4) [HY24,](#page-56-3) [Lam24\]](#page-57-3) has found broad applications in quantum resource theory. However, the optimal Stein exponents derived in these works are often challenging to compute, limiting their applicability. In contrast, our new generalized Stein's lemma, presented in Theorem [33,](#page-26-1) offers computational efficiency. To leverage this advantage, even when the task of interest does not directly satisfy Assumption [25,](#page-15-1) one can follow the general methodology of relaxing the set of interest to one that does, particularly regarding the polar assumption in (A.4). For example, in entanglement theory, the set of separable states can be relaxed to the Rains set, which satisfies all necessary assumptions. Similarly, in fault-tolerant quantum computing, the set of stabilizer states can be relaxed to the set of states with non-positive mana, which also fulfills the required conditions.

In the following, we provide several examples to illustrate this idea. Generally, this approach can be applied by verifying the conditions of the relevant theory and performing the necessary relaxations when required. As a result, we anticipate that this approach has the potential for far-reaching applications beyond the specific cases discussed here.

#### <span id="page-46-0"></span>8.1 Quantum entanglement distillation

One of the most significant applications of the generalized Stein's lemma is in entanglement distillation, an essential quantum information processing task that involves converting multiple copies of noisy entangled states into a smaller number of Bell states. The distillable entanglement of a bipartite state  $\rho_{AB}$ , denoted by  $E_{D,\Omega}(\rho_{AB})$ , represents the maximum number of Bell states that can be extracted from the given state with asymptotically vanishing error under the operation class  $\Omega$ . It has been established that the distillable entanglement under asymptotically non-entanglement generating operations, denoted by  $E_{D, \text{ANE}}$ , is given by the regularized relative entropy of entanglement [\[BP10\]](#page-55-4),

$$
E_{D,\text{ANE}}(\rho_{AB}) = D^{\infty}(\rho_{AB}||\text{SEP}) := \lim_{n \to \infty} \frac{1}{n} D(\rho_{AB}^{\otimes n}||\text{SEP}(A^n : B^n))
$$
(325)

where SEP( $A^n : B^n$ ) denotes the set of all separable states between  $\mathcal{H}_A^{\otimes n}$  and  $\mathcal{H}_B^{\otimes n}$ . Evaluating this quantity is hard in general, as it involves a limit as well as the separability problem, which is known to be computationally hard [\[Gur03\]](#page-56-15).

Here, we can utilize our generalized AEP to derive an efficient lower bound for  $D^{\infty}(\rho_{AB}||\text{SEP})$ . Note that a frequently-used relaxation of SEP is given by the Rains set [\[Rai01,](#page-58-14) [ADMVW02\]](#page-54-7)  $^6$  $^6$ 

$$
\text{Rains}(A:B) := \left\{ \sigma \in \mathcal{H}_+(AB) : \|\sigma^{\mathsf{T}_B}\|_1 \le 1 \right\}. \tag{326}
$$

We can show that it satisfies Assumption [25.](#page-15-1) This is because we have, by the SDP duality, that

$$
h_{\text{Rains}}(\omega) = \sup_{\sigma \in \text{Rains}(A:B)} \text{Tr}[\omega \sigma] = \inf_{\gamma \ge \omega} ||\gamma^{\mathsf{T}_B}||_{\infty},\tag{327}
$$

where  $\|\cdot\|_{\infty}$  is the spectral norm. By the multiplicativity of  $\|\cdot\|_{\infty}$  we can easily check that  $h_{\text{Rains}}$  is sub-multiplicative under tensor product, which is equivalent to the polar assumption (A.4) by Lemma [8.](#page-8-2) The rest of assumptions can also be easily verified.

Therefore, we have the relaxation

$$
D^{\infty}(\rho_{AB} \|\text{SEP}) \ge D^{\infty}(\rho_{AB} \|\text{Rains}),\tag{328}
$$

where the right-hand side known as the regularized Rains bound can be efficiently estimated as discussed in Remark [2](#page-16-8) by considering  $\mathscr{A}_n = \{\rho^{\otimes n}\}\$  and  $\mathscr{B}_n =$  Rains $(A_{[n]} : B_{[n]})$  which has efficient SDP representations [\[FWTD19\]](#page-56-2).

Similar to the regularized relative entropy of entanglement, the regularized Rains bound has the operational meaning that it is the distillable entanglement under Rains-preserving operations  $E_{D, \text{Rains}}$ , that is, [\[RFWG19\]](#page-58-15)

$$
E_{D,\text{Rains}}(\rho_{AB}) = D^{\infty}(\rho_{AB} || \text{Rains}). \tag{329}
$$

This marks the first time that an operational regularized entanglement measure has been shown to be efficiently computable, even when expressed as a regularized formula, beyond zero-error settings. Previous work in [\[LMR24\]](#page-57-12) studied the zero-error entanglement cost under PPT operations and proved that it is efficiently computable despite the absence of a closed-form formula.

As the Rains-preserving operations is a superset of the local operation and classical communication (LOCC) operations, the regularized Rains bound also gives an upper bound on the distillable entanglement under LOCC operations  $E_{D,LOCC}$ , that is,

$$
E_{D,\text{LOCC}}(\rho_{AB}) \le D^{\infty}(\rho_{AB} \|\text{Rains}) \le E_{D,\text{ANE}}(\rho_{AB})
$$
\n(330)

This improves the best known efficiently computable bound for  $E_{D,LOCC}$  as well.

<span id="page-46-1"></span> $6\text{ The set of PPT states does not satisfy Assumption 25, see discussion after Eq. (334).}$  $6\text{ The set of PPT states does not satisfy Assumption 25, see discussion after Eq. (334).}$  $6\text{ The set of PPT states does not satisfy Assumption 25, see discussion after Eq. (334).}$  $6\text{ The set of PPT states does not satisfy Assumption 25, see discussion after Eq. (334).}$  $6\text{ The set of PPT states does not satisfy Assumption 25, see discussion after Eq. (334).}$ 

#### <span id="page-47-0"></span>8.2 Magic state distillation

The above argument for entanglement distillation also applies to magic states, which is a key resource for fault-tolerant quantum computing [\[BK05,](#page-55-14) [VMGE14,](#page-58-16) [FL20\]](#page-56-16). The task of magic state distillation aims to extract as many copies of the target magic state as possible with asymptotically vanishing error. The distillable magic is denoted by  $M_{D,\Omega}$ , where  $\Omega$  represents the set of allowed operations. Typically, the most natural choice of operations involves stabilizer operations, and the corresponding distillable magic is denoted by  $M_{D,STAB}$ . However, characterizing this set of operations is challenging.

Motivated by the idea of the Rains bound from entanglement theory, Wang et al. [\[WWS20\]](#page-59-3) relaxed the set of all stabilizer states to a set of sub-normalized states with non-positive mana,  $W(\mathcal{H}) := \{ \sigma \in$  $\mathscr{H}_+(\mathcal{H}) : \|\sigma\|_{W,1} \leq 1$ , where  $\|\cdot\|_{W,1}$  denotes the Wigner trace norm. Based on the set W, Wang et al. introduced a magic measure  $D(\rho||\mathcal{W})$ , called thauma, and proved that it serves as an upper bound for the distillable magic under stabilizer operations, that is,

$$
M_{D,\text{STAB}}(\rho) \le D(\rho \| \mathcal{W})c(T),\tag{331}
$$

where  $c(T)$  is a constant that depends on the target magic state T.

Here, we can verify that W satisfies our Assumption  $25$ . Applying the generalized AEP, we can obtain an improved bound through regularization while keeping the computational efficiency.

To see this, it is straightforward to show that the support function of  $W$  is given by

$$
h_{\mathcal{W}}(\omega) = \sup_{\sigma \in \mathcal{W}} \text{Tr}[\omega \sigma] = \inf_{\gamma \ge \omega} ||\gamma||_{W,\infty},\tag{332}
$$

where the  $\|\cdot\|_{W,\infty}$  is the Wigner spectral norm and the second equality follows from the SDP duality. Since  $\|\cdot\|_{W,\infty}$  is multiplicative under tensor product, we can verify that the support function  $h_{\mathcal{W}}$  is submultiplicative under tensor product as well. Hence, the polar set  $\mathcal{W}^{\circ}$  is closed under tensor product by Lemma [8,](#page-8-2) and the remaining assumptions in Assumption [25](#page-15-1) can also be verified. Then, we can consider the regularization and get

$$
M_{D,\text{STAB}}(\rho) \le D^{\infty}(\rho \| \mathcal{W})c(T),\tag{333}
$$

where  $D^{\infty}(\rho||\mathcal{W})$  is the reguarlized thauma which remains efficiently computable by applying Remark [2](#page-16-8) with  $\mathscr{A}_n = \{\rho^{\otimes n}\}\$  and  $\mathscr{B}_n = \mathcal{W}(\mathcal{H}^{\otimes n})$ . This improves the best-known estimation of magic state distillation under stablizer opeations.

#### <span id="page-47-1"></span>8.3 Entanglement cost for quantum states and channels

The study of entanglement cost is dual to the task of entanglement distillation. Here, the goal is to prepare a given quantum state using the fewest possible Bell states. The *entanglement cost* of a quantum state, denoted as  $E_{C,\Omega}$ , is the minimum number of Bell states required to prepare one copy of this state under a class of operations Ω. Of particular interest is the set of LOCC operations. It is known that computing  $E_{\text{CLOCC}}$  is NP-hard in general [\[Hua14,](#page-56-17) Theorem 1]. Therefore, finding efficiently computable lower and upper bounds to estimate  $E_{C,LOC}$  is of fundamental importance. Here, we focus on deriving lower bounds, which indicate that no matter what preparation strategies are used, the amount of entanglement consumed cannot be smaller than this value.

There are several lower bounds in the literature, but they are unsatisfactory for different reasons. One such lower bound is given by the regularized PPT-relative entropy of entanglement [\[Hay17,](#page-56-18) Eq. (8.235)],

<span id="page-47-2"></span>
$$
E_{C,\text{LOCC}}(\rho) \ge D^{\infty}(\rho \|\text{PPT}) := \lim_{n \to \infty} \frac{1}{n} D(\rho^{\otimes n} \|\text{PPT}(A^n : B^n)),\tag{334}
$$

which is difficult to evaluate due to its regularization. Note that the PPT set does not satisfy the as-sumption (A.4) and therefore cannot directly apply Remark [2.](#page-16-8) A counter-example can be given by the projector on the  $\mathbb{C}^3 \otimes \mathbb{C}^3$  antisymmetric subspace, denoted as  $\rho^a$ . Since the support function  $h_{\mathscr{C}_{\text{PPT}}}(\cdot)$  is given by a semidefinite program, we find numerically that  $h_{PPT}((\rho^a)^{\otimes 2}) - h_{PPT}(\rho^a)^2 \approx 0.0093 > 0$ .

A single-letter lower bound for entanglement cost is provided by the quantum squashed entangle-ment [\[CW04\]](#page-55-15), but its computability remains uncertain due to the unbounded dimension of the extension system. Wang and Duan have proposed two single-letter SDP lower bounds in separate works:

$$
E_{\text{WD,1}}(\rho_{AB}) := -\log \max \text{Tr} \, \Pi_{\rho} V_{AB} \quad \text{s.t. } \|V_{AB}^{\mathsf{T}_B}\|_1 = 1, \ V_{AB} \ge 0, \quad [\text{WD17b}] \tag{335}
$$

$$
E_{\text{WD},2}(\rho_{AB}) := -\log \min \ \|Y_{AB}^{\mathsf{T}_B}\|_{\infty} \quad \text{s.t.} \ -Y_{AB} \le \Pi_{\rho}^{\mathsf{T}_B} \le Y_{AB}. \quad \text{[WD17a]} \tag{336}
$$

It has been shown that for any  $\rho \in \mathcal{D}(AB)$ ,

$$
\max\left\{E_{\text{WD,1}}(\rho), E_{\text{WD,2}}(\rho)\right\} \le D^{\infty}(\rho \|\text{PPT}) \le E_{C,\text{LOCC}}(\rho). \tag{337}
$$

By the definition of min-relative entropy and the Rains set, it is clear that  $E_{WD,1}$  can be written as  $^7$  $^7$ 

<span id="page-48-1"></span>
$$
E_{\text{WD,1}}(\rho_{AB}) = D_{\min}(\rho_{AB} || \text{Rains}(A:B)). \tag{338}
$$

After a detailed examination of the dual program for  $E_{WD,2}$ , we can also reformulate it in a comparable structure by introducing an appropriate set of operators.

**Lemma 45.** *For any*  $\rho \in \mathcal{D}(AB)$ *, it holds* 

$$
E_{\text{WD},2}(\rho_{AB}) = D_{\min}(\rho_{AB}||\text{WD}(A:B)),\tag{339}
$$

*with the set of Hermitian operators defined by*

$$
WD(A:B) := \left\{ \sigma \in \mathcal{H}(AB) : \exists Y \in \mathcal{H}(AB), \text{ s.t. } -Y \leq \sigma^{T_B} \leq Y, ||Y^{T_B}||_1 \leq 1 \right\}. \tag{340}
$$

*Proof.* Using the Lagrangian method, we have the dual SDP of  $E_{WD,2}$  as

$$
E_{\text{WD,2}}(\rho) = -\log \max \text{Tr} \Pi_{\rho} (V - F)^{\mathsf{T}_{B}} \quad \text{s.t.}
$$
  

$$
V + F = (W - X)^{\mathsf{T}_{B}}, \text{Tr}(W + X) \le 1, V, F, W, X \ge 0.
$$
 (341)

Let  $(V - F)^{T_B} = \sigma$ ,  $V + F = Y$ . By the definition of  $D_{\text{min}}$  and  $\mathcal{C}_{\text{WD}}$ , we have the desired result.  $\Box$ 

As discussed above, both SDP bounds  $E_{WD,1}$  and  $E_{WD,2}$  are essentially entanglement measures induced by the min-relative entropy. However, a significant limitation of these bounds is that they vanish for any full-rank state.

<span id="page-48-2"></span>**Proposition 46.** *For any full rank state*  $\rho \in \mathcal{D}(AB)$ , *it holds* 

$$
E_{\text{WD},1}(\rho) = E_{\text{WD},2}(\rho) = 0. \tag{342}
$$

*Proof.* If  $\rho_{AB}$  is full rank, then  $\Pi_o = I_{AB}$  and thus

$$
E_{\text{WD,1}}(\rho_{AB}) = -\log \max \left\{ \text{Tr} \,\sigma_{AB} : \,\sigma_{AB} \ge 0, \|\sigma_{AB}^{\mathsf{T}_B}\|_1 \le 1 \right\}. \tag{343}
$$

It is clear that for any feasible solution  $\sigma_{AB}$  it holds that  $\text{Tr } \sigma_{AB} = \text{Tr } \sigma_{AB}^{\mathsf{T}_B} \le ||\sigma_{AB}^{\mathsf{T}_B}||_1 \le 1$ . On the other hand, there is a feasible solution  $\sigma_{AB} = I_{AB}/|AB|$  such that  $\text{Tr} \sigma_{AB} = 1$ . Thus the maximization is taken at Tr  $\sigma_{AB} = 1$  and thus  $E_{WD,1}(\rho_{AB}) = 0$ . Similarly, given full rank  $\rho_{AB}$ , it holds

$$
E_{\text{WD},2}(\rho_{AB}) = -\log \max \left\{ \text{Tr}\,\sigma_{AB} : \sigma, Y \in \mathcal{H}(AB), -Y_{AB} \leq \sigma_{AB}^{\mathsf{T}_B} \leq Y_{AB}, \|Y^{\mathsf{T}_B}\|_1 \leq 1 \right\}. \tag{344}
$$

Then for any feasible solutions  $\sigma_{AB}$ ,  $Y_{AB}$ , it holds  $\text{Tr} \sigma_{AB} = \text{Tr} \sigma_{AB}^{\text{T}_B} \leq \text{Tr} Y_{AB} = \text{Tr} Y_{AB}^{\text{T}_B} \leq$  $||Y^{\mathsf{T}_B}||_1 \leq 1$ . On the other hand, considering the feasible solution  $\sigma_{AB} = Y_{AB} = I_{AB}/|AB|$ , we have Tr  $\sigma_{AB} = 1$ . Thus the maximization is taken at Tr  $\sigma_{AB} = 1$  and  $E_{WD,2}(\rho_{AB}) = 0$ .

<span id="page-48-0"></span><sup>&</sup>lt;sup>7</sup> The original definition of  $E_{WD,1}$  imposes the condition  $||V_{AB}^{T_B}||_1 = 1$ . However, it is equivalent to optimize over the condition  $||V_{AB}^{T_B}||_1 \leq 1$ , as the optimal solution can always be chosen at the boundary.

Recently, Wang et al. introduced the PPT<sub>k</sub> set [\[WJZ23\]](#page-59-6) as

<span id="page-49-0"></span>
$$
PPTk(A:B) := \left\{ \sigma \in \mathcal{H}_+(AB) : \text{Tr } f^k(\sigma) \le 1 \right\},\tag{345}
$$

where  $f(\sigma) := |\sigma^{T_B}|$  and  $k \in \mathbb{N}_+$ . Building on this set, the authors introduced an efficiently computable lower bound for entanglement cost [\[WJZ23\]](#page-59-6),

$$
E_{C,\text{LOCC}}(\rho_{AB}) \ge E_{\text{WIZ}}(\rho_{AB}) := D_{\text{S},1/2}(\rho_{AB} \|\text{PPT}_k(A:B)).\tag{346}
$$

In the following, we show that  $PPT_k$  satisfies Assumption [25](#page-15-1) and thus we can apply our Remark [2](#page-16-8) to get an improved bound.

<span id="page-49-2"></span>**Lemma 47.** *Let*  $k \geq 2$ *. The* PPT<sub>k</sub> *set defined in Eq.* [\(345\)](#page-49-0) *satisfies Assumption [25.](#page-15-1)* 

*Proof.* By the SDP duality, we have the support function

$$
h_{\text{PPT}_k}(\omega) = \inf \left\{ \|\gamma_k^{\mathsf{T}_B}\|_{\infty} : \gamma_1 \ge \omega, \ -\gamma_{i+1} \le \gamma_i^{\mathsf{T}_B} \le \gamma_{i+1}, \forall i \in \{1, 2, \cdots, k-1\} \right\}.
$$
 (347)

Then for any  $\omega_1, \omega_2$ , assume their optimal solutions in the dual program are respectively given by  $X_i$ and  $Y_i$  for all  $i \in \{1, 2, \dots, k\}$ . Then we have  $X_1 \otimes Y_1 \ge \omega_1 \otimes \omega_2$ . Since  $-X_{i+1} \le X_i^{T_B} \le X_{i+1}$  and  $-Y_{i+1} \le Y_i^{\mathsf{T}_B} \le Y_{i+1}$ , we have  $-I \le X_{i+1}^{-1/2} X_i^{\mathsf{T}_B} X_{i+1}^{-1/2} \le I$  and  $-I \le Y_{i+1}^{-1/2} Y_i^{\mathsf{T}_B} Y_{i+1}^{-1/2} \le I$  where the inverses are taken on the supports. This gives  $-I \leq X_{i+1}^{-1/2} X_i^{\mathsf{T}_B} X_{i+1}^{-1/2} \otimes Y_{i+1}^{-1/2} Y_i^{\mathsf{T}_B} Y_{i+1}^{-1/2} \leq I$ which is equivalent to  $-X_{i+1} \otimes Y_{i+1} \leq X_i^{\mathsf{T}_B} \otimes Y_i^{\mathsf{T}_B} \leq -X_{i+1} \otimes Y_{i+1}$ . So  $\{X_i \otimes Y_i\}_i$  is a feasible solution to the dual program of  $h_{\text{PPT}_k}(\omega_1 \otimes \omega_2)$ . By the multiplicativity of the infinity norm under tensor product, we know that  $h_{\text{PPT}_k}$  is sub-multiplicative under tensor product. This proves that PPT<sub>k</sub> satisfies assumption (A.4) by Lemma [8.](#page-8-2) The rest of assumptions can also be easily checked.  $□$ 

**Theorem 48.** Let  $\rho \in \mathcal{D}(AB)$  and  $k \geq 2$ . Let  $(*) = \max\{E_{WD,1}(\rho), E_{WD,2}(\rho), E_{WJZ}(\rho)\}\)$  represent *the previously known bounds. Then it holds*

$$
(*) \leq D_{\mathcal{M}}(\rho \|\text{PPT}_k) \leq D^{\infty}(\rho \|\text{PPT}_k) \leq D^{\infty}(\rho \|\text{PPT}) \leq E_{C,\text{LOCC}}(\rho), \tag{348}
$$

*Moreover, both*  $D_M(\rho \| \text{PPT}_k)$  *and*  $D^\infty(\rho \| \text{PPT}_k)$  *can be efficiently computed.* 

*Proof.* It has been shown that [\[WJZ23,](#page-59-6) Corollary 4],

<span id="page-49-1"></span>
$$
PPT(A:B) \subseteq PPT_k(A:B) \subseteq \cdots \subseteq Rains(A:B). \tag{349}
$$

It is also clear from their definitions that  $PPT_2(A : B) \subseteq WD(A : B)$ . This implies

$$
PPT(A:B) \subseteq PPT_k(A:B) \subseteq \cdots PPT_2(A:B) \subseteq WD(A:B) \cap Rains(A:B). \tag{350}
$$

Therefore, the first two inequalities of the asserted result follow from the relation of divergences in Lemma [16](#page-10-3) and the relation of sets in Eq. [\(350\)](#page-49-1). The second inequality follows from the superadditivity in Lemma [24](#page-15-2) and the asymptotic equivalence in Lemma [29.](#page-18-1) The third inequality follows from the relation in Eq. [\(350\)](#page-49-1). The last equality is known from Eq. [\(337\)](#page-48-1). The computability of  $D_M(\rho||PPT_k)$  and  $D^{\infty}(\rho||PPT_k)$  follows from Lemma 47 and Remark 2.  $D^{\infty}(\rho||PPT_k)$  follows from Lemma [47](#page-49-2) and Remark [2.](#page-16-8)

**Remark 6** Following the same argument in [\[WJZ23\]](#page-59-6), we can show that the new measures  $D_M(\rho||PPT_k)$ and  $D^{\infty}(\rho||PPT_k)$  also satisfy the properties such as normalization, faithfulness and (super-)additivity.

Besides the bounds mentioned, which are established on PPT, there is another efficiently computable lower bound on  $E_{\text{CLOCC}}$  given by Lami and Regula in [\[LR23\]](#page-57-13),

$$
E_{\text{LR}}(\rho) := \log \sup \left\{ \text{Tr} \, X \rho : \| X^{\mathsf{T}_B} \|_{\infty} \le 1, \| X \|_{\infty} = \text{Tr} [X \rho] \right\}. \tag{351}
$$

In the following, we compare our new bounds with previously established ones through several examples, including Isotropic states and Werner states. To the best of our knowledge, the entanglement costs for these states remain unresolved. Additionally, we use randomly generated quantum states to showcase the broad applicability and improvement of our bound across unstructured quantum states. In all cases, our experiments clearly demonstrate the superiority of our bound over existing ones.

Example 1 (Isotropic states and Werner states.) The Isotropic state is defined by a convex mixture of the maximally entangled state and its orthogonal complement,

$$
\rho_{I,p} := p|\Phi\rangle\langle\Phi| + \frac{1-p}{d^2 - 1}(I - |\Phi\rangle\langle\Phi|),\tag{352}
$$

where  $|\Phi\rangle = \frac{1}{\sqrt{2}}$  $\frac{d}{dt} \sum_{i=1}^{d} |ii\rangle$  is the *d*-dimensional maximally entangled state. The PPT-relative entropy of entanglement for  $\rho_{I,p}$  and its regularization are given by [\[Rai98,](#page-58-17) Theorem 7]

$$
D^{\infty}(\rho_{I,p}||\text{PPT}) = D(\rho_{I,p}||\text{PPT}) = \begin{cases} 0 & \text{if } 0 \le p \le \frac{1}{d}, \\ \log d + p \log p + (1-p) \log \frac{1-p}{d-1} & \text{if } \frac{1}{d} \le p \le 1. \end{cases}
$$
(353)

The Werner state is defined by a convex mixture of the normalized projectors on the symmetric  $(\rho^s)$  and anti-symmetric  $(\rho^a)$  subspaces,

$$
\rho_{W,p} := (1-p)\rho^s + p\rho^a = \frac{1-p}{d(d+1)}(I+S) + \frac{p}{d(d-1)}(I-S),\tag{354}
$$

where  $S = \sum_{i,j=1}^{d} |ij\rangle\langle ji|$  is the SWAP operator of dimension d. The regularized PPT-relative entropy of entanglement for  $\rho_{W,p}$  is given by [\[AEJ](#page-54-8)<sup>+</sup>01]

$$
D^{\infty}(\rho_{W,p}||\text{PPT}) = \begin{cases} 0, & \text{if } 0 \le p \le \frac{1}{2}, \\ 1 - h(p), & \text{if } \frac{1}{2} < p \le \frac{d+2}{2d}, \\ \log \frac{d+2}{d} + (1-p) \log \frac{d-2}{d+2}, & \text{if } p > \frac{d+2}{2d}. \end{cases}
$$
(355)

Note that both the Isotropic states and the Werner states are full-rank states for any  $p \in (0, 1)$ . As a result, the previous bounds  $E_{WD,1}$  and  $E_{WD,2}$  reduce to zero, as shown in Proposition [46.](#page-48-2) We then compare our new bound  $D_M(\rho||PPT_2)$  with the previously established bounds  $E_{WJZ}$  [\[WJZ23\]](#page-59-6),  $E_{LR}$  [\[LR23\]](#page-57-13) and the analytical bound  $D^{\infty}(\rho||PPT)$  in Figure [7.](#page-51-0) It turns out that  $D_M(\rho||PPT_2)$  coincides with the analytical bound  $D^{\infty}(\rho||PPT)$  for both cases, significantly improving the numerical bounds  $E_{\text{WJZ}}$  and  $E_{\text{LR}}$ .

<span id="page-51-0"></span>

 $E_{LR}$  [\[LR23\]](#page-57-13) and  $D^{\infty}(\rho||$ PPT) [Rai98, AEJ<sup>+</sup>01] for (a) Isotropic states and (b) Werner states. The Figure 7: Comparison of the new lower bound  $D_M(\rho||PPT_2)$  with previous bounds  $E_{WJZ}$  [\[WJZ23\]](#page-59-6), horizontal axis is the state parameter  $p$  and the vertical axis is the value of the entanglement measure.

general, we root sold comparison nere with  $E_K$  (ER25) by generating random orpartite states according<br>to the Hilbert-Schmidt measure, with varying ranks. For each rank, we generate 500 quantum states of  $\frac{1}{2}$ **EXAMPLE 2** (Kandom quantum states.) Since  $D_M(p||1|12)$  has been proved to be better than  $D_{WJZ}$  in general, we focus our comparison here with  $E_{LR}$  [\[LR23\]](#page-57-13) by generating random bipartite states according **Example 2** (Random quantum states.) Since  $D_M(\rho||PPT_2)$  has been proved to be better than  $E_{WJZ}$  in dimension 3 ⊗ 3. The comparison is presented in Figure [8.](#page-51-1) It is evident that  $D_M(\rho||PT_2)$  outperforms  $E_{LR}$  in most cases, particularly for higher-rank states.

<span id="page-51-1"></span>

Figure 8: Comparison of the new bound  $D_M(\rho||PPT_2)$  with the previously known bound  $E_{LR}$  [\[LR23\]](#page-57-13) for randomly generated quantum states with different ranks.

Similar to the entanglement cost for quantum states, the entanglement cost of a quantum channel, denoted by  $E_{\text{CLOCC}}(\mathcal{N})$ , represents the minimal rate at which entanglement (between the sender and receiver) is required to simulate multiple copies of the channel, given the availability of free classical communication. It is known that [\[BBCW13\]](#page-54-9)

$$
E_{C,\text{LOCC}}(\mathcal{N}) \ge \sup_{\rho \in \mathcal{D}(AA')} E_{C,\text{LOCC}}(\mathcal{N}_{A \to B}(\rho_{AA'})),\tag{356}
$$

where system  $\mathcal{H}_{A'}$  is isomorphic to system  $\mathcal{H}_A$ . Wang et al. introduced a lower bound for the entanglement cost of a quantum channel in [\[WJZ23\]](#page-59-6),

$$
E_{C,\text{LOCC}}(\mathcal{N}) \ge E_{\text{WJZ}}(\mathcal{N}) \ge E_{\text{WJZ}}(\mathcal{N}_{A \to B}(\Phi_{AA'}))
$$
\n(357)

where  $E_{\text{WJZ}}(\mathcal{N}) := \sup_{\rho \in \mathscr{D}(AA')} E_{\text{WJZ}}(\mathcal{N}_{A \to B}(\rho_{AA'}))$  and  $\Phi_{AA'}$  is the maximally entangled state. This lower bound has been used to demonstrate that the resource theory of entanglement is irreversible for amplitude damping channels.

Specifically, the amplitude damping channel is defined by

$$
\mathcal{N}_{\text{ad}}(\rho) = E_0 \rho E_0^\dagger + E_1 \rho E_1^\dagger,\tag{358}
$$

with Kraus operators  $E_0 = |0\rangle\langle 0| + \sqrt{1-\gamma}|1\rangle\langle 1|$  and  $E_1 = \sqrt{\gamma}|0\rangle\langle 1|$ . Its quantum capacity, i.e., the maximal rate at which entanglement can be generated from the channel, is known as [\[GF05\]](#page-56-19)

$$
Q(\mathcal{N}_{\text{ad}}) = \max_{p \in [0,1]} h_2((1 - \gamma)p) - h_2(\gamma p), \tag{359}
$$

where  $h_2$  is the binary entropy. It has been shown in [\[WJZ23\]](#page-59-6) that for  $0.25 \lesssim \gamma < 1$ ,

$$
E_{C,\text{LOCC}}(\mathcal{N}_{\text{ad}}) \ge E_{\text{WJZ}}(\mathcal{N}_{\text{ad}}(\Phi_{AA'})) > Q(\mathcal{N}_{\text{ad}}),\tag{360}
$$

Here, we can improve this bound by

$$
E_{C,\text{LOCC}}(\mathcal{N}_{\text{ad}}) \ge D_{\text{M}}(\mathcal{N}_{\text{ad}}(\Phi_{AA'}) \|\text{PPT}_2) > Q(\mathcal{N}_{\text{ad}})
$$
(361)

and show that the strict gap exists across the entire parameter region  $0 < \gamma < 1$  in Figure [9.](#page-53-1)

<span id="page-53-1"></span>

Figure 9: Comparison of the new lower bound  $D_M(\mathcal{N}_{ad}(\Phi_{AA'})||PPT_2)$  with the previously known bounds  $E_{\text{WJZ}}$  [\[WJZ23\]](#page-59-6),  $E_{\text{LR}}$  [\[LR23\]](#page-57-13) and the quantum capacity Q for amplitude damping channel. The horizontal axis is the channel parameter  $\gamma$  and the vertical axis is the value of the entanglement measure.

# <span id="page-53-0"></span>9 Conclusion

entry for measured relative entropy between two sets of states, the enant rule property for quantum<br>relative entropy, and minimum output channel divergence—are of independent interest and are likely to The significance of this work is multifaceted. First, we established a very general quantum AEP with explicit convergence guarantees and computational efficiency. Given that the AEP is central to information theory and is evident in its broad applications in both classical and quantum information theory, we expect that our generalized quantum AEP plays a similar role and advances the study of quantum information science further. Second, our generalized quantum Stein's lemma addresses open questions raised by Brandão et al. [[BHLP20,](#page-54-2) Section 3.5] and Mosonyi et al. [\[MSW22,](#page-57-4) Section VI], which seek a Stein's lemma with computational efficiency. We provide a general methodology for approximating the problem of interest when the original problem is difficult to solve and our result is not directly applicable, along with illustrations in quantum resource theories. Due to the generality of this methodology, we anticipate that this approach has the potential for far-reaching applications beyond the specific cases discussed here. Third, we introduce a fully quantum version of adversarial hypothesis testing and a relative entropy accumulation theorem, which might find applications in quantum cryptography. Finally, we believe that the technical tools established in this work—including the variational formula, superadditivity for measured relative entropy between two sets of states, the chain rule property for quantum have further applications.

Many problems remain open for future investigation. For example, we have demonstrated that the regularized divergence can be efficiently computed using convex optimization techniques. However, developing a more explicit algorithm and implementing numerical solutions for these optimization programs is an area for further exploration. Additionally, designing a general algorithm to construct the smallest superset of a given set that satisfies the polar assumption presents an intriguing challenge. This would extend the applicability of our generalized AEP to boarder areas.

While we have obtained explicit bounds on the second-order term for the generalized AEP of order  $O(n^{2/3} \log n)$ , such bounds are not tight in general. Can one bound the second-order term by  $O(\sqrt{n})$  as in the AEP? Sharpening the second-order term would provide a more comprehensive understanding of the convergence behavior and enhance its accuracy in practical applications. This work also opens new directions by considering adversarial quantum channel discrimination and establishing its connection with the minimum output channel divergence. We can keep exploring the strong converse exponent and error exponent as well as quantum channel discrimination in the Chernoff setting. We anticipate that the quantities of interest will also be characterized by the corresponding minimum output channel divergence. We also proposed a new framework for quantum resource theory in which state transformations are performed without requiring precise characterization of the states being manipulated, making it more robust to imperfections. We proved the asymptotic reversibility of this theory under asymptotic resource non-generating operations, while leaving the systematic study of other different operational classes and settings for future exploration.

Acknowledgements. We would like to thank Kerry He for the help of using QICS package. K.F. is supported by the University Development Fund (Grant No. UDF01003565) from The Chinese University of Hong Kong, Shenzhen. O.F. acknowledges support by the European Research Council (ERC Grant AlgoQIP, Agreement No. 851716), by the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 101017733 (VERIqTAS) and by the Agence Nationale de la Recherche under the Plan France 2030 with the reference ANR-22-PETQ-0009.

# References

<span id="page-54-9"></span><span id="page-54-8"></span><span id="page-54-7"></span><span id="page-54-6"></span><span id="page-54-5"></span><span id="page-54-4"></span><span id="page-54-3"></span><span id="page-54-2"></span><span id="page-54-1"></span><span id="page-54-0"></span>

<span id="page-55-18"></span><span id="page-55-17"></span><span id="page-55-16"></span><span id="page-55-15"></span><span id="page-55-14"></span><span id="page-55-13"></span><span id="page-55-12"></span><span id="page-55-11"></span><span id="page-55-10"></span><span id="page-55-9"></span><span id="page-55-8"></span><span id="page-55-7"></span><span id="page-55-6"></span><span id="page-55-5"></span><span id="page-55-4"></span><span id="page-55-3"></span><span id="page-55-2"></span><span id="page-55-1"></span><span id="page-55-0"></span>

- <span id="page-56-13"></span>[FF21a] K. Fang and H. Fawzi. Geometric Renyi divergence and its applications in quantum ´ channel capacities. *Communications in Mathematical Physics*, 384(3):1615–1677, 2021.
- <span id="page-56-6"></span>[FF21b] H. Fawzi and O. Fawzi. Defining quantum divergences via convex optimization. *Quantum*, 5:387, 2021.
- <span id="page-56-4"></span>[FFRS20] K. Fang, O. Fawzi, R. Renner, and D. Sutter. Chain rule for the quantum relative entropy. *Physical Review Letters*, 124(10):100501, 2020.
- <span id="page-56-10"></span>[FGW21] K. Fang, G. Gour, and X. Wang. Ultimate limits of quantum channel discrimination. *arXiv preprint arXiv:2110.14842*, 2021.
- <span id="page-56-16"></span>[FL20] K. Fang and Z.-W. Liu. No-go theorems for quantum resource purification. *Physical Review Letters*, 125(6):060405, 2020.
- <span id="page-56-8"></span>[FS23] H. Fawzi and J. Saunderson. Optimal self-concordant barriers for quantum relative entropies. *SIAM Journal on Optimization*, 33(4):2858–2884, 2023.
- <span id="page-56-9"></span>[FST22] O. Fawzi, A. Shayeghi, and H. Ta. A hierarchy of efficient bounds on quantum capacities exploiting symmetry. *IEEE Transactions on Information Theory*, 68(11):7346–7360, November 2022.
- <span id="page-56-2"></span>[FWTD19] K. Fang, X. Wang, M. Tomamichel, and R. Duan. Non-asymptotic entanglement distillation. *IEEE Transactions on Information Theory*, 65(10):6454–6465, 2019.
- <span id="page-56-19"></span>[GF05] V. Giovannetti and R. Fazio. Information-capacity description of spin-chain correlations. *Physical Review A—Atomic, Molecular, and Optical Physics*, 71(3):032314, 2005.
- <span id="page-56-12"></span>[Gou24] G. Gour. Resources of the quantum world. *arXiv preprint arXiv:2402.05474*, 2024.
- <span id="page-56-15"></span>[Gur03] L. Gurvits. Classical deterministic complexity of edmonds' problem and quantum entanglement. In *Proceedings of the thirty-fifth annual ACM symposium on Theory of computing*, pages 10–19, 2003.
- <span id="page-56-18"></span>[Hay17] M. Hayashi. Quantum information theory. *Graduate Texts in Physics, Springer*, 2017.
- <span id="page-56-11"></span>[HMO08] F. Hiai, M. Mosonyi, and T. Ogawa. Error exponents in hypothesis testing for correlated states on a spin chain. *Journal of Mathematical Physics*, 49(3), 2008.
- <span id="page-56-1"></span>[HP91] F. Hiai and D. Petz. The proper formula for relative entropy and its asymptotics in quantum probability. *Communications in Mathematical Physics*, 143:99–114, 1991.
- <span id="page-56-7"></span>[HR11] T. Holenstein and R. Renner. On the randomness of independent experiments. *IEEE Transactions on Information Theory*, 57(4):1865–1871, 2011.
- <span id="page-56-14"></span>[HSF24] K. He, J. Saunderson, and H. Fawzi. Qics: Quantum information conic solver. *arXiv preprint arXiv:2410.17803*, 2024.
- <span id="page-56-5"></span>[HT16] M. Hayashi and M. Tomamichel. Correlation detection and an operational interpretation of the Rényi mutual information. *Journal of Mathematical Physics*, 57(10), 2016.
- <span id="page-56-17"></span>[Hua14] Y. Huang. Computing quantum discord is NP-complete. *New Journal of Physics*, 16(3):33027, 2014.
- <span id="page-56-0"></span>[HV93] T. S. Han and S. Verdu. Approximation theory of output statistics. ´ *IEEE Transactions on Information Theory*, 39(3):752–772, 1993.
- <span id="page-56-3"></span>[HY24] M. Hayashi and H. Yamasaki. Generalized quantum Stein's lemma and second law of quantum resource theories. *arXiv preprint arXiv:2408.02722*, 2024.
- <span id="page-57-16"></span>[IRS16] R. Iten, J. M. Renes, and D. Sutter. Pretty good measures in quantum information theory. *IEEE Transactions on Information Theory*, 63(2):1270–1279, 2016.
- <span id="page-57-7"></span>[KZ05] A. Kurdila and M. Zabarankin. *Convex Functional Analysis*. Systems & Control: Foundations & Applications. Birkhäuser Basel, 2005.
- <span id="page-57-3"></span>[Lam24] L. Lami. A solution of the generalised quantum Stein's lemma. *arXiv preprint arXiv: 2408.06410*, 2024.
- <span id="page-57-14"></span>[LKDW18] F. Leditzky, E. Kaur, N. Datta, and M. M. Wilde. Approaches for approximate additivity of the holevo information of quantum channels. *Physical Review A*, 97(1):012332, 2018.
- <span id="page-57-11"></span>[LLS+23] F. Leditzky, D. Leung, V. Siddhu, G. Smith, and J. A. Smolin. The platypus of the quantum channel zoo. *IEEE Transactions on Information Theory*, 69(6):3825–3849, June 2023.
- <span id="page-57-2"></span>[LM02] E. Levitan and N. Merhav. A competitive Neyman-Pearson approach to universal hypothesis testing with applications. *IEEE Transactions on Information Theory*, 48(8):2215– 2229, 2002.
- <span id="page-57-12"></span>[LMR24] L. Lami, F. A. Mele, and B. Regula. Computable entanglement cost. *arXiv preprint arXiv:2405.09613*, 2024.
- <span id="page-57-13"></span>[LR23] L. Lami and B. Regula. No second law of entanglement manipulation after all. *Nature Physics*, 19(2):184–189, 2023.
- <span id="page-57-0"></span>[McM53] B. McMillan. The basic theorems of information theory. *The Annals of mathematical statistics*, pages 196–219, 1953.
- <span id="page-57-5"></span>[MFSR22] T. Metger, O. Fawzi, D. Sutter, and R. Renner. Generalised entropy accumulation. In *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 844–850. IEEE, 2022.
- <span id="page-57-9"></span>[MH11] M. Mosonyi and F. Hiai. On the quantum rényi relative entropies and related capacity formulas. *IEEE Transactions on Information Theory*, 57(4):2474–2487, April 2011.
- <span id="page-57-8"></span>[MH23] M. Mosonyi and F. Hiai. Some continuity properties of quantum Rényi divergences. *arXiv preprint arXiv: 2209.00646*, 2023.
- <span id="page-57-6"></span> $[MLDS+13]$  M. Müller-Lennert, F. Dupuis, O. Szehr, S. Fehr, and M. Tomamichel. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*, 54(12):122203, 2013.
- <span id="page-57-10"></span>[MO15] M. Mosonyi and T. Ogawa. Two approaches to obtain the strong converse exponent of quantum hypothesis testing for general sequences of quantum states. *IEEE Transactions on Information Theory*, 61(12):6975–6994, 2015.
- <span id="page-57-4"></span>[MSW22] M. Mosonyi, Z. Szilagyi, and M. Weiner. On the error exponents of binary state discrimination with composite hypotheses. *IEEE Transactions on Information Theory*, 68(2):1032–1067, February 2022.
- <span id="page-57-1"></span>[NO00] H. Nagaoka and T. Ogawa. Strong converse and Stein's lemma in quantum hypothesis testing. *IEEE Transactions on Information Theory*, 46(7):2428–2433, feb 2000.
- <span id="page-57-15"></span>[NS09] M. Nussbaum and A. Szkoła. The Chernoff Lower Bound For Symmetric Quantum Hypothesis Testing. *The Annals of Statistics*, 37(2):1040–1057, 2009.

<span id="page-58-18"></span><span id="page-58-17"></span><span id="page-58-16"></span><span id="page-58-15"></span><span id="page-58-14"></span><span id="page-58-13"></span><span id="page-58-12"></span><span id="page-58-11"></span><span id="page-58-10"></span><span id="page-58-9"></span><span id="page-58-8"></span><span id="page-58-7"></span><span id="page-58-6"></span><span id="page-58-5"></span><span id="page-58-4"></span><span id="page-58-3"></span><span id="page-58-2"></span><span id="page-58-1"></span><span id="page-58-0"></span>[Pet86] D. Petz. Quasi-entropies for finite quantum systems. *Reports on Mathematical Physics*, 23(1):57–65, 1986. [Pia09] M. Piani. Relative entropy of entanglement and restricted measurements. *Physical Review Letters*, 103(16):160504, 2009. [OWW18] H. Qi, Q. Wang, and M. M. Wilde. Applications of position-based coding to classical communication over quantum channels. *Journal of Physics A: Mathematical and Theoretical*, 51(44):444002, 2018. [Rai98] E. Rains. An improved bound on distillable entanglement. *arXiv preprint quantph/9809082*, 1998. [Rai01] E. M. Rains. A semidefinite program for distillable entanglement. *IEEE Transactions on Information Theory*, 47(7):2921–2933, 2001. [Ren05] R. Renner. *Security of QKD*. PhD thesis, PhD thesis, ETH, 2005, quant-ph/0512258, 2005. [RFWG19] B. Regula, K. Fang, X. Wang, and M. Gu. One-shot entanglement distillation beyond local operations and classical communication. *New Journal of Physics*, 21(10):103017, 2019. [RSB24] T. Rippchen, S. Sreekumar, and M. Berta. Locally-measured Rényi divergences.  $arXiv$ *preprint arXiv:2405.05037*, 2024. [Sch95] B. Schumacher. Quantum coding. *Physical Review A*, 51(4):2738, 1995. [Sha48] C. E. Shannon. A Mathematical Theory of Communication. *Bell System Technical Journal*, 27(3):379–423, jul 1948. [Sio58] M. Sion. On general minimax theorems. *Pacific Journal of Mathematics*, 8(1):171–176, mar 1958. [TBH14] M. Tomamichel, M. Berta, and M. Hayashi. Relating different quantum generalizations of the conditional rényi entropy. *Journal of Mathematical Physics*, 55(8), 2014. [TCR09] M. Tomamichel, R. Colbeck, and R. Renner. A fully quantum asymptotic equipartition property. *IEEE Transactions on Information Theory*, 55(12):5840–5847, 2009. [Tom16] M. Tomamichel. *Quantum Information Processing with Finite Resources*, volume 5 of *SpringerBriefs in Mathematical Physics*. Springer International Publishing, Cham, 2016. [Ume54] H. Umegaki. Conditional expectation in an operator algebra. *Tohoku Mathematical Journal, Second Series*, 6(2-3):177–181, 1954. [VH97] S. Verdu and T. S. Han. The role of the asymptotic equipartition property in noiseless source coding. *IEEE Transactions on Information Theory*, 43(3):847–857, 1997. [VMGE14] V. Veitch, S. H. Mousavian, D. Gottesman, and J. Emerson. The resource theory of stabilizer quantum computation. *New Journal of Physics*, 16(1):013009, 2014. [VV95] S. Vembu and S. Verdú. Generating random bits from an arbitrary source: Fundamental limits. *IEEE Transactions on Information Theory*, 41(5):1322–1332, 1995. [Wat09] J. Watrous. Semidefinite programs for completely bounded norms. *Theory of Computing*, 5(1):217–238, 2009.

- <span id="page-59-1"></span>[Wat18] J. Watrous. *The theory of quantum information*. Cambridge University Press, 2018.
- <span id="page-59-2"></span>[WBHK20] M. M. Wilde, M. Berta, C. Hirche, and E. Kaur. Amortized channel divergence for asymptotic quantum channel discrimination. *Letters in Mathematical Physics*, 110:2277– 2336, 2020.
- <span id="page-59-5"></span>[WD17a] X. Wang and R. Duan. Irreversibility of Asymptotic Entanglement Manipulation under Quantum Operations Completely Preserving Positivity of Partial Transpose. *Physical Review Letters*, 119(18):1–7, 2017.
- <span id="page-59-4"></span>[WD17b] X. Wang and R. Duan. Nonadditivity of Rains' bound for distillable entanglement. *Physical Review A*, 95(6):1–5, 2017.
- <span id="page-59-6"></span>[WJZ23] X. Wang, M. Jing, and C. Zhu. Computable and faithful lower bound for entanglement cost. *arXiv preprint arXiv:2311.10649*, 2023.
- <span id="page-59-3"></span>[WWS20] X. Wang, M. M. Wilde, and Y. Su. Efficiently computable bounds for magic state distillation. *Physical Review Letters*, 124(9):090505, 2020.
- <span id="page-59-0"></span>[WWY14] M. M. Wilde, A. Winter, and D. Yang. Strong converse for the classical capacity of entanglement-breaking and hadamard channels via a sandwiched Rényi relative entropy. *Communications in Mathematical Physics*, 331(2):593–622, 2014.

# Appendices

In Section [A](#page-60-0) we present the variational formula for the measured relative entropy with different parameters  $\alpha$ . In Section [B](#page-62-0) we provide an alternative proof for Lemma [32.](#page-25-1) In Section [C,](#page-63-0) we present a counter-example to a potential improvement of the chain rule. In Section [D,](#page-64-0) we will see that proving the superadditivity of  $D_M(\mathscr{A}|\mathscr{B})$  directly from its variational formula requires the partial trace assumption. Some useful properties are presented in Section [E.](#page-64-1)

# <span id="page-60-0"></span>A Variational formula for  $D_{\text{M},\alpha}(\mathscr{A}||\mathscr{B})$

Let  $\bar{h}_{\mathscr{C}}(X) := \inf_{Y \in \mathscr{C}} \text{Tr}[XY]$  and  $\mathscr{C}^{\star} := \{ X \in \mathscr{H}_{+} : \text{Tr}[XY] \geq 1, \forall Y \in \mathscr{C} \}.$ **Lemma 49.** Let  $\mathcal{A} \subseteq \mathcal{D}$  and  $\mathcal{B} \subseteq \mathcal{H}$  be two compact convex sets. Then it holds

$$
D_{\mathbf{M},\alpha}(\mathscr{A}||\mathscr{B}) = \frac{\alpha}{\alpha - 1} \log(*) \tag{362}
$$

*with*

$$
(*) = \begin{cases} \inf_{W,V} h_{\mathscr{A}}(W) & \text{s.t. } W \in \mathscr{H}_{++}, \ V \in \mathscr{B}^{\circ}, \ W^{\frac{\alpha}{\alpha-1}} \le V, & \text{if } \alpha \in (0,1/2), \\ \inf_{W,V} h_{\mathscr{A}}(V) & \text{s.t. } W \in \mathscr{H}_{++}, \ W \in \mathscr{B}^{\circ}, \ W^{\frac{\alpha-1}{\alpha}} \le V, & \text{if } \alpha \in [1/2,1), \\ \sup_{W,V} \bar{h}_{\mathscr{A}}(V) & \text{s.t. } W \in \mathscr{H}_{++}, \ W \in \mathscr{B}^{\star}, \ W^{\frac{\alpha-1}{\alpha}} \ge V, & \text{if } \alpha \in (1,+\infty), \end{cases}
$$
(363)

*where the right-hand side is a convex program.*

*Proof.* The case for  $\alpha \in [1/2, 1)$  has been proved in the main text. Now we prove the other two cases. Let  $\alpha \in (0, 1/2)$ . For any fixed  $\rho \in \mathscr{A}$  and  $\sigma \in \mathscr{B}$ , we have by Eq. [\(29\)](#page-10-1) that

$$
Q_{\mathcal{M},\alpha}(\rho||\sigma) := \inf_{W \in \mathcal{H}_{++}} \alpha \operatorname{Tr}[\rho W] + (1 - \alpha) \operatorname{Tr}[\sigma W^{\frac{\alpha}{\alpha - 1}}]
$$
(364)

$$
= \inf_{\substack{W \in \mathcal{H}_+ \\ W^{\alpha-1} \le V}} \alpha \operatorname{Tr} \left[ \rho W \right] + (1 - \alpha) \operatorname{Tr} [\sigma V], \tag{365}
$$

where the second equality follows by introducing an additional variable  $V$ . Then we have

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \inf_{\substack{W \in \mathscr{H}_{++} \\ \sigma \simeq 1 \le V}} \alpha \operatorname{Tr}[\rho W] + (1 - \alpha) \operatorname{Tr}[\sigma V]. \tag{366}
$$

Note that all  $\mathscr{A}, \mathscr{B}$  and  $\{(W, V) : W \in \mathscr{H}_{++}, W^{\frac{\alpha}{\alpha-1}} \leq V\}$  are convex sets, with  $\mathscr{A}, \mathscr{B}$  being compact. Moreover, the objective function is linear in  $(\rho, \sigma)$ , and also linear in  $(W, V)$ . So we can apply Sion's minimax theorem [\[Sio58,](#page-58-11) Corollary 3.3] to exchange the infimum and supremum and get

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{\alpha-1} \le V}} \sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \alpha \operatorname{Tr} [\rho W] + (1 - \alpha) \operatorname{Tr} [\sigma V] \tag{367}
$$

$$
= \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{\alpha-1} \leq V}} \alpha \sup_{\rho \in \mathcal{A}} \text{Tr}[\rho W] + (1 - \alpha) \sup_{\sigma \in \mathcal{B}} \text{Tr}[\sigma V]. \tag{368}
$$

By using the definition of the support function, we have

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{\alpha-1} \le V}} \alpha h_{\mathscr{A}}(W) + (1 - \alpha) h_{\mathscr{B}}(V) \tag{369}
$$

$$
= \inf_{\substack{W \in \mathcal{H}_{++} \\ W^{\alpha-1} \le V}} h_{\mathscr{A}}(W)^{\alpha} h_{\mathscr{B}}(V)^{1-\alpha},
$$
\n(370)

where the second line follows from the weighed arithmetic-geometric mean inequality  $\alpha x + (1 \alpha$ )y  $\leq x^{\alpha}y^{1-\alpha}$  (with equality if and only if  $x = y$ ) and the fact that  $(W, V)$  is a feasible solution implies  $(kW, k^{\frac{\alpha}{\alpha-1}}V)$  is also a feasible solution for any  $k \ge 0$ . Therefore, we can choose  $k =$  $(h_{\mathscr{A}}(W)/h_{\mathscr{B}}(W))^{\alpha-1}$ , which implies  $h_{\mathscr{A}}(kW) = h_{\mathscr{B}}(k^{\frac{\alpha}{\alpha-1}}V)$  and therefore the equality of the weighed arithmetic-geometric mean is achieved. Similarly, for any feasible solution  $(W, V)$  we can always construct a new solution  $(Wh_{\mathscr{B}}(V)^{\frac{1-\alpha}{\alpha}}, V/h_{\mathscr{B}}(V))$  achieves the same objective value. This implies

$$
\sup_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{\frac{\alpha}{\alpha-1}} \leq V \\ h_{\mathscr{B}}(V)=1}} (h_{\mathscr{A}}(W))^{\alpha} = \inf_{\substack{W \in \mathscr{H}_{++} \\ W^{\frac{\alpha}{\alpha-1}} \leq V \\ h_{\mathscr{B}}(V) \leq 1}} (h_{\mathscr{A}}(W))^{\alpha},
$$
\n(371)

where the second equality follows by the same reasoning. Finally, noting that  $h_{\mathscr{B}}(V) \leq 1$  if and only if  $V \in \mathscr{B}^{\circ}$ , we have the asserted result in Eq. [\(53\)](#page-13-0). It is easy to check that the objective function  $h_{\mathscr{A}}(W)$ is convex in W and the feasible set is also a convex set.

Let  $\alpha \in (1, +\infty)$ . For any fixed  $\rho \in \mathcal{A}$  and  $\sigma \in \mathcal{B}$ , we have by Eq. [\(29\)](#page-10-1) that

$$
Q_{\mathcal{M},\alpha}(\rho\|\sigma) := \sup_{W\in\mathscr{H}_{++}} \alpha \operatorname{Tr}\left[\rho W^{\frac{\alpha-1}{\alpha}}\right] + (1-\alpha) \operatorname{Tr}[\sigma W] \tag{372}
$$

$$
= \sup_{\substack{W \in \mathcal{H}_{+} \\ W^{\frac{\alpha-1}{\alpha}} \ge V \ge 0}} \alpha \operatorname{Tr} [\rho V] + (1 - \alpha) \operatorname{Tr} [\sigma W], \tag{373}
$$

where the second equality follows by introducing an additional variable  $V$ . Then we have

$$
\inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \sup_{\substack{W \in \mathscr{H}_{++} \\ W \frac{\alpha-1}{\alpha} \ge V \ge 0}} \alpha \operatorname{Tr} [\rho V] + (1 - \alpha) \operatorname{Tr} [\sigma W]. \tag{374}
$$

Note that all  $\mathscr{A}, \mathscr{B}$  and  $\{(W, V) : W \in \mathscr{H}_{++}, W^{\frac{\alpha-1}{\alpha}} \geq V \geq 0\}$  are convex sets, with  $\mathscr{A}, \mathscr{B}$  being compact. Moreover, the objective function is linear in  $(\rho, \sigma)$ , and also linear in  $(W, V)$ . So we can apply Sion's minimax theorem [\[Sio58,](#page-58-11) Corollary 3.3] to exchange the infimum and supremum and get

$$
\inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \sup_{\substack{W \in \mathscr{H}_{++} \\ W^{\frac{\alpha-1}{\alpha}} \ge V \ge 0}} \inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} \alpha \operatorname{Tr} [\rho V] + (1 - \alpha) \operatorname{Tr} [\sigma W] \tag{375}
$$

$$
= \sup_{\substack{W \in \mathcal{H}_{++} \\ W^{\frac{\alpha-1}{\alpha}} \ge V \ge 0}} \alpha \inf_{\rho \in \mathcal{A}} \text{Tr}[\rho V] + (1 - \alpha) \inf_{\sigma \in \mathcal{B}} \text{Tr}[\sigma W]. \tag{376}
$$

Let  $\bar{h}_{\mathscr{C}}(X) := \inf_{\rho \in \mathscr{C}} \text{Tr}[X\rho]$ , we have

$$
\inf_{\substack{\rho \in \mathscr{A} \\ \sigma \in \mathscr{B}}} Q^{\mathcal{M}}_{\alpha}(\rho \| \sigma) = \sup_{\substack{W \in \mathscr{H}_{++} \\ W^{\frac{\alpha-1}{\alpha}} \ge V \ge 0}} \alpha \bar{h}_{\mathscr{A}}(V) + (1 - \alpha) \bar{h}_{\mathscr{B}}(W) \tag{377}
$$

$$
= \sup_{\substack{W \in \mathcal{H}_{++} \\ W^{\frac{\alpha-1}{\alpha}} \ge V \ge 0}} \bar{h}_{\mathscr{A}}(V)^{\alpha} \bar{h}_{\mathscr{B}}(W)^{1-\alpha}, \tag{378}
$$

where the second line follows from the weighed arithmetic-geometric mean inequality  $\alpha x + (1 \alpha$ )y  $\leq x^{\alpha}y^{1-\alpha}$  (with equality if and only if  $x = y$ ) and the fact that  $(W, V)$  is a feasible solution implies  $(kW, k^{\frac{\alpha-1}{\alpha}}V)$  is also a feasible solution for any  $k \ge 0$ . Therefore, we can choose  $k =$  $(\bar{h}_{\mathscr{A}}(V)/\bar{h}_{\mathscr{B}}(W))^{\alpha}$ , which implies  $h_{\mathscr{A}}(kW) = h_{\mathscr{B}}(k^{\frac{\alpha-1}{\alpha}}V)$  and therefore the equality of the weighed arithmetic-geometric mean is achieved. Similarly, for any feasible solution  $(W, V)$  we can always construct a new solution  $(W/\bar{h}_{\mathscr{B}}(W), V\bar{h}_{\mathscr{B}}(W)^{\frac{1-\alpha}{\alpha}})$  achieves the same objective value. This implies

$$
\inf_{\rho \in \mathscr{A}} Q_{\alpha}^{\mathcal{M}}(\rho \| \sigma) = \sup_{\substack{W \in \mathscr{H}_{++} \\ \sigma \in \mathscr{B}}} (\bar{h}_{\mathscr{A}}(V))^{\alpha} = \sup_{\substack{W \in \mathscr{H}_{++} \\ W_{\alpha}^{\alpha-1} \ge V \ge 0}} (\bar{h}_{\mathscr{A}}(V))^{\alpha},
$$
\n
$$
\lim_{\substack{\sigma \in \mathscr{B} \\ \bar{h}_{\mathscr{B}}(W) = 1}} (\bar{h}_{\mathscr{A}}(V))^{\alpha},
$$
\n(379)

where the second equality follows by the same reasoning. Finally, noting that  $\bar{h}_{\mathscr{B}}(W) \geq 1$  if and only if  $W \in \mathscr{B}^{\star} := \{ X \in \mathscr{H}_{+} : \text{Tr}[XY] \geq 1, \forall Y \in \mathscr{B} \}$ , we have the asserted result in Eq. [\(53\)](#page-13-0). It is easy to check that the objective function  $\bar{h}_{\mathscr{A}}(V)$  is concave in V and the feasible set is also a convex set.  $\Box$ 

## <span id="page-62-0"></span>B Alternative proof for Lemma [32](#page-25-1)

Here we give an alternative proof for Lemma [32,](#page-25-1) which follows the argument outlined in [\[BP10,](#page-55-4) Section III]. Denote

$$
f(\mathscr{A}, \sigma, \varepsilon) := \sup_{M} \{ \text{Tr}[\sigma M] : 0 \le M \le I, \text{Tr}[\rho M] \le \varepsilon, \forall \rho \in \mathscr{A} \}.
$$
 (380)

By using the definition of dual cone of a convex set, we get

$$
f(\mathscr{A}, \sigma, \varepsilon) = \sup_{M} \{ \text{Tr}[\sigma M] : 0 \le M \le I, \varepsilon I - M \in \mathscr{A}^* \},
$$
\n(381)

where  $\mathscr{A}^* := \{X : \text{Tr}[X\rho] \geq 0, \forall \rho \in \mathscr{A}\}$  is the dual cone of  $\mathscr{A}$ . Then introducing the Lagrange multipliers  $X > 0$ ,  $Y > 0$  and  $Z \in \text{Cone}(\mathscr{A})$ , with  $\text{Cone}(\mathscr{A})$  being the cone generated by  $\mathscr{A}$ , we get the Lagrangian of  $f(\mathscr{A}, \sigma, \varepsilon)$  as

$$
L(\sigma, \varepsilon, M, X, Y, Z) := \text{Tr}[\sigma M] + \text{Tr}[XM] + \text{Tr}[Y(I - M)] + \text{Tr}[\varepsilon I - M]Z \tag{382}
$$

<span id="page-62-3"></span><span id="page-62-1"></span>
$$
= \operatorname{Tr}[M(\sigma + X - Y - Z)] + \operatorname{Tr}[Y] + \varepsilon \operatorname{Tr}[Z]. \tag{383}
$$

It is easy to see that  $M = (\varepsilon/2)I$  is a strictly feasible solution to the optimization problem in Eq. [\(381\)](#page-62-1). Therefore, by Slater's condition [\[BV04\]](#page-55-16),  $f(\mathscr{A}, \sigma, \varepsilon)$  is equal to its dual program, which is given by

$$
f(\mathscr{A}, \sigma, \varepsilon) = \inf_{Y, Z} \{ \text{Tr}[Y] + \varepsilon \text{Tr}[Z] : \sigma \le Y + Z, Y \ge 0, Z \in \text{Cone}(\mathscr{A}) \}.
$$
 (384)

Using that  $Tr[V_{+}] = min \{Tr[W] : W \ge 0, W \ge V\}$  with  $V_{+}$  being the positive part of V, we have

$$
f(\mathscr{A}, \sigma, \varepsilon) = \inf_{Z} \{ \text{Tr}[(\sigma - Z)_{+}] + \varepsilon \text{Tr}[Z] : Z \in \text{Cone}(\mathscr{A}) \}
$$
(385)

<span id="page-62-2"></span>
$$
= \inf_{\rho \in \mathscr{A}} \inf_{x>0} \{ \text{Tr}[(\sigma - x\rho)_+] + \varepsilon x \}
$$
\n(386)

where the second line follows by introducing  $Z = x\rho$  with  $\rho \in \mathscr{A}$  and the fact that  $\mathscr{A} \subseteq \mathscr{D}$  is a set of normalized quantum states. Applying the above result to the case that  $\mathscr{A} = \{\rho\}$ , we get

$$
f(\rho, \sigma, \varepsilon) = \sup_{M} \left\{ \text{Tr}[\sigma M] : 0 \le M \le I, \, \text{Tr}[\rho M] \le \varepsilon \right\} = \inf_{x > 0} \left\{ \text{Tr}[(\sigma - x\rho)_+] + \varepsilon x \right\}. \tag{387}
$$

Taking this to Eq.  $(386)$ , we have

$$
f(\mathscr{A}, \sigma, \varepsilon) = \inf_{\rho \in \mathscr{A}} \sup_M \{ \text{Tr}[\sigma M] : 0 \le M \le I, \text{Tr}[\rho M] \le \varepsilon \}. \tag{388}
$$

Replacing M with  $I - M$  in Eqs. [\(380\)](#page-62-3) and [\(388\)](#page-62-4), we get

$$
\inf_{M} \{ \text{Tr}[\sigma M] : 0 \le M \le I, \text{Tr}[\rho (I - M)] \le \varepsilon, \forall \rho \in \mathscr{A} \}
$$
\n(389)

$$
= \sup_{\rho \in \mathscr{A}} \inf_{M} \left\{ \text{Tr}[\sigma M] : 0 \le M \le I, \text{Tr}[\rho (I - M)] \le \varepsilon \right\}.
$$
 (390)

In a simpler notation, this shows

<span id="page-62-5"></span><span id="page-62-4"></span>
$$
\beta_{\varepsilon}(\mathscr{A}||\{\sigma\}) = \sup_{\rho \in \mathscr{A}} \beta_{\varepsilon}(\rho||\sigma). \tag{391}
$$

Now we are ready to prove the asserted result. Note that both B and  $\{M : 0 \le M \le I, \alpha(\mathcal{A}, M) \le \varepsilon\}$ are convex with the later being compact. Moreover, the objective function  $\text{Tr}[\sigma M]$  is linear in both  $\sigma$  and M. Therefore, we can apply Sion's minimax theorem [ $Sio58$ , Corollary 3.3] to exchange the infimum and supremum and get

$$
\beta_{\varepsilon}(\mathscr{A}||\mathscr{B}) = \inf_{\substack{0 \le M \le I \\ \alpha(\mathscr{A}, M) \le \varepsilon}} \sup_{\sigma \in \mathscr{B}} \text{Tr}[\sigma M] = \sup_{\sigma \in \mathscr{B}} \inf_{\substack{0 \le M \le I \\ \alpha(\mathscr{A}, M) \le \varepsilon}} \text{Tr}[\sigma M]. \tag{392}
$$

Combining with Eq.  $(391)$ , we have

$$
\beta_{\varepsilon}(\mathscr{A}||\mathscr{B}) = \sup_{\sigma \in \mathscr{B}} \beta_{\varepsilon}(\mathscr{A}||\{\sigma\}) = \sup_{\sigma \in \mathscr{B}} \sup_{\rho \in \mathscr{A}} \beta_{\varepsilon}(\rho||\sigma). \tag{393}
$$

This completes the proof.

## <span id="page-63-0"></span>C Counter-example to a potential improvement of the chain rule

The quantum channel divergence studied in most existing literatures [\[LKDW18\]](#page-57-14) use the same test states for both channels. So it may be expected that we can enhance the chain rule in Lemma [38](#page-36-2) by using the same test states as well. However, we show here that this is not possible by giving a counter-example. That is, the chain rule cannot be enhanced to

$$
D_{\scriptscriptstyle{\mathrm{M}}}(\mathcal{N}_{A\to B}(\rho_{RA})||\mathcal{M}_{A\to B}(\sigma_{RA})) \ge D_{\scriptscriptstyle{\mathrm{M}}}(\rho_{R}||\sigma_{R}) + D_{\scriptscriptstyle{\mathrm{M}}}^{\text{inf}}(\mathcal{N}||\mathcal{M})
$$
\n(394)

where the channel divergence takes the same input state

$$
D_{\scriptscriptstyle{\rm M}}^{\scriptscriptstyle{\rm inf}}(\mathcal{N}\|\mathcal{M}) := \inf_{\rho \in \mathscr{D}(A)} D_{\scriptscriptstyle{\rm M}}(\mathcal{N}_{A \to B}(\rho_A) \|\mathcal{M}_{A \to B}(\rho_A)). \tag{395}
$$

To see this, consider the generalized amplitude damping (GAD) channel, which is defined as

<span id="page-63-1"></span>
$$
\mathcal{A}_{\gamma,N}(\rho) = \sum_{i=1}^4 A_i \rho A_i^{\dagger}, \quad \gamma, N \in [0,1],
$$
\n(396)

with the Kraus operator

$$
A_1 = \sqrt{1 - N} (|0\rangle\langle 0| + \sqrt{1 - \gamma} |1\rangle\langle 1|), \qquad A_2 = \sqrt{\gamma(1 - N)} |0\rangle\langle 1|,
$$
 (397)

$$
A_3 = \sqrt{N}(\sqrt{1-\gamma}|0\rangle\langle 0| + |1\rangle\langle 1|), \qquad A_4 = \sqrt{\gamma N}|1\rangle\langle 0|.
$$
 (398)

Using convex optimization, we can numerically evaluate each terms  $D_M(\mathcal{N}_{A\to B}(\rho_{RA})||\mathcal{M}_{A\to B}(\sigma_{RA}))$ ,  $D_M(\rho_R || \sigma_R)$  and  $\tilde{D}_M^{\text{inf}}(\mathcal{N} || \mathcal{M})$ . Then in Figure [10\(](#page-64-2)a), we show that the channel divergence  $D_M^{\text{inf}}$  is subadditive under tensor product of channels. That is, it does not inherit the properties of the state divergence, making it not a suitable channel extension. Moreover, in Figure [10\(](#page-64-2)b), we show that the chain rule property in Eq. [\(394\)](#page-63-1) does not hold, as there are cases such that  $y < x_2$  in the plot.

(a) Subadditivity for channel divergence

(b) Random states with  $d_A = d_B = d_R = 2$ 

<span id="page-64-2"></span>

Figure 10: (a) Subadditivity for the channel divergence  $D^{M,\text{inf}}'$  where  $\mathcal{A}_{0.5,0}$  and  $\mathcal{A}_{p,0.9}$  are the GAD channels and  $p$  ranges from 0 to 1; (b) Random test for the chain rule property, where the quantum channels are chosen as  $A_{0.5,0.9}$  and  $A_{0.5,0.9}$ , the quantum states are 500 randomly generated quantum states with real entries,  $x_1 = D_M(\rho_R || \sigma_R) + D_M^{\text{inf}}(\mathcal{A}_{0.5,0} || \mathcal{A}_{0.5,0.9})$  and  $x_2 = D_M(\rho_R || \sigma_R) +$  $D_{\rm M}^{\rm inf'}({\cal A}_{0.5,0}\|{\cal A}_{0.5,0.9}).$ 

# <span id="page-64-0"></span>D Proving the superadditivity of  $D_M(\mathscr{A}|\mathscr{B})$  from its variational formula

Here we show that if we attempt to prove the superadditivity of Lemma [24](#page-15-2) by directly using the variational formula in Lemma [20,](#page-11-2)

$$
D_M(\mathscr{A}||\mathscr{B}) = \sup_{W \in \mathscr{B}_{++}^{\circ}} -h_{\mathscr{A}}(-\log W),\tag{399}
$$

we will require the stability assumption of  $\mathscr{A}_n$  under partial trace. Let  $W_1$  and  $W_2$  be any feasible solutions to  $D_M(\mathscr{A}_1|\mathscr{B}_1)$  and  $D_M(\mathscr{A}_2|\mathscr{B}_2)$ , respectively. It is clear that  $W_1 \otimes W_2$  is also a feasible solution to  $D_M(\mathscr{A}_{12}||\mathscr{B}_{12})$  by the polar assumption  $(\mathscr{B}_1)_+^{\circ} \otimes (\mathscr{B}_2)_+^{\circ} \subseteq (\mathscr{B}_{12})_+^{\circ}$  $\int_{+}^{\infty}$ . Now we aim to argue that this achieves an objective value no smaller than those by  $W_1$  and  $W_2$ . Note that  $-h_{\mathscr{A}}(- \log W) =$  $\inf_{\rho \in \mathcal{A}}$  Tr[ $\rho$  log W]. We hope to show

$$
\inf_{\rho_{12}\in\mathscr{A}_{12}} \text{Tr}[\rho_{12}\log(W_1\otimes W_2)] \ge \inf_{\rho_1\in\mathscr{A}_1} \text{Tr}[\rho_1\log W_1] + \inf_{\rho_2\in\mathscr{A}_2} \text{Tr}[\rho_2\log W_2].\tag{400}
$$

For any  $\rho_{12} \in \mathcal{A}_{12}$ , we have

$$
\text{Tr}[\rho_{12}\log(W_1 \otimes W_2)] = \text{Tr}[\rho_{12}(\log W_1) \otimes I] + \text{Tr}[\rho_{12}I \otimes (\log W_2)] \tag{401}
$$

<span id="page-64-3"></span>
$$
= \text{Tr}[\text{Tr}_2 \,\rho_{12}(\log W_1)] + \text{Tr}[\text{Tr}_1 \,\rho_{12}(\log W_2)]. \tag{402}
$$

If we had  $Tr_2 \rho_{12} \in \mathcal{A}_1$  and  $Tr_1 \rho_{12} \in \mathcal{A}_2$ , then we can conclude Eq. [\(400\)](#page-64-3). It is however not clear how to reach this conclusion without this assumption. This justifies our use of continuity to prove superadditivity in Lemma [24,](#page-15-2) thereby avoiding the partial trace assumption and placing fewer constraints on the subsequent theories.

## <span id="page-64-1"></span>E Useful properties

We establish an explicit continuity for  $D_{\alpha}$  was  $\alpha \to 1$  with  $\alpha < 1$ . It is analogous to [\[TCR09,](#page-58-2) Lemma 8] and [\[DFR20,](#page-55-5) Lemma B.8] and the proof is almost the same.

<span id="page-65-0"></span>**Lemma 50** (Explicit continuity of Rényi divergences as  $\alpha \to 1$  from below). Let  $\rho \in \mathcal{D}(\mathcal{H})$  and  $\sigma \in \mathscr{H}_{+}(\mathcal{H})$ *. Let*  $\eta = \left( \max(4, 2^{2D_{\text{P},3/2}(\rho\|\sigma)} + 2^{-2D_{\text{P},1/2}(\rho\|\sigma)} + 1) \right)^2$ *. For*  $\alpha \in (1 - 1/\log \eta, 1)$ *, we have*

$$
0 \le D(\rho \|\sigma) - D_{P,\alpha}(\rho \|\sigma) \le (1 - \alpha)(\log \eta)^2. \tag{403}
$$

*As a result, we also have*

$$
0 \le D(\rho \|\sigma) - D_{s,\alpha}(\rho\|\sigma) \le (1 - \alpha) \left( (\log \eta)^2 + D(\rho\|\sigma) + \log \text{Tr}(\sigma) \right). \tag{404}
$$

*Proof.* We show the result for classical distributions  $P$ , Q over  $X$  and then the result follows immediately for the Petz Rényi divergence by applying it to the corresponding Nussbaum-Szkoła distributions [[NS09\]](#page-57-15) as was done in [\[DF19\]](#page-55-17). We follow the proof from [\[TCR09,](#page-58-2) Lemma 8]. Letting  $\beta = \alpha - 1 < 0$ , we have

$$
D_{\alpha}(P||Q) = \frac{1}{\beta} \log \sum_{x \in \mathcal{X}} P(x) \left(\frac{P(x)}{Q(x)}\right)^{\beta}.
$$
 (405)

Letting  $t^{\beta} = 1 + \beta \ln t + r_{\beta}(t)$ , where  $r_{\beta}(t) = t^{\beta} - \beta \ln t - 1$ , we can write

$$
D_{\alpha}(P||Q) = \frac{1}{\beta} \log \left( 1 + \beta \sum_{x \in \mathcal{X}} P(x) \ln \frac{P(x)}{Q(x)} + \sum_{x \in \mathcal{X}} P(x) r_{\beta} \left( \frac{P(x)}{Q(x)} \right) \right).
$$
 (406)

We now use the fact that  $ln(1 + y) \leq y$ . Thus,

$$
D_{\alpha}(P||Q) \ge \frac{1}{\beta} \frac{1}{\ln 2} \left( \beta \sum_{x \in \mathcal{X}} P(x) \ln \frac{P(x)}{Q(x)} + \sum_{x \in \mathcal{X}} P(x) r_{\beta} \left( \frac{P(x)}{Q(x)} \right) \right)
$$
(407)

$$
= D(P||Q) + \frac{1}{\beta \ln 2} \sum_{x \in \mathcal{X}} P(x)r_{\beta} \left(\frac{P(x)}{Q(x)}\right). \tag{408}
$$

We now define  $s_\beta(t) = 2(\cosh(\beta \ln t) - 1)$  as in [\[TCR09\]](#page-58-2), but recall that we have  $\beta < 0$ . Note that  $s_\beta(t) = s_{\beta}(t)$  so we also have that it is monotonically increasing for  $t > 1$  and concave in t for  $\beta > -1/2$  and  $t \ge 3$ . In addition, we have  $r_\beta(t) = e^{\beta \ln t} + (-\beta \ln t + 1) - 2 \le e^{\beta \ln t} + e^{-\beta \ln t} - 2 = s_\beta(t)$ for all  $t \geq 0$ . Note that we also have  $s_{\beta}(t) = s_{\beta}(1/t)$  and  $s_{\beta}(t^2) = s_{2\beta}(t)$ . As a result,  $s_{\beta}(t) \leq$  $s_{2\beta}(1+\sqrt{t}+\sqrt{1/t})$ . As a consequence, because we assumed  $\alpha > 1 - 1/\log 16$ , we have  $2\beta \ge -1/2$ and  $1 + \sqrt{t} + \sqrt{1/t} \ge 3$  and we can use the concavity of  $s_{2\beta}$  and get

$$
D_{\alpha}(P||Q) \ge D(P||Q) + \frac{1}{\beta \ln 2} \sum_{x \in \mathcal{X}: P(x) > 0} P(x) s_{2\beta} \left( 1 + \sqrt{\frac{P(x)}{Q(x)}} + \sqrt{\frac{Q(x)}{P(x)}} \right)
$$
(409)

$$
\geq D(P||Q) + \frac{1}{\beta \ln 2} s_{2\beta} \left( \sum_{x \in \mathcal{X}: P(x) > 0} P(x) \left( 1 + \sqrt{\frac{P(x)}{Q(x)}} + \sqrt{\frac{Q(x)}{P(x)}} \right) \right) \tag{410}
$$

$$
\geq D(P||Q) + \frac{1}{\beta \ln 2} s_{2\beta} \left( 1 + 2^{2D_{3/2}(P||Q)} + 2^{-2D_{1/2}(P||Q)} \right) \tag{411}
$$

$$
\geq D(P\|Q) + \frac{1}{\beta \ln 2} s_{2\beta} \left(\sqrt{\eta}\right). \tag{412}
$$

Note that by Taylor's theorem around  $z = 0$ , we have  $e^z + e^{-z} - 1 \leq z^2 (e^z + e^{-z})$ . As a result  $\frac{1}{\beta \ln 2} s_{2\beta}(\sqrt{\eta}) \ge \frac{1}{\beta \ln 2} (2\beta \ln \sqrt{\eta})^2 \cosh(2\beta \ln \sqrt{\eta}) = \beta(\log \eta)^2 \ln 2 \cosh(\beta \ln \eta) \ge \beta(\log \eta)^2$  where we used the fact that  $\ln 2 \cosh(\ln 2) < 1$ , which proves that

$$
D_{\alpha}(P||Q) \ge D(P||Q) - (1 - \alpha)(\log \eta)^{2}.
$$
\n(413)

To prove the bound on the sandwiched Rényi divergence, we use the relation proved in [[IRS16\]](#page-57-16) between the different divergences in the regime  $\alpha \in (0,1)$ :

$$
D_{s,\alpha}(\rho||\sigma) \ge \alpha D_{P,\alpha}(\rho||\sigma) + (1-\alpha)(\log \text{Tr}(\rho) - \log \text{Tr}(\sigma)).\tag{414}
$$

This implies

$$
D_{\mathcal{S},\alpha}(\rho||\sigma) \ge D_{\mathcal{P},\alpha}(\rho||\sigma) - (1-\alpha)D_{\mathcal{P},\alpha}(\rho||\sigma) - (1-\alpha)\log \text{Tr}(\sigma) \tag{415}
$$

$$
\geq D(\rho \|\sigma) - (1 - \alpha) \left( (\log \eta)^2 + D(\rho \|\sigma) + \log \text{Tr}(\sigma) \right) \tag{416}
$$

⊓⊔

For  $\alpha > 1$ , we use [\[TCR09,](#page-58-2) Lemma 8], with a slightly different parametrization. The proof is the same as Lemma [50.](#page-65-0)

<span id="page-66-0"></span>**Lemma 51.** *Let*  $\rho \in \mathcal{D}(\mathcal{H})$  *and*  $\sigma \in \mathcal{H}_+(\mathcal{H})$ *. For any*  $\alpha \in (1, 1 + 1/\log n)$ *, we have* 

$$
D_{\mathcal{S},\alpha}(\rho\|\sigma) \le D(\rho\|\sigma) + (\alpha - 1)\left(\log \eta\right)^2,\tag{417}
$$

 $where \eta = \left(\max(4, 2^{D_{\text{P},3/2}(\rho||\sigma)} + 2^{-D_{\text{P},1/2}(\rho||\sigma)} + 1)\right)^2$ .

<span id="page-66-1"></span>**Lemma 52.** *Let*  $\alpha \in [1/2, 1]$ *,*  $\rho_{AB} \in \mathscr{D}(AB)$  *and*  $\sigma_B^{(\alpha)} = \frac{(\text{Tr}_A \rho_{AB}^{\alpha})^{\frac{1}{\alpha}}}{Z}$  $\frac{\rho_{AB}^{\alpha}\right)^{\alpha}}{Z}$ , where  $Z = \text{Tr}\left((\text{Tr}_A \rho_{AB}^{\alpha})^{\frac{1}{\alpha}}\right)$ . *Then*

$$
D_{\mathbf{P},\frac{3}{2}}(\rho_{AB}||\mathrm{id}_A \otimes \sigma_B^{(\alpha)}) \le 4\log d_A. \tag{418}
$$

Note that the choice of the parameter  $\frac{3}{2}$  is not arbitrary and this lemma does not hold for higher values. See the discussion in the proof of [\[DF19,](#page-55-17) Corollary III.5].

*Proof.* We have

$$
\operatorname{Tr}\left(\rho_{AB}^{3/2}(\operatorname{id}_A\otimes\sigma_B^{(\alpha)})^{-1/2}\right) = Z^{1/2}\operatorname{Tr}\left(\rho_{AB}^{3/2}\operatorname{id}_A\otimes(\operatorname{Tr}_A\rho_{AB}^{\alpha})^{-\frac{1}{2\alpha}}\right). \tag{419}
$$

We start by showing that  $Z \leq d_A$ . In fact, we use the operator Jensen inequality for the operator concave function  $x \mapsto x^{\alpha}$  as follows:

$$
\text{Tr}_A(\rho_{AB}^{\alpha}) = d_A \sum_a \frac{\langle a |}{\sqrt{d_A}} \rho_{AB}^{\alpha} \frac{|a\rangle}{\sqrt{d_A}}
$$
(420)

$$
\leq d_A \left( \sum_a \frac{\langle a |}{\sqrt{d_A}} \rho_{AB} \frac{|a\rangle}{\sqrt{d_A}} \right)^{\alpha} \tag{421}
$$

$$
=d_A^{1-\alpha}\rho_B^{\alpha}.\tag{422}
$$

As a result, we get

$$
Z \le \text{Tr}\left((d_A^{1-\alpha}\rho_B^{\alpha})^{\frac{1}{\alpha}}\right) = d_A^{\frac{1-\alpha}{\alpha}} \le d_A. \tag{423}
$$

For the inequality, we used the fact that the function  $x \mapsto x^{1/\alpha}$  is monotone and continuous and thus  $X \mapsto \text{Tr}(X^{\frac{1}{\alpha}})$  is monotone (see e.g., [\[Car10,](#page-55-18) Section 2.2]).

Now let us consider

$$
\operatorname{Tr}\left(\rho_{AB}^{3/2}\operatorname{id}_A\otimes\left(\operatorname{Tr}_A\rho_{AB}^\alpha\right)^{\frac{1}{\alpha}}\right)\leq\operatorname{Tr}\left(\rho_{AB}\operatorname{id}_A\otimes\left(\operatorname{Tr}_A\rho_{AB}\right)^{-\frac{1}{2\alpha}}\right)\tag{424}
$$

$$
\leq \text{Tr}(\rho_{AB}\rho_B^{-1})\tag{425}
$$

$$
\leq d_A,\tag{426}
$$

where we used for the first inequality the fact that  $\rho_{AB}^{3/2} \le \rho_{AB}$ ,  $\rho_{AB}^{\alpha} \ge \rho_{AB}$  and the operator antimonotonicity of the function  $x \mapsto x^{-\frac{1}{2\alpha}}$ , and for the second inequality the fact that  $\rho_B^{\frac{1}{2\alpha}} \ge \rho_B$ . As a result,

$$
D_{P,\frac{3}{2}}(\rho_{AB}||id_A \otimes \sigma_B^{(\alpha)}) \le 2\log(d_A^{1/2}d_A) \le 4\log d_A.
$$
 (427)

⊓⊔

<span id="page-67-0"></span>**Proposition 53.** For  $0 \leq \varepsilon < \frac{1}{2}$ , we have

$$
H_{\max}^{\sqrt{2\varepsilon}}(B|C)_{\rho} \leq -\inf_{\sigma_C \in \mathscr{D}(C)} D_{\mathrm{H},\varepsilon}(\rho_{BC} \|\mathrm{id}_B \otimes \sigma_C). \tag{428}
$$

*Proof.* Using Lemma [32](#page-25-1) with  $\mathscr{A} = \{ \rho_{BC} \}$  and  $\mathscr{B} = \{ id_B \otimes \sigma_C : \sigma_C \in \mathscr{D}(C) \}$ , we can write

$$
-\inf_{\sigma_C \in \mathcal{D}(C)} D_{\mathrm{H},\varepsilon}(\rho_{BC} \|\mathrm{id}_B \otimes \sigma_C) = \log \inf_{0 \le M \le I} \{ \sup_{\sigma_C} \{ \mathrm{Tr}(M I_B \otimes \sigma_C) : \mathrm{Tr}(M \rho_{BC}) \ge 1 - \varepsilon \} \tag{429}
$$

Let M be such that  $\text{Tr}(M\rho_{BC})$ , define  $\tilde{\rho}_{BC} = \sqrt{M}\rho_{BC}\sqrt{M}$ . Then, by the gentle measurement lemma, we have  $F(\rho, \tilde{\rho}) \ge 1 - \sqrt{2\varepsilon}$  (see e.g., [\[DKF](#page-55-12)<sup>+</sup>14, Lemma A.3]). Then, using Lemma [54,](#page-67-1) we get

$$
\sup_{\sigma_C \in \mathscr{D}(C)} \left\| (\sqrt{M} \rho_{BC} \sqrt{M})^{\frac{1}{2}} \mathrm{id}_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2 \leq \sup_{\sigma_C} \text{Tr}(MI \otimes \sigma_C) \tag{430}
$$

As such,

$$
-\inf_{\sigma_C \in \mathscr{D}(C)} D_{\mathrm{H},\varepsilon}(\rho_{BC} \|\mathrm{id}_B \otimes \sigma_C) \tag{431}
$$

$$
\geq \log \inf_{0 \leq M \leq I} \sup_{\sigma_C} \left\{ \left\| (\sqrt{M} \rho_{BC} \sqrt{M})^{\frac{1}{2}} \mathrm{id}_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2 : \text{Tr}(M \rho_{BC}) \geq 1 - \varepsilon \right\} \tag{432}
$$

$$
\geq \log \inf_{\substack{\tilde{\rho}_{BC} \in \mathcal{H}_+(BC) \\ \text{Tr}(\tilde{\rho}) \leq 1}} \sup_{\sigma_C \in \mathcal{D}(C)} \left\| \tilde{\rho}_{BC}^{\frac{1}{2}} \text{id}_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2 \tag{433}
$$

$$
=H_{\max}^{\sqrt{2\varepsilon}}(B|C)_{\rho}.
$$
\n(434)

⊓⊔

<span id="page-67-1"></span>**Lemma 54.** *Let*  $\rho \in \mathcal{D}(A)$  *and*  $\sigma \in \mathcal{H}_+(A)$ *.* 

$$
\left\| \left( \sqrt{M} \rho \sqrt{M} \right)^{\frac{1}{2}} \sigma^{\frac{1}{2}} \right\|_{1}^{2} \leq \text{Tr}(M\sigma). \tag{435}
$$

*Proof.* This fact is used in [\[DKF](#page-55-12)<sup>+</sup>14, Proposition 4.2]. It uses the semidefinite program for the fidelity  $\|\sqrt{\omega}\sqrt{\theta}\|_1^2 = \min\{\text{Tr}(Z\theta) : \omega_{AE} \le Z \otimes I_E, Z \ge 0\}$  [\[Wat09,](#page-58-18) Corollary 7], where  $\omega_{AE}$  is a purification of  $\omega$ . Let  $\rho_{AE}$  be a purification of  $\rho_A$ . Then we have  $\sqrt{M_A} \rho_{AE} \sqrt{M_A} \le M_A \otimes I_E$  and  $\sqrt{M_A} \rho_{AE} \sqrt{M_A}$ is a purification of  $\sqrt{M} \rho_A \sqrt{M}$ . As a result  $Z = M$  is feasible for the semidefinite program above and we get the desired result. □