

Variational expressions and Uhlmann theorem for measured divergences

Kun Fang ^{*1}, Hamza Fawzi ^{†2}, and Omar Fawzi ^{‡3}

¹School of Data Science, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, China

²Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, United Kingdom

³Univ Lyon, Inria, ENS Lyon, UCBL, LIP, Lyon, France

Abstract

We present a variational expression for measured f -divergences for convex functions f . Under a suitable operator convexity condition, this variational expression leads to a convex optimization problem recovering the known expression in the case of Rényi divergences. As an application, we establish an Uhlmann theorem for a wide class of measured f -divergences including measured α -Rényi divergences for all $\alpha \geq 0$. The well-known Uhlmann theorem for the fidelity corresponds to the special case $\alpha = \frac{1}{2}$.

1 Introduction

For a convex function $f : \mathbb{R}_{>0} \rightarrow \mathbb{R}$, the f -divergence between discrete probability vectors P and Q is defined as¹

$$S_f(P\|Q) := \sum_x Q(x) f\left(\frac{P(x)}{Q(x)}\right). \quad (1)$$

This family already appears in Rényi's work [Rén61] and was further studied by Csiszár [Csi67] and Ali and Silvey [AS66]. Important examples include the Kullback-Leibler divergence (also called relative entropy) $D(\cdot\|\cdot)$ corresponding to $f_1(t) = t \log t$ with $D(P\|Q) = S_{f_1}(P\|Q)$ and the α -Rényi divergences $D_\alpha(\cdot\|\cdot)$ corresponding to $f_\alpha(t) = \text{sign}(\alpha - 1)t^\alpha$ with $D_\alpha(P\|Q) = \frac{1}{\alpha-1} \log(\text{sign}(\alpha - 1)S_{f_\alpha}(P\|Q))$. Such divergences play an important role in various areas of statistics and information theory, we refer to [PW25, Chapter 7] for an overview.

For quantum states modeled by positive semidefinite operators ρ and σ , there are multiple ways of extending (1); see [HM17] for a systematic exposition. The main focus on this paper is on the *measured* [Don86, HP91] f -divergence which is obtained by performing a measurement M on the states ρ and σ and computing the f -divergence between the resulting probability distributions $P_{\rho,M}$ and $P_{\sigma,M}$. We then take the supremum over all possible measurements or all projective measurements:

$$S_{M,f}(\rho\|\sigma) := \sup_{M \text{ POVM}} S_f(P_{\rho,M}\|P_{\sigma,M}) \quad (2)$$

$$S_{M,f}^P(\rho\|\sigma) := \sup_{M \text{ PVM}} S_f(P_{\rho,M}\|P_{\sigma,M}), \quad (3)$$

^{*}kunfang@cuhk.edu.cn

[†]h.fawzi@damtp.cam.ac.uk

[‡]omar.fawzi@ens-lyon.fr

¹ We make the following standard conventions to define this expression: $f(0) = \lim_{t \downarrow 0} f(t)$, $0f(\frac{0}{0}) = 0$ and $0f(\frac{a}{0}) = \lim_{t \downarrow 0} tf(\frac{a}{t})$ for $a > 0$. Note that the limits may be $+\infty$.

where in (2) the supremum is over positive operator-valued measures (POVM) M described by a discrete set \mathcal{X} and positive semidefinite operators $\{M_x\}_{x \in \mathcal{X}}$ satisfying $\sum_{x \in \mathcal{X}} M_x = I$, the supremum in (3) is over projection-valued measures (PVM) M , i.e., POVMs satisfying in addition that M_x is an orthogonal projector for all $x \in \mathcal{X}$ and $P_{\rho, M}(x) = \text{Tr}[\rho M_x]$.

1.1 Main results

The objective of this paper is to understand some of the fundamental properties of both (2) and (3). First, concerning the data-processing inequality (DPI), i.e., the property $S(T(\rho)||T(\sigma)) \leq S(\rho||\sigma)$ for any completely positive and trace-preserving map T , it is clear that $S_{M, f}$ satisfies it for any convex f , but when does $S_{M, f}^P$ satisfy DPI? By [HM17, Corollary 4.20], this is equivalent to asking when $S_{M, f} = S_{M, f}^P$? Our first result (Theorem 2) shows that provided the Fenchel conjugate f^* of f is operator convex, we have $S_{M, f} = S_{M, f}^P$. The condition is in fact more general: it suffices to find a one-to-one function ψ such that $-\psi$ and $f^* \circ \psi$ are operator convex. In order to prove this, we follow the same strategy as [BFT17], namely we establish a variational expression for $S_{M, f}^P(\rho||\sigma)$ for any convex function f (Proposition 1). Then, we show that under the operator convexity condition, this variational expression can be written as a convex optimization program. Then, applying the operator Jensen inequality, we establish the desired equality $S_{M, f} = S_{M, f}^P$. Note that choosing $f = f_\alpha$, we recover the result for the measured Rényi divergences [BFT17] with a streamlined proof.

The second property we consider is the Uhlmann property that is well known for the fidelity [Uhl76]: given a quantum state ρ_A on A and a state σ_{AR} on $A \otimes R$, is there an extension ρ_{AR} of ρ_A such that $S_{M, f}(\rho_{AR}||\sigma_{AR}) = S_{M, f}(\rho_A||\sigma_A)$? We show in Theorem 3 that if f^* is operator convex and operator monotone and the domain of f^* is unbounded from below, then Uhlmann's property holds. Similarly, if $(f^*)^{-1}$ is operator concave and operator monotone and the range of f^* is unbounded from above, then Uhlmann's property holds with the roles of ρ and σ exchanged. In the special case of Rényi divergences, this shows the Uhlmann property for all $\alpha \geq 0$ (Theorem 4). This was previously known for $\alpha = \frac{1}{2}$ [Uhl76] and in the limit $\alpha \rightarrow \infty$ [MFSR24]. Our result can also be used to recover the regularized Uhlmann theorem of [MFSR24].

We conclude this paper in Section 4 with an intriguing “duality” property (Proposition 6) relating the measured Kullback-Leibler divergence $D_M(\rho||\mathcal{C})$ between ρ and a convex set \mathcal{C} and the Umegaki quantum relative entropy $D(\rho||\mathcal{C}_{++}^\circ)$ between ρ and the corresponding polar set \mathcal{C}_{++}° .

Notation: For a finite-dimensional Hilbert space A , we let $\mathcal{H}(A)$ denote the space of Hermitian operators on A , $\mathcal{H}_+(A) = \{\omega \in \mathcal{H}(A) : \omega \geq 0\}$ the set of positive semidefinite operators and $\mathcal{H}_{++}(A) = \{\omega \in \mathcal{H}(A) : \omega > 0\}$ the set of positive definite operators. The set of density operators is denoted by $\mathcal{D}(A) = \{\rho \in \mathcal{H}_+(A) : \text{Tr}[\rho] = 1\}$. We drop the Hilbert space from the notation when it is clear from the context. For $\omega \in \mathcal{H}$, $\text{spec}(\omega)$ denotes the spectrum of ω . The measured Rényi divergences for $\alpha \geq 0$ are defined as

$$D_{M, \alpha}(\rho||\sigma) = \frac{1}{\alpha - 1} \log(\text{sign}(\alpha - 1) S_{M, f_\alpha}(\rho||\sigma)) \quad \text{for } \alpha \neq 1 \quad (4)$$

$$D_M(\rho||\sigma) = S_{M, f_1}(\rho||\sigma) \quad \text{for } \alpha = 1. \quad (5)$$

In the limit $\alpha \rightarrow \infty$, it is known that $D_{M, \alpha} \rightarrow D_{\max}$, see e.g., [MO15, Appendix A].

Independent work: We note that in independent and concurrent work, Mazzola, Renner and Sutter [MSR25] established a regularized Uhlmann theorem for the Sandwiched Rényi divergence for $\alpha \geq \frac{1}{2}$ as well as an Uhlmann inequality for the measured Rényi divergences i.e., they show the existence of an extension satisfying $D_{M, \alpha}(\rho_{AR}||\sigma_{AR})$ is upper bounded by the corresponding Sandwiched Rényi divergence.

2 Variational expression for measured divergences

In the classical setting, variational expressions for f -divergences play an important role (see [PW25, Chapter 7]), but they are also fundamental in the quantum case in particular for the measured divergences as they are at the core of many applications of such divergences; see e.g., [BFT17, SBT17, FFF24].

In order to state our variational expression, we need to introduce the Fenchel conjugate of a function. For a function $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$, the Fenchel conjugate (also called Legendre transform) is defined as [BV04]

$$f^*(z) := \sup_{t \in \text{dom}(f)} \{tz - f(t)\}. \quad (6)$$

where $\text{dom}(f) = \{t \in \mathbb{R} : f(t) < +\infty\}$. The domain of f^* is defined in the same way as

$$\text{dom}(f^*) = \{z \in \mathbb{R} : f^*(z) < +\infty\}.$$

As f should define a divergence, we assume throughout this paper that $(0, \infty) \subseteq \text{dom}(f)$ and that f is convex. We assume in addition that f lower-semicontinuous. Note that the f -divergence only depends on the values f takes on $(0, \infty)$, but the Fenchel conjugate (and hence the variational expressions we obtain) depends on the the values of f on $\text{dom}(f)$.

We prove the following general variational formula for the projective measured f -divergence under the sole condition that f is convex and lower-semicontinuous. This can be seen as a quantum analog of [PW25, Theorem 7.26].

Proposition 1. *Let $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ be a convex lower-semicontinuous function and $(0, \infty) \subseteq \text{dom}(f)$. For any $\rho, \sigma \in \mathcal{H}_+$,*

$$S_{M,f}^P(\rho \parallel \sigma) = \sup_{\omega \in \mathcal{H}, \text{spec}(\omega) \subset \text{dom}(f^*)} \text{Tr}[\rho\omega] - \text{Tr}[\sigma f^*(\omega)], \quad (7)$$

where f^* is the Fenchel conjugate of f as defined in (6).

Proof. Since f is convex and lower-semicontinuous, we know that $f = (f^*)^*$ [Roc70, Theorem 12.2], i.e., for any $t \in \text{dom}(f)$

$$f(t) = \sup_{z \in \text{dom}(f^*)} \{tz - f^*(z)\}.$$

Hence for any $r, s > 0$ we get

$$sf(r/s) = \sup_{z \in \text{dom}(f^*)} \{rz - sf^*(z)\}.$$

Plugging into (3), we get

$$S_{M,f}^P(\rho \parallel \sigma) = \sup_{\{M_x\} \text{ PVM}} \sum_x \text{Tr}[M_x \sigma] f\left(\frac{\text{Tr}[M_x \rho]}{\text{Tr}[M_x \sigma]}\right) \quad (8)$$

$$= \sup_{\{M_x\} \text{ PVM}} \sum_x \sup_{z_x \in \text{dom}(f^*)} \text{Tr}[M_x \rho] z_x - \text{Tr}[M_x \sigma] f^*(z_x) \quad (9)$$

$$= \sup_{\{M_x\} \text{ PVM}} \sup_{\{z_x\}} \sum_x \text{Tr}[z_x M_x \rho] - \text{Tr}[f^*(z_x) M_x \sigma] \quad (10)$$

$$= \sup_{\{M_x\} \text{ PVM}} \sup_{\{z_x\}} \text{Tr}\left[\sum_x z_x M_x \rho\right] - \text{Tr}\left[f^*\left(\sum_x z_x M_x\right) \sigma\right] \quad (11)$$

$$= \sup_{\omega \in \mathcal{H}, \text{spec}(\omega) \subset \text{dom}(f^*)} \text{Tr}[\rho\omega] - \text{Tr}[\sigma f^*(\omega)], \quad (12)$$

$f(t)$	$\text{dom}(f)$	$f^*(z)$	$\text{dom}(f^*)$	Props. of f^* on $\text{dom}(f^*)$	Props. of $(f^*)^{-1}$ on $\text{range}(f^*)$
$-t^\alpha$ ($\alpha \in [0, 1)$)	$\mathbb{R}_{>0}$	$(1 - \alpha) \left(\frac{-z}{\alpha}\right)^{-\frac{\alpha}{1-\alpha}}$	$\mathbb{R}_{<0}$	$\alpha \in [0, \frac{1}{2}]$: op. convex and monotone	$\alpha \in [\frac{1}{2}, 1)$: op. concave and monotone
$\begin{cases} t^\alpha & t \geq 0 \\ 0 & t < 0 \end{cases}$ ($\alpha > 1$)	\mathbb{R}	$(\alpha - 1) \left(\frac{z}{\alpha}\right)^{\frac{\alpha}{\alpha-1}}$	$\mathbb{R}_{>0}$	$\alpha \geq 2$: op. convex	$\alpha > 1$: op. concave and monotone
$t \log(t)$	$\mathbb{R}_{>0}$	e^{z-1}	\mathbb{R}	Not applicable	Op. concave and monotone
$\frac{1}{2} t - 1 $	\mathbb{R}	z	$[-\frac{1}{2}, \frac{1}{2}]$	Linear monotone	Linear monotone

Table 1: Convex functions f and their Fenchel conjugates

where everywhere $\{M_x\}$ is a projective measurement and $z_x \in \text{dom}(f^*)$. Note that in the third line we used the fact that M_x are orthogonal projectors. \square

Table 1 shows the Fenchel conjugate of some common functions.

Remark 1 We can recover Rényi divergences for $\alpha \in [0, 1)$ by choosing

$$f_\alpha(t) = -t^\alpha, \quad \text{dom}(f_\alpha) = \mathbb{R}_{>0}$$

whose Fenchel conjugate is given in Table 1. In this case, the variational expression (7) becomes:

$$S_{M, f_\alpha}^P(\rho \parallel \sigma) = \sup_{\omega < 0} \text{Tr}[\rho\omega] - (1 - \alpha) \text{Tr} \left[\sigma \left(-\frac{\omega}{\alpha} \right)^{-\frac{\alpha}{1-\alpha}} \right],$$

which becomes after the change of variable $\gamma = -\frac{\omega}{\alpha}$:

$$S_{M, f_\alpha}^P(\rho \parallel \sigma) = \sup_{\gamma > 0} -\alpha \text{Tr}[\rho\gamma] - (1 - \alpha) \text{Tr} \left[\sigma \gamma^{-\frac{\alpha}{1-\alpha}} \right].$$

Then, since $D_{M, \alpha}(\rho \parallel \sigma) = \frac{1}{\alpha-1} \log \left(-S_{M, f_\alpha}^P(\rho \parallel \sigma) \right)$, we get the variational expression established in [BFT17] for $\alpha \in (0, 1)$ (up to a simple change of variable that is described in Remark 3). Note that for $\alpha = 0$, we interpret γ^0 as the projector on the support of γ .

When $\alpha > 1$, we consider the function²

$$f_\alpha(t) = \begin{cases} t^\alpha & t \geq 0 \\ 0 & t < 0, \end{cases} \quad \text{dom}(f_\alpha) = \mathbb{R} \quad (13)$$

whose Fenchel conjugate is given in Table 1. The expression (7) gives

$$S_{M, f_\alpha}^P(\rho \parallel \sigma) = \sup_{\omega > 0} \alpha \text{Tr}[\rho\omega] + (1 - \alpha) \text{Tr} \left[\sigma \omega^{\frac{\alpha}{1-\alpha}} \right],$$

which also matches with the expression found in [BFT17].

For the measured Kullback-Leibler divergence, we choose $f_1(t) = t \log t$ and $\text{dom}(f_1) = \mathbb{R}_{>0}$. This gives

$$S_{M, f_1}^P(\rho \parallel \sigma) = \sup_{\omega \in \mathcal{H}} \text{Tr}[\rho\omega] - \text{Tr} \left[\sigma e^{\omega-1} \right], \quad (14)$$

which again is equivalent to the expressions obtained in [Pet08, BFT17].

² The reason we consider the function (13) defined on the whole of \mathbb{R} instead of just $\mathbb{R}_{>0}$ is that the resulting Fenchel conjugate is defined on $\mathbb{R}_{>0}$ instead of the whole of \mathbb{R} which is useful in the application of Theorem 2

The variational expression (7) holds for all convex lower-semicontinuous functions f . However, the optimization program as written is in general not a convex one unless additional conditions are imposed, such as operator convexity of f^* . The theorem below gives a general sufficient condition under which $S_{M,f}^P(\rho\|\sigma)$ can be expressed as a convex program and which guarantees that both the measured divergence as well as the one restricted to projective measurements match.

Theorem 2. *Let $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ be a convex lower-semicontinuous function such that $(0, \infty) \subset \text{dom}(f)$, and let f^* be its Fenchel conjugate with domain $\text{dom}(f^*)$.*

Assume $\psi : J \rightarrow \text{dom}(f^)$ is a one-to-one map from an interval $J \subset \mathbb{R}$ to $\text{dom}(f^*)$ such that*

- ψ is operator concave on J
- $f^* \circ \psi$ is operator convex on J .

Then we have, for any $\rho, \sigma \in \mathcal{H}_+$

$$S_{M,f}(\rho\|\sigma) = S_{M,f}^P(\rho\|\sigma) = \sup_{\gamma \in \mathcal{H}, \text{spec}(\gamma) \subset J} \{ \text{Tr}[\rho\psi(\gamma)] - \text{Tr}[\sigma(f^* \circ \psi)(\gamma)] \}. \quad (15)$$

Remark 2 Combining Theorem 2 with [HM17, Corollary 4.20], it follows that under the conditions of the theorem, the data-processing inequality holds.

Remark 3 For all the examples mentioned in Remark 1, such a ψ can be found. In fact, for $\alpha \in (0, \frac{1}{2}]$, $f_\alpha^*(z) = (1 - \alpha)(\frac{-z}{\alpha})^{-\frac{\alpha}{1-\alpha}}$ is already operator convex so $\psi(z) = z$ works. For $\alpha \in [\frac{1}{2}, 1)$, we can choose $J = \mathbb{R}_{>0}$ and $\psi(\lambda) = (f_\alpha^*)^{-1}(\lambda) = -\alpha(\frac{\lambda}{1-\alpha})^{-\frac{1-\alpha}{\alpha}}$. For $\alpha > 1$, we can also choose $J = \mathbb{R}_{>0}$ and $\psi(\lambda) = (f_\alpha^*)^{-1}(\lambda) = \alpha(\frac{\lambda}{\alpha-1})^{\frac{\alpha-1}{\alpha}}$. Similarly, for $f(t) = t \log t$, we choose $J = \mathbb{R}_{>0}$ and $\psi(\lambda) = (f^*)^{-1}(\lambda) = 1 + \log \lambda$. For all these examples, the result (15) was already established in [BFT17].

Another example is the measured total variation distance which is obtained by taking $f_{TV}(t) = \frac{1}{2}|t - 1|$ and $\text{dom}(f) = \mathbb{R}$. With this choice $f_{TV}^*(z) = z$ for $z \in \text{dom}(f_{TV}^*) = [-\frac{1}{2}, \frac{1}{2}]$. Expression (15) then gives the well-known variational formulation of the trace distance.

Proof of Theorem 2. The second equality in (15) is a simple change of variables to (7), namely $\omega = \psi(\gamma)$. The inequality $S_{M,f}(\rho\|\sigma) \geq S_{M,f}^P(\rho\|\sigma)$ is clear and so it remains to show the reverse inequality. Fix $\{M_x\}$ any POVM and note that we can write, using the same argument as in (9)

$$\sum_x \text{Tr}[M_x\sigma] f\left(\frac{\text{Tr}[M_x\rho]}{\text{Tr}[M_x\sigma]}\right) = \sup_{\{z_x\} \subset \text{dom}(f^*)} \text{Tr}\left[\sum_x z_x M_x \rho\right] - \text{Tr}\left[\sum_x f^*(z_x) M_x \sigma\right].$$

Since ψ is a one-to-one map from J to $\text{dom}(f^*)$, we can do the change of variables $z_x = \psi(\lambda_x)$ where $\lambda_x \in J$, and so we get

$$\sum_x \text{Tr}[M_x\sigma] f\left(\frac{\text{Tr}[M_x\rho]}{\text{Tr}[M_x\sigma]}\right) = \sup_{\{\lambda_x\} \subset J} \text{Tr}\left[\sum_x \psi(\lambda_x) M_x \rho\right] - \text{Tr}\left[\sum_x f^*(\psi(\lambda_x)) M_x \sigma\right]. \quad (16)$$

By assumption, ψ is operator concave, and so using the operator Jensen inequality [HP03]

$$\psi\left(\sum_x \lambda_x M_x\right) = \psi\left(\sum_x \sqrt{M_x}(\lambda_x I)\sqrt{M_x}\right) \geq \sum_x \psi(\lambda_x) M_x$$

and similarly, since $f^* \circ \psi$ is operator convex we have

$$\sum_x f^*(\psi(\lambda_x))M_x \geq (f^* \circ \psi) \left(\sum_x \lambda_x M_x \right).$$

Combining the last two inequalities in (16) gives

$$\begin{aligned} \sum_x \text{Tr}[M_x \sigma] f \left(\frac{\text{Tr}[M_x \rho]}{\text{Tr}[M_x \sigma]} \right) &\leq \sup_{\{\lambda_x\} \subset J} \text{Tr} \left[\psi \left(\sum_x \lambda_x M_x \right) \rho \right] - \text{Tr} \left[f^* \circ \psi \left(\sum_x \lambda_x M_x \right) \sigma \right] \\ &\leq \sup_{\gamma \in \mathcal{H}, \text{spec}(\gamma) \subset J} \text{Tr}[\psi(\gamma)\rho] - \text{Tr}[(f^* \circ \psi)(\gamma)\sigma] \\ &= S_{M,f}^P(\rho \parallel \sigma) \end{aligned}$$

where in the second inequality above we used $\gamma = \sum_x \lambda_x M_x$ whose spectrum is in J since $M_x \geq 0$ and $\sum_x M_x = I$. \square

3 Uhlmann's theorem for measured divergences

Uhlmann's theorem is a fundamental result in quantum information theory that describes the relationship between the fidelity of mixed quantum states and their extensions [Uhl76]. More specifically, it states that for any two quantum states ρ_A and σ_{AR} , there exists an extension ρ_{AR} of ρ_A such that the fidelity $F(\rho_{AR}, \sigma_{AR})$ equals the fidelity $F(\rho_A, \sigma_A)$. The following result generalizes the renowned Uhlmann's theorem to measured f -divergences in general.

Theorem 3 (Uhlmann's theorem for measured f -divergences). *Let f be a convex lower-semicontinuous function. If f^* is operator convex and operator monotone on $\text{dom}(f^*)$ and $\inf\{z : z \in \text{dom}(f^*)\} = -\infty$, then for any $\rho_A \in \mathcal{D}(A)$ and $\sigma_{AR} \in \mathcal{H}_+(AR)$, we have*

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} S_{M,f}(\rho_{AR} \parallel \sigma_{AR}) = S_{M,f}(\rho_A \parallel \sigma_A). \quad (17)$$

Similarly, if $f^ : \text{dom}(f^*) \rightarrow \text{range}(f^*)$ is one-to-one and $(f^*)^{-1}$ is operator concave and operator monotone on $\text{range}(f^*)$ and $\sup\{\lambda : \lambda \in \text{range}(f^*)\} = +\infty$, then for any $\rho_{AR} \in \mathcal{D}(AR)$ and $\sigma_A \in \mathcal{H}_+(A)$, we have*

$$\inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} S_{M,f}(\rho_{AR} \parallel \sigma_{AR}) = S_{M,f}(\rho_A \parallel \sigma_A). \quad (18)$$

In addition, the infimum in both equations is attained.

Remark 4 We leave it as an open question whether such a result can be extended to other choices of f . However, some of the conditions are necessary. For example, the trace distance does not satisfy the Uhlmann property, but as mentioned in Remark 3, we have $f_{TV}^*(z) = z$ for $z \in \text{dom}(f^*) = [-\frac{1}{2}, +\frac{1}{2}]$ which is clearly operator convex and monotone. This means that the condition on $\text{dom}(f^*)$ is needed in this context.

Proof. We establish the result for Eq. (17) first. Recall the variational expression in Eq. (15) with $\psi = \text{id}$:

$$S_{M,f}(\rho \parallel \sigma) = \sup_{\substack{\gamma \in \mathcal{H} \\ \text{spec}(\gamma) \subset \text{dom}(f^*)}} \{\text{Tr}[\rho\gamma] - \text{Tr}[\sigma f^*(\gamma)]\}. \quad (19)$$

We now use Sion's minimax theorem [Sio58, Corollary 3.3] to get

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) = \sup_{\substack{\gamma_{AR} \in \mathcal{H}(AR) \\ \text{spec}(\gamma_{AR}) \subset \text{dom}(f^*)}} \inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} \{\text{Tr}[\rho_{AR} \gamma_{AR}] - \text{Tr}[\sigma_{AR} f^*(\gamma_{AR})]\}.$$

In fact, the set $\{\rho_{AR} \in \mathcal{H}_+(AR) : \text{Tr}_R \rho_{AR} = \rho_A\}$ is compact and convex and $\{\gamma_{AR} \in \mathcal{H}(AR) : \text{spec}(\gamma_{AR}) \subset \text{dom}(f^*)\}$ is convex. In addition the expression $\text{Tr}[\rho_{AR} \gamma_{AR}] - \text{Tr}[\sigma_{AR} f^*(\gamma_{AR})]$ is concave in γ (because f^* is operator convex) and linear in ρ . Now, by semidefinite programming duality, we have for a fixed $\gamma_{AR} \in \mathcal{H}(AR)$

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} \text{Tr}[\rho_{AR} \gamma_{AR}] = \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \Lambda_A \otimes I_R \leq \gamma_{AR}}} \text{Tr}[\rho_A \Lambda_A]. \quad (20)$$

In fact it is clear that the infimum is finite as $0 \leq \rho_{AR} \leq \text{Tr}[\rho_A] I_{AR}$ and $\Lambda_A = -(\|\gamma_{AR}\|_\infty + 1) I_A$ is strictly feasible for the dual. As a result,

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) = \sup_{\substack{\gamma_{AR} \in \mathcal{H}(AR) \\ \text{spec}(\gamma_{AR}) \subset \text{dom}(f^*)}} \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \Lambda_A \otimes I_R \leq \gamma_{AR}}} \{\text{Tr}[\rho_A \Lambda_A] - \text{Tr}[\sigma_{AR} f^*(\gamma_{AR})]\}. \quad (21)$$

Note that we can obtain a lower bound on the supremum (the easy direction) by restricting ourselves to $\gamma_{AR} = \Lambda_A \otimes I_R$ and observing that $f^*(\Lambda_A \otimes I_R) = f^*(\Lambda_A) \otimes I_R$:

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) \geq \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \text{spec}(\Lambda_A) \subset \text{dom}(f^*)}} \{\text{Tr}[\rho_A \Lambda_A] - \text{Tr}[\sigma_A f^*(\Lambda_A)]\}. \quad (22)$$

$$= S_{M,f}(\rho_A \| \sigma_A). \quad (23)$$

For the other direction, as f^* is operator monotone we have for any feasible (γ, Λ) , it holds that $f^*(\gamma_{AR}) \geq f^*(\Lambda_A) \otimes I_R$. This then implies

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) \leq \sup_{\substack{\gamma_{AR} \in \mathcal{H}(AR) \\ \text{spec}(\gamma_{AR}) \subset \text{dom}(f^*)}} \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \Lambda_A \otimes I_R \leq \gamma_{AR}}} \{\text{Tr}[\rho_A \Lambda_A] - \text{Tr}[\sigma_A f^*(\Lambda_A)]\} \quad (24)$$

$$\leq \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \text{spec}(\Lambda_A) \subset \text{dom}(f^*)}} \{\text{Tr}[\rho_A \Lambda_A] - \text{Tr}[\sigma_A f^*(\Lambda_A)]\}. \quad (25)$$

$$= S_{M,f}(\rho \| \sigma). \quad (26)$$

In the second inequality, we used the condition $\inf \text{dom}(f^*) = -\infty$ to get that $\Lambda_A \otimes I_R \leq \gamma_{AR}$ and $\text{spec}(\gamma_{AR}) \subset \text{dom}(f^*)$ implies $\text{spec}(\Lambda_A) \subset \text{dom}(f^*)$. This establishes (17).

To prove that the infimum is attained, for any $n \geq 1$, we can take $\rho_{AR}^{(n)} \in \mathcal{H}_+(AR)$ satisfying $\text{Tr}_R \rho_{AR}^{(n)} = \rho_A$ such that $0 \leq S_{M,f}(\rho_{AR}^{(n)} \| \sigma_{AR}) - S_{M,f}(\rho_A \| \sigma_A) \leq \frac{1}{n}$. As the set $\{\rho_{AR} \in \mathcal{H}_+(AR) : \text{Tr}_R \rho_{AR} = \rho_A\}$ is compact, we can find a converging subsequence $\lim_{i \rightarrow \infty} \rho_{AR}^{n_i} = \rho_{AR}^\infty$. By construction, $\text{Tr}_R \rho_{AR}^\infty = \rho_A$ so the data processing inequality (see Remark 2) gives $S_{M,f}(\rho_{AR}^\infty \| \sigma_{AR}) \geq S_{M,f}(\rho_A \| \sigma_A)$. As $S_{M,f}$ can be written as the supremum of lower-semicontinuous functions, it is also lower semi-continuous. This implies

$$S_{M,f}(\rho_A \| \sigma_A) = \lim_{i \rightarrow \infty} S_{M,f}(\rho_{AR}^{n_i} \| \sigma_{AR}) \geq S_{M,f}(\rho_{AR}^\infty \| \sigma_{AR}). \quad (27)$$

So the infimum is attained at ρ_{AR}^∞ .

The proof of Eq. (18) is similar. Recall the variational expression in Eq. (15) with $\psi = (f^*)^{-1}$:

$$S_{M,f}(\rho \| \sigma) = \sup_{\substack{\gamma \in \mathcal{H} \\ \text{spec}(\gamma) \subset \text{range}(f^*)}} \{\text{Tr}[\rho (f^*)^{-1}(\gamma)] - \text{Tr}[\sigma \gamma]\}. \quad (28)$$

We now use Sion's minimax theorem [Sio58, Corollary 3.3] to get

$$\inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) = \sup_{\substack{\gamma_{AR} \in \mathcal{H}(AR) \\ \text{spec}(\gamma_{AR}) \subset \text{range}(f^*)}} \inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} \{ \text{Tr}[\rho_{AR}(f^*)^{-1}(\gamma_{AR})] - \text{Tr}[\sigma_{AR}\gamma_{AR}] \}. \quad (29)$$

In fact, the set $\{\sigma_{AR} \in \mathcal{H}_+(AR) : \text{Tr}_R \sigma_{AR} = \sigma_A\}$ is compact and convex and $\{\gamma_{AR} \in \mathcal{H}(AR) : \text{spec}(\gamma_{AR}) \subset \text{range}(f^*)\}$ is convex. In addition the expression $\text{Tr}[\rho_{AR}(f^*)^{-1}(\gamma_{AR})] - \text{Tr}[\sigma_{AR}\gamma_{AR}]$ is concave in γ and linear in σ . Now, by semidefinite programming duality, we have

$$\sup_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} \text{Tr}[\sigma_{AR}\gamma_{AR}] = \inf_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \Lambda_A \otimes I_R \geq \gamma_{AR}}} \text{Tr}[\sigma_A \Lambda_A]. \quad (30)$$

In fact it is clear that the supremum is finite as $0 \leq \sigma_{AR} \leq \text{Tr}[\sigma_A]I_{AR}$ and $\Lambda_A = (\|\gamma_{AR}\|_\infty + 1)I_A$ is strictly feasible for the dual. As a result,

$$\inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) = \sup_{\substack{\gamma_{AR} \in \mathcal{H}(AR) \\ \text{spec}(\gamma_{AR}) \subset \text{range}(f^*)}} \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \Lambda_A \otimes I_R \geq \gamma_{AR}}} \{ \text{Tr}[\rho_{AR}(f^*)^{-1}(\gamma_{AR})] - \text{Tr}(\sigma_A \Lambda_A) \}. \quad (31)$$

Note that we can in particular choose $\gamma_{AR} = \Lambda_A \otimes I_R$ and get the easy direction:

$$\inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) \geq \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \text{spec}(\Lambda_A) \subset \text{range}(f^*)}} \{ \text{Tr}[\rho_A(f^*)^{-1}(\Lambda_A)] - \text{Tr}[\sigma_A \Lambda_A] \}. \quad (32)$$

$$= S_{M,f}(\rho \| \sigma). \quad (33)$$

For the other direction, as $(f^*)^{-1}$ is operator monotone we have for any feasible (γ, Λ) , it holds that $(f^*)^{-1}(\gamma_{AR}) \leq (f^*)^{-1}(\Lambda_A) \otimes I_R$. In addition, as $\sup \text{range}(f^*) = +\infty$ and $\text{spec}(\gamma_{AR}) \subset \text{range}(f^*)$, we have that $\text{spec}(\Lambda_A) \subset \text{range}(f^*)$. In fact, f^* is continuous on its domain since it is a univariate lower-semicontinuous convex function, and thus $\text{range}(f^*)$ is an interval, which by our additional assumption, is of the form $[a, +\infty)$ or $(a, +\infty)$. This then implies

$$\inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} S_{M,f}(\rho_{AR} \| \sigma_{AR}) \leq \sup_{\substack{\Lambda_A \in \mathcal{H}(A) \\ \text{spec}(\Lambda_A) \subset \text{range}(f^*)}} \{ \text{Tr}[\rho_A(f^*)^{-1}(\Lambda_A)] - \text{Tr}[\sigma_A \Lambda_A] \}. \quad (34)$$

$$= S_{M,f}(\rho_A \| \sigma_A). \quad (35)$$

This establishes (18). The attainment of the infimum follows the same proof as before. \square

As an immediate corollary, we obtain the following result of Rényi divergences.

Corollary 4 (Uhlmann's theorem for measured Rényi divergences). *Let $\rho_A \in \mathcal{D}(A)$, $\sigma_{AR} \in \mathcal{H}_+(AR)$, and $\alpha \in [0, 1/2]$, then*

$$\inf_{\substack{\rho_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \rho_{AR} = \rho_A}} D_{M,\alpha}(\rho_{AR} \| \sigma_{AR}) = D_{M,\alpha}(\rho_A \| \sigma_A). \quad (36)$$

Similarly, let $\rho_{AR} \in \mathcal{D}(AR)$, $\sigma_A \in \mathcal{H}_+(A)$ and $\alpha \in [\frac{1}{2}, +\infty]$ ³, then

$$\inf_{\substack{\sigma_{AR} \in \mathcal{H}_+(AR) \\ \text{Tr}_R \sigma_{AR} = \sigma_A}} D_{M,\alpha}(\rho_{AR} \| \sigma_{AR}) = D_{M,\alpha}(\rho_A \| \sigma_A). \quad (37)$$

In addition, the infimum in both equations is attained.

Remark 5 Note that for $\alpha \geq 1$, it is easy to see that Uhlmann's property in the first argument (i.e., (36)) cannot hold in general. In fact, we can choose $\rho_A = |0\rangle\langle 0|$ and $\sigma_{AR} = |\Phi\rangle\langle\Phi|$ where $|\Phi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$, then $D_{M,\alpha}(\rho_A\|\sigma_A) = \log 2$ but $D_{M,\alpha}(\rho_{AR}\|\sigma_{AR}) = +\infty$ for any extension ρ_{AR} of ρ_A .

Proof. For $\alpha \in [0, \frac{1}{2}]$, as previously mentioned, we have $f_\alpha^*(z) = (1-\alpha)(\frac{-z}{\alpha})^{-\frac{\alpha}{1-\alpha}}$ and $\text{dom}(f^*) = \mathbb{R}_{<0}$. In this case, f_α^* is operator convex, operator monotone and the domain is unbounded from below.

For $\alpha \in [\frac{1}{2}, 1)$, we also have $f_\alpha^*(z) = (1-\alpha)(\frac{-z}{\alpha})^{-\frac{\alpha}{1-\alpha}}$ and $\text{dom}(f_\alpha^*) = \mathbb{R}_{<0}$. But in this case, we check the conditions for (18): we have f_α^* is one-to-one with $\text{range}(f_\alpha^*) = \mathbb{R}_{>0}$ and $(f_\alpha^*)^{-1}(\lambda) = -\alpha(\frac{\lambda}{1-\alpha})^{-\frac{1-\alpha}{\alpha}}$ which is operator concave, operator monotone and we have $\text{sup range}(f_\alpha^*) = +\infty$.

For $\alpha > 1$, we have $f_\alpha^*(z) = (\alpha-1)(\frac{z}{\alpha})^{\frac{\alpha}{\alpha-1}}$ with $\text{dom}(f^*) = \text{range}(f^*) = \mathbb{R}_{>0}$. Thus, $(f_\alpha^*)^{-1}(\lambda) = \alpha(\frac{\lambda}{\alpha-1})^{\frac{\alpha-1}{\alpha}}$ is operator concave, operator monotone and $\text{sup range}(f_\alpha^*) = +\infty$.

For $\alpha = 1$, recall that $f_1(t) = t \log t$, and thus we have $f_1^*(z) = e^{z-1}$ with $\text{dom}(f_1^*) = \mathbb{R}$ and $\text{range}(f_1^*) = \mathbb{R}_{>0}$. Thus $(f_1^*)^{-1}(\lambda) = 1 + \log \lambda$ is operator concave, operator monotone and $\text{sup range}(f_1^*) = +\infty$.

For $\alpha = +\infty$, we take supremum over $\alpha \in [1, +\infty)$. Note that $D_{M,\alpha}(\rho\|\sigma)$ is lower semicontinuous in (ρ, σ) for every $\alpha \in [1, +\infty)$ [MH23, Proposition III.11] and monotonic increasing in α for every (ρ, σ) . So we can use the minimax theorem in [MH11, Corollary A.2] to exchange the supremum and infimum. Finally, noting that $\sup_{\alpha \geq 1} D_{M,\alpha}(\rho\|\sigma) = D_{\max}(\rho\|\sigma)$, we have the desired result. \square

The case $\alpha = \frac{1}{2}$ corresponds to the fidelity as $D_{M,\frac{1}{2}} = -2 \log F$, where F is the fidelity; see [Fuc96, MO15]. Thus, Uhlmann's theorem is the special case $\alpha = \frac{1}{2}$ of this theorem. The case $\alpha = +\infty$ was shown in [MFSR24]. Moreover, the regularized Uhlmann's theorem from [MFSR24] follows easily from this theorem. Note that the result in [MFSR24] is shown for $\alpha > 1$, whereas the following result applies more generally.

Let $D_{S,\alpha}(\rho\|\sigma) := \frac{1}{\alpha-1} \log \text{Tr} \left[\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}} \right]^\alpha$ be the Sandwiched Rényi divergence [MLDS⁺13, WWY14].

Corollary 5. Let $\rho_{AR} \in \mathcal{D}(AR)$, $\sigma_A \in \mathcal{H}_+(A)$ and $\alpha \in [1/2, +\infty]$, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \inf_{\substack{\sigma_{A^n R^n} \in \mathcal{H}_+(A^n R^n) \\ \text{Tr}_{R^n} \sigma_{A^n R^n} = \sigma_A^{\otimes n}}} D_{S,\alpha}(\rho_{AR}^{\otimes n} \|\sigma_{A^n R^n}) = D_{S,\alpha}(\rho_A \|\sigma_A). \quad (38)$$

Proof. Applying Corollary 4 to the states $\rho_{AR}^{\otimes n}$ and $\sigma_A^{\otimes n}$, we get

$$\inf_{\substack{\sigma_{A^n R^n} \in \mathcal{H}_+(A^n R^n) \\ \text{Tr}_{R^n} \sigma_{A^n R^n} = \sigma_A^{\otimes n}}} D_{M,\alpha}(\rho_{AR}^{\otimes n} \|\sigma_{A^n R^n}) = D_{M,\alpha}(\rho_A^{\otimes n} \|\sigma_A^{\otimes n}). \quad (39)$$

By the asymptotic equivalence of the measured Rényi divergence and the Sandwiched Rényi divergence [FFF24, Lemma 29] applied with $\mathcal{B}_n = \{\sigma_{A^n R^n} \in \mathcal{H}_+(A^n R^n) : \text{Tr}_{R^n} \sigma_{A^n R^n} = \sigma_A^{\otimes n}\}$ (which is convex, compact and permutation invariant) gives

$$\lim_{n \rightarrow \infty} \frac{1}{n} D_{S,\alpha}(\rho_{AR}^{\otimes n} \|\mathcal{B}_n) = \lim_{n \rightarrow \infty} \frac{1}{n} D_{M,\alpha}(\rho_{AR}^{\otimes n} \|\mathcal{B}_n) = D_{S,\alpha}(\rho_A \|\sigma_A). \quad (40)$$

\square

³ When $\alpha = 1$, the equality is interpreted as D_M . When $\alpha = +\infty$, the equality is interpreted as D_{\max} .

Remark 6 Note that the set \mathcal{B}_n defined in the proof above satisfies all assumptions in [FFF24, Assumption 25]. In fact, (A.1) and (A.2) are clear and $\text{Tr}_{R_1 R_2}(\sigma_{A_1 R_1}^{(1)} \otimes \sigma_{A_2 R_2}^{(2)}) = \sigma_{A_1}^{(1)} \otimes \sigma_{A_2}^{(2)}$ so (A.3) is satisfied. The condition (A.4) states that the support function is submultiplicative, i.e., $h_{\mathcal{B}_{m+k}}(X_{A^m R^m} \otimes X_{A^k R^k}) \leq h_{\mathcal{B}_m}(X_{A^m R^m})h_{\mathcal{B}_k}(X_{A^k R^k})$. For that, recall that we have obtained a dual formulation for the support function of the sets \mathcal{B}_n in (30). We use this same dual formulation to establish the desired submultiplicativity:

$$\begin{aligned}
h_{\mathcal{B}_{m+k}}(X_{A^m R^m} \otimes X_{A^k R^k}) &= \sup_{\substack{\sigma_{A^{m+k} R^{m+k}} \in \mathcal{H}_+(A^{m+k} R^{m+k}) \\ \text{Tr}_{R^{m+k}} \sigma_{A^{m+k} R^{m+k}} = \sigma_A^{\otimes m+k}}} \text{Tr}(\sigma_{A^{m+k} R^{m+k}} X_{A^m R^m} \otimes X_{A^k R^k}) \\
&= \inf_{\substack{\Lambda_{A^{m+k}} \in \mathcal{H}(A^{m+k}) \\ \Lambda_{A^{m+k}} \otimes I_{R^{m+k}} \geq X_{A^m R^m} \otimes X_{A^k R^k}}} \text{Tr}(\sigma_A^{\otimes m+k} \Lambda_{A^{m+k}}) \\
&\leq \inf_{\substack{\Lambda_{A^m} \in \mathcal{H}(A^m), \Lambda_{A^k} \in \mathcal{H}(A^k) \\ \Lambda_{A^m} \otimes \Lambda_{A^k} \otimes I_{R^{m+k}} \geq X_{A^m R^m} \otimes X_{A^k R^k}}} \text{Tr}(\sigma_A^{\otimes m+k} \Lambda_{A^m} \otimes \Lambda_{A^k}) \\
&\leq \inf_{\substack{\Lambda_{A^m} \in \mathcal{H}(A^m), \Lambda_{A^k} \in \mathcal{H}(A^k) \\ \Lambda_{A^m} \otimes I_{R^m} \geq X_{A^m R^m} \\ \Lambda_{A^k} \otimes I_{R^k} \geq X_{A^k R^k}}} \text{Tr}(\sigma_A^{\otimes m} \Lambda_{A^m}) \text{Tr}(\sigma_A^{\otimes k} \Lambda_{A^k}) \\
&= h_{\mathcal{B}_m}(X_{A^m R^m})h_{\mathcal{B}_k}(X_{A^k R^k}).
\end{aligned}$$

Thus, the results of [FFF24] characterize the asymmetric hypothesis testing problem between $\{\rho_{AR}^{\otimes n}\}$ and \mathcal{B}_n . A related hypothesis testing scenario was considered in [HT16]. The authors of [HT16] studied hypothesis testing between $\{\rho_{AR}^{\otimes n}\}$ and $\{\rho_A^{\otimes n} \otimes \sigma_{R^n} : \sigma \in \mathcal{D}(R^n)\}$ and demonstrated that the Rényi mutual information is the strong converse exponent in this context. Similarly, we expect Corollary 5 can be used to show establish the strong converse exponent for hypothesis testing between $\rho_{AR}^{\otimes n}$ and \mathcal{B}_n .

4 Duality with the quantum relative entropy

This section is specific to the measured Kullback-Leibler divergence (i.e., $f(t) = t \log t$) also called the measured relative entropy: we utilize the variational expression for D_M to derive a duality relation with the Umegaki quantum relative entropy which is defined by $D(\rho\|\sigma) = \text{Tr}(\rho \log \rho) - \text{Tr}(\rho \log \sigma)$ when the support of ρ is included in the support of σ and $+\infty$ otherwise. For a quantum divergence \mathbb{D} and a set \mathcal{C} of positive operators, we write $\mathbb{D}(\rho\|\mathcal{C}) := \inf_{\sigma \in \mathcal{C}} \mathbb{D}(\rho\|\sigma)$.

Proposition 6. *Let $\rho \in \mathcal{D}$ and $\mathcal{C} \subseteq \mathcal{H}_+$ be a compact convex set, it holds*

$$D(\rho\|\mathcal{C}) + D_M(\rho\|\mathcal{C}_{++}^\circ) = D(\rho\|I), \quad (41)$$

where $\mathcal{C}^\circ := \{X : \text{Tr}[XY] \leq 1, \forall Y \in \mathcal{C}\}$ is the polar set and $\mathcal{C}_{++}^\circ := \mathcal{C}^\circ \cap \mathcal{H}_{++}$.

Proof. By the variational expression for the measured Kullback-Leibler divergence [Pet08, BFT17] also presented in (14), we have

$$D_M(\rho\|\sigma) = \sup_{\omega \in \mathcal{H}_{++}} \text{Tr}[\rho \log \omega] + 1 - \text{Tr}[\sigma \omega] \quad (42)$$

$$= \sup_{\omega \in \mathcal{H}_{++}} -D(\rho\|\omega) - \text{Tr}[\sigma \omega] + 1 + D(\rho\|I) \quad (43)$$

$$= g_\rho^*(-\sigma) + 1 + D(\rho\|I), \quad (44)$$

where the second line follows from the definition of Umegaki relative entropy and the third line follows by denoting g_ρ^* as the Fenchel conjugate of $g_\rho : \omega \mapsto D(\rho\|\omega)$, with $\text{dom}(g_\rho) = \{\omega \geq 0 : \rho \ll \omega\}$, i.e., for $\tau \in \mathcal{H}$, $g_\rho^*(\tau) = \sup_{\omega > 0} \text{Tr}[\omega\tau] - D(\rho\|\omega)$.⁴ As g_ρ is lower-semicontinuous, the biduality relation of Fenchel conjugate [Roc70, Theorem 12.2] gives

$$D(\rho\|\sigma) = g_\rho(\sigma) = \sup_{\omega \in \text{dom}(g_\rho^*)} \text{Tr}[\sigma\omega] - g_\rho^*(\omega) \quad (45)$$

$$= \sup_{\omega < 0} \text{Tr}[\sigma\omega] - g_\rho^*(\omega) \quad (46)$$

$$= \sup_{\omega < 0} \text{Tr}[\sigma\omega] - D_{\text{M}}(\rho\|-\omega) + D(\rho\|I) + 1 \quad (47)$$

$$= \sup_{\omega > 0} -\text{Tr}[\sigma\omega] - D_{\text{M}}(\rho\|\omega) + D(\rho\|I) + 1. \quad (48)$$

In the second line we used the fact that for any $\omega \in \text{dom}(g_\rho^*)$

$$g_\rho^*(\omega) = \lim_{\epsilon \downarrow 0} g_\rho^*((1-\epsilon)\omega + \epsilon(-I))$$

which follows by applying [Roc70, Theorem 7.5]. Note that the theorem applies because $-I$ is in the interior of $\text{dom}(g_\rho^*)$ and g_ρ^* is lower-semicontinuous. The third line (47) follows from Eq. (44), and the last line follows by replacing ω with $-\omega$. Optimizing over $\sigma \in \mathcal{C}$, we have

$$D(\rho\|\mathcal{C}) = \inf_{\sigma \in \mathcal{C}} D(\rho\|\sigma) = \inf_{\sigma \in \mathcal{C}} \sup_{\omega \in \mathcal{H}_{++}} -\text{Tr}[\sigma\omega] - D_{\text{M}}(\rho\|\omega) + 1 + D(\rho\|I). \quad (49)$$

Since $D_{\text{M}}(\rho\|\omega)$ is convex in ω , the above objective function is concave in ω and convex in σ . Moreover, since \mathcal{C} is a compact convex set by assumption, we can use Sion's minimax theorem [Sio58, Corollary 3.3] to exchange the infimum and supremum. This gives

$$D(\rho\|\mathcal{C}) = \sup_{\omega \in \mathcal{H}_{++}} \inf_{\sigma \in \mathcal{C}} -\text{Tr}[\sigma\omega] - D_{\text{M}}(\rho\|\omega) + 1 + D(\rho\|I). \quad (50)$$

Let $h_{\mathcal{C}}(\omega) := \sup_{\sigma \in \mathcal{C}} \text{Tr}[\sigma\omega]$ be the support function of \mathcal{C} . We get

$$D(\rho\|\mathcal{C}) = \sup_{\omega \in \mathcal{H}_{++}} -h_{\mathcal{C}}(\omega) - D_{\text{M}}(\rho\|\omega) + 1 + D(\rho\|I) \quad (51)$$

$$\stackrel{(a)}{=} \sup_{\substack{\omega \in \mathcal{H}_{++} \\ h_{\mathcal{C}}(\omega)=1}} -D_{\text{M}}(\rho\|\omega) + D(\rho\|I) \quad (52)$$

$$\stackrel{(b)}{=} \sup_{\substack{\omega \in \mathcal{H}_{++} \\ h_{\mathcal{C}}(\omega) \leq 1}} -D_{\text{M}}(\rho\|\omega) + D(\rho\|I) \quad (53)$$

$$\stackrel{(c)}{=} -D_{\text{M}}(\rho\|\mathcal{C}_{++}^\circ) + D(\rho\|I), \quad (54)$$

To see why (a) holds, let ω be a solution in the first line and define $\tilde{\omega} = \omega/h_{\mathcal{C}}(\omega)$. Then $h_{\mathcal{C}}(\tilde{\omega}) = 1$ and we will see that $\tilde{\omega}$ achieves an objective value no smaller than ω as

$$-D_{\text{M}}(\rho\|\tilde{\omega}) = -D_{\text{M}}(\rho\|\omega) - \log h_{\mathcal{C}}(\omega) \geq -D_{\text{M}}(\rho\|\omega) - h_{\mathcal{C}}(\omega) + 1 \quad (55)$$

where the inequality follows from the fact that $\log x \leq x - 1$. The step (b) follows by the same argument. Suppose ω is a solution in the third line, let $\tilde{\omega} = \omega/h_{\mathcal{C}}(\omega)$ and we can check that $\tilde{\omega}$ gives an objective value no smaller than the second line. Finally, we have the step (c) by the fact that $h_{\mathcal{C}}(\omega) \leq 1$ if and only if $\omega \in \mathcal{C}^\circ$ and the notation that $\mathcal{C}_{++}^\circ = \mathcal{C}^\circ \cap \mathcal{H}_{++}$. This completes the proof. \square

⁴ Note that $g_\rho^*(\tau) := \sup_{\omega \in \text{dom}(g_\rho)} \text{Tr}[\omega\tau] - D(\rho\|\omega) = \sup_{\omega > 0} \text{Tr}[\omega\tau] - D(\rho\|\omega)$ since for any $\omega \geq 0$, $D(\rho\|\omega) = \lim_{\epsilon \downarrow 0} D(\rho\|\omega + \epsilon I)$

Remark 7 This duality result may assist in proving the subadditivity or superadditivity properties of one term on the left-hand side by establishing the corresponding property for the other term. For instance, a concrete example is as follows: if \mathcal{C} is closed under the tensor product, then it is easy to verify that $D(\rho\|\mathcal{C})$ is subadditive. This, in turn, implies that $D_{\text{M}}(\rho\|\mathcal{C}_{++}^{\circ})$ is superadditive. This reasoning also applies when exchanging the roles of D and D_{M} . If we can demonstrate the superadditivity of $D_{\text{M}}(\rho\|\mathcal{C}_{++}^{\circ})$, then the duality implies the subadditivity of $D(\rho\|\mathcal{C})$, which may be difficult to prove directly in some cases.

Since the measured relative entropy coincides with the quantum relative entropy in the classical case, we can replace D_{M} with D , obtaining an equality expressed solely in terms of the relative entropy. In the quantum case, if we consider the regularized divergence $\mathbb{D}^{\infty}(\rho\|\mathcal{C}) := \lim_{n \rightarrow \infty} \mathbb{D}(\rho^{\otimes n}\|\mathcal{C}_n)$, we get a self-duality for the quantum relative entropy as follows.

Corollary 7. *Let $\rho \in \mathcal{D}$. Let $\{\mathcal{C}_n\}_{n \in \mathbb{N}}$ be a sequences of sets satisfying $\mathcal{C}_n \subseteq \mathcal{H}_+(\mathcal{H}^{\otimes n})$ and $D_{\text{max}}(\rho^{\otimes n}\|\mathcal{C}_n) \leq cn$, for all $n \in \mathbb{N}$ and a constant $c \in \mathbb{R}_+$. In addition, each \mathcal{C}_n is convex, compact and permutation-invariant, and $\mathcal{C}_m \otimes \mathcal{C}_k \subseteq \mathcal{C}_{m+k}$ for any $m, k \in \mathbb{N}$. Then it holds*

$$D^{\infty}(\rho\|\mathcal{C}) + D^{\infty}(\rho\|\mathcal{C}_{++}^{\circ}) = D(\rho\|I). \quad (56)$$

Proof. By Proposition 6, we know that

$$\frac{1}{n}D(\rho^{\otimes n}\|\mathcal{C}_n) + \frac{1}{n}D_{\text{M}}(\rho^{\otimes n}\|(\mathcal{C}_n)_{++}^{\circ}) = D(\rho\|I). \quad (57)$$

Since the first term on the left-hand side is non-increasing in n (see [FFF24, Lemma 27]), its limit exists when n goes to infinity. By the equality relation, the limit for the second term on the left-hand side also exists. Therefore, we have

$$\left[\lim_{n \rightarrow \infty} \frac{1}{n}D(\rho^{\otimes n}\|\mathcal{C}_n) \right] + \left[\lim_{n \rightarrow \infty} \frac{1}{n}D_{\text{M}}(\rho^{\otimes n}\|(\mathcal{C}_n)_{++}^{\circ}) \right] = D(\rho\|I). \quad (58)$$

Finally, we complete the proof by noting that the second term on the left hand side is equal to the one with quantum relative entropy (see [FFF24, Lemma 29]). \square

Acknowledgements. We would like to thank David Sutter for a comment that helped us clarify Corollary 5 and Remark 6. K.F. is supported by the National Natural Science Foundation of China (grant No. 12404569 and 92470113), the Shenzhen Science and Technology Program (grant No. JCYJ20240813113519025) and the University Development Fund (grant No. UDF01003565) from the Chinese University of Hong Kong, Shenzhen. H.F. was partially funded by UK Research and Innovation (UKRI) under the UK government’s Horizon Europe funding guarantee EP/X032051/1. O.F. acknowledges support by the European Research Council (ERC Grant AlgoQIP, Agreement No. 851716), by the European Union’s Horizon 2020 research and innovation programme under Grant Agreement No 101017733 (VERIQ-TAS) and by the Agence Nationale de la Recherche under the Plan France 2030 with the reference ANR-22-PETQ-0009.

References

- [AS66] S. M. Ali and S. D. Silvey. A general class of coefficients of divergence of one distribution from another. *Journal of the Royal Statistical Society: Series B (Methodological)*, 28(1):131–142, 1966.

- [BFT17] M. Berta, O. Fawzi, and M. Tomamichel. On variational expressions for quantum relative entropies. *Letters in Mathematical Physics*, 107(12):2239–2265, 2017.
- [BV04] S. Boyd and L. Vandenberghe. *Convex optimization*. Cambridge University Press, 2004.
- [Csi67] I. Csiszár. On information-type measure of difference of probability distributions and indirect observations. *Studia Sci. Math. Hungar.*, 2:299–318, 1967.
- [Don86] M. J. Donald. On the relative entropy. *Communications in Mathematical Physics*, 105:13–34, 1986. <https://doi.org/10.1007/BF01212339>.
- [FFF24] K. Fang, H. Fawzi, and O. Fawzi. Generalized quantum asymptotic equipartition. *arXiv preprint arXiv:2411.04035*, 2024.
- [Fuc96] C. A. Fuchs. Distinguishability and accessible information in quantum theory. *arXiv preprint quant-ph/9601020*, 1996.
- [HM17] F. Hiai and M. Mosonyi. Different quantum f -divergences and the reversibility of quantum operations. *Reviews in Mathematical Physics*, 29(07):1750023, 2017. <https://doi.org/10.1142/S0129055X17500234>.
- [HP91] F. Hiai and D. Petz. The proper formula for relative entropy and its asymptotics in quantum probability. *Communications in Mathematical Physics*, 143:99–114, 1991. <https://doi.org/10.1007/BF02100287>.
- [HP03] F. Hansen and G. K. Pedersen. Jensen’s operator inequality. *Bulletin of the London Mathematical Society*, 35(4):553–564, 2003.
- [HT16] M. Hayashi and M. Tomamichel. Correlation detection and an operational interpretation of the Rényi mutual information. *Journal of Mathematical Physics*, 57(10), 2016.
- [MFSR24] T. Metger, O. Fawzi, D. Sutter, and R. Renner. Generalised entropy accumulation. *Communications in Mathematical Physics*, 405(11):261, 2024.
- [MH11] M. Mosonyi and F. Hiai. On the quantum Rényi relative entropies and related capacity formulas. *IEEE Transactions on Information Theory*, 57(4):2474–2487, 2011.
- [MH23] M. Mosonyi and F. Hiai. Some continuity properties of quantum Rényi divergences. *arXiv preprint arXiv: 2209.00646*, 2023.
- [MLDS⁺13] M. Müller-Lennert, F. Dupuis, O. Szehr, S. Fehr, and M. Tomamichel. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*, 54(12), 2013.
- [MO15] M. Mosonyi and T. Ogawa. Quantum hypothesis testing and the operational interpretation of the quantum rényi relative entropies. *Communications in Mathematical Physics*, 334:1617–1648, 2015.
- [MSR25] G. Mazzola, D. Sutter, and R. Renner. Uhlmann’s theorem for relative entropies. *arXiv preprint arXiv:2502.01749*, 2025.
- [Pet08] D. Petz. *Quantum Information Theory and Quantum Statistics*. Springer, 2008.
- [PW25] Y. Polyanskiy and Y. Wu. *Information theory: From coding to learning*. Cambridge university press, 2025.

- [Rén61] A. Rényi. On measures of entropy and information. In *Proceedings of the fourth Berkeley symposium on mathematical statistics and probability, volume 1: contributions to the theory of statistics*, volume 4, pages 547–562. University of California Press, 1961.
- [Roc70] R. T. Rockafellar. *Convex analysis*, 1970.
- [SBT17] D. Sutter, M. Berta, and M. Tomamichel. Multivariate trace inequalities. *Communications in Mathematical Physics*, 352:37–58, 2017.
- [Sio58] M. Sion. On general minimax theorems. *Pacific Journal of Mathematics*, 8(1):171–176, mar 1958.
- [Uhl76] A. Uhlmann. The “transition probability” in the state space of a*-algebra. *Reports on Mathematical Physics*, 9(2):273–279, 1976.
- [WWY14] M. M. Wilde, A. Winter, and D. Yang. Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy. *Communications in Mathematical Physics*, 331:593–622, 2014. <https://doi.org/10.1007/s00220-014-2122-x>.